Areas of Interest and Soil Boring L	ocations	San	nple Infor	mation				General	Chemical Rad C	Parameters hem ^(e)	GRO &	Analysis	Sample Type	
Soil Boring Easting Ore Processing Area	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	<u>На</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	Additional Comments
Hazardous Waste Storage Building														
	392437.85	SB-01 (0-1)	0-1	8/13/2010	1151	X	Χ	Χ					INVE	
		SB-01 (1-2)		8/13/2010		X	Χ	Χ	Χ	Χ			INVE	
		SB-01 (4-5)	4-5	8/13/2010	1154	Χ	Χ	Χ					INVE	
		SB-01 (9-10)	9-10	8/13/2010	1156							Hold	INVE	
		SB-01 (11-12)		8/13/2010	1200							Hold	INVE	Refusal at 12 feet bgs in basalt
SB-2 824082.42	392417.40		0-1	8/12/2010		X	Х	Χ					INVE	
		SB-02 (1-2)	1-2	8/12/2010		Χ	Χ	Χ	X				INVE	
		SB-102 (1-2)	1-2	8/12/2010					X				DUP	
		SB-02 (4-5)	4-5	8/12/2010		X	Χ	Χ					INVE	
		SB-02 (9-10)		8/12/2010		X		X				Hold	INVE	
		SB-02 (11.5-12.5)	11.5-12.5	8/12/2010	1310	X		Χ				Hold	INVE	Refusal at 12.5 feet bgs in basalt
SB-3 824118.55	392396.86	SB-03 (0-1)	0-1	8/12/2010	1130	Χ	X	Χ					INVE	
		SB-03 (1-2)		8/12/2010		X	X	X	X				INVE	
		SB-03 (4-5)	4-5	8/12/2010		X	X	X					INVE	
		SB-03 (9-10)		8/12/2010		Χ		Χ				Hold	INVE	
		SB-03 (12-13)		8/12/2010		Χ		X				Hold	INVE	Refusal at 13.5 feet bgs in basalt
SB-4 824140.18	392416.10	SB-04 (0-1)	0-1	8/12/2010	1050	Χ	Х	Χ					INVE	
		SB-04 (1-2)		8/12/2010	1054	Χ	X	Χ	Χ				INVE	
		SB-04 (4-5)	4-5	8/12/2010	1056	Χ	X	Χ					INVE	
		SB-104 (4-5)	4-5	8/12/2010	1001	X	Χ	X					DUP	
		SB-04 (9-10)	9-10	8/12/2010	1100	X		X				Hold	INVE	
		SB-04 (12-13)	12-13	8/12/2010	1110	Χ		X				Hold	INVE	Refusal at 13 feet bgs in basalt
North Sulfuric Acid Plant Neutralization tank & unloading														
_	391900.56	SB-05 (0-1)	0-1	8/6/2010	809	X	Χ	Χ					INVE	
		SB-05 (1-2)		8/6/2010	810	Χ	Χ	Χ	Χ		Χ		INVE	
		SB-05 (4-5)	4-5	8/6/2010	811	Χ	Χ	Χ					INVE	Textile Liner at 4.5 foot bgs
		SB-05 (9-10)		8/6/2010	813	Χ		Χ				Hold	INVE	· ·
		SB-05 (14-15)		8/6/2010	815	Χ		Χ				Hold	INVE	
		SB-05 (19-20)		8/6/2010	817	Χ		Χ				Hold	INVE	
		SB-05 (21-22)		8/6/2010	820	Χ		X				Hold	INVE	Refusal at 21.75 feet bgs in weathered basalt
SB-6 824286.93	391773.72			8/6/2010	905	X	Х	Χ					INVE	
		SB-06 (1-2)		8/6/2010	906	X	Χ	X	X		Χ		INVE	
		SB-06 (4-5)	4-5	8/6/2010	907	X	Χ	X					INVE	
		SB-06 (9-10)	9-10	8/6/2010	915	X		X				Hold	INVE	
		SB-06 (14-15)		8/6/2010	920	X		X				Hold	INVE	
		SB-06 (18-19)	18-19	8/6/2010	930	Χ		X				Hold	INVE	Refusal at 19 feet bgs in weathered basalt

Areas of Interest and Soil Boring L	ocations	San	nple Inform	nation				General	Chemical Rad Ch	Parameters nem ^(e)	GRO &	Analysis	Sample Type	
<u>Soil Boring</u> <u>Easting</u> North Sulfuric Acid Plant (continued) Acid Plant Building	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	Нq	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	Additional Comments
SB-7 824290.81	391669.72	SB-07 (0-1)	0-1	8/6/2010	1000	X	X	Χ					INVE	
		SB-107 (0-1)	0-1	8/6/2010	1200	Χ	Χ	Χ					DUP	
		SB-07 (1-2)	1-2	8/6/2010	1001	X	Χ	Χ	Χ		X		INVE	
		SB-07 (4-5)	4-5	8/6/2010	1003	X	Χ	X					INVE	
		SB-07 (9-10)	9-10	8/6/2010	1005	X		Χ				Hold	INVE	
		SB-07 (14-15)	14-15	8/6/2010	1007	X		X				Hold	INVE	
		SB-07 (18-19)	18-19	8/6/2010	1015	Х		Χ				Hold	INVE	Refusal at 19.5 feet bgs in weathered basalt. Wet from 19 to 19.5 feet bgs
SB-8 824405.73	391673 18	SB-08 (0-1)	0-1	8/4/2010	859	Χ	Χ	X			Х		INVE	
05 0 021100.10	001070110	SB-08 (1-2)	1-2	8/4/2010	900	X	X	X	Χ	Χ	X		INVE	
		SB-108 (1-2)	1-2	8/4/2010	902				X	X	X		DUP	
		SB-08 (4-5)	4-5	8/4/2010	800	X	X	Χ			Χ		INVE	
		SB-08 (9-10)	9-10	8/4/2010	923	Χ		Χ			X	Hold	INVE	
		SB-08 (14-15)	14-15	8/4/2010	930	Χ		Χ			X	Hold	INVE	
		SB-08 (19-20)	19-20	8/4/2010	1015	Χ		Χ			Х	Hold	INVE	Refusal at 21 feet bgs in weathered basalt
SB-9 824487.31	391762.59	SB-09 (0-1)	0-1	8/6/2010	829	Χ	Χ	X			Χ		INVE	
		SB-09 (1-2)	1-2	8/6/2010	830	X	X	X	Χ	Χ	X		INVE	
		SB-109 (1-2)	1-2	8/6/2010	800				Χ	Χ			DUP	
		SB-09 (4-5)	4-5	8/6/2010	835	X	Χ	Χ			X		INVE	
		SB-109 (4-5)	4-5	8/6/2010	800	X	Χ	Χ			X		DUP	
		SB-09 (9-10)	9-10	8/6/2010	840	Χ		Χ			X	Hold	INVE	
		SB-09 (14-15)		8/6/2010	845	X		X			Χ	Hold	INVE	
		SB-09 (19-20)	19-20	8/6/2010	900	Х		Χ			Х	Hold	INVE	Refusal at 21 feet bgs in weathered basalt
East Sulfuric Acid Plant Control room and tanks														
SB-10 824262.64	391570.58	SB-10 (0-1)	0-1	8/4/2010	1159	X	Χ	Χ					INVE	
		SB-10 (1-2)	1-2	8/4/2010	1200	X	Χ	X	Χ				INVE	
		SB-10 (4-5)	4-5	8/4/2010	1202	X	X	X					INVE	
		SB-10 (9-10)	9-10	8/4/2010	1205	X		X				Hold	INVE	
		SB-10 (14-15)	14-15	8/4/2010	1208	Х		Χ				Hold	INVE	Refusal at 17.5 feet bgs in weathered basalt
SB-11 824271.46	391516.27	SB-11 (0-1)	0-1	8/4/2010	1333	X	Χ	Χ					INVE	
		SB-111 (0-1)	0-1	8/4/2010	1230	X	Χ	Χ					DUP	
		SB-11 (1-2)	1-2	8/4/2010	1335	X	Χ	Χ	Χ				INVE	
		SB-11 (4-5)		8/4/2010	1337	X	Χ	X					INVE	
		SB-11 (9-10)		8/4/2010		X		X				Hold	INVE	
	5	SB-11 (12.5-13.5)	12.5-13.5	8/4/2010	1345	Χ		Χ				Hold	INVE	Refusal at 15.5 feet bgs in weathered basalt
SB-12 824388.94	391528.69	SB-12 (0-1)	0-1	8/4/2010	1113	Χ	Χ	Χ					INVE	
		SB-12 (1-2)	1-2	8/4/2010	1115	X	Χ	X	Χ				INVE	
		SB-12 (4-5)		8/4/2010		X	Χ	X					INVE	
		SB-12 (9-10)		8/4/2010		X		X				Hold	INVE	
		SB-12 (14-15)	14-15	8/4/2010	1130	Χ		Χ				Hold	INVE	Refusal at 16.5 feet bgs in weathered basalt

Areas of Interest and Soil Boring L	_ocations	San	nple Inforn	mation				General	Chemical Rad Ch	Parameters nem ^(e)	GRO &	Analysis	Sample Type	
Soil Boring Easting East Sulfuric Acid Plant (continued)	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	<u>рН</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold (b)	(INVE/DUP)	Additional Comments
SB-13 824278.54	391434.67	SB-13 (0-1)	0-1	8/4/2010	1414	X	X	Χ					INVE	
		SB-13 (1-2)	1-2	8/4/2010	1415	Χ	Χ	Χ	Χ	X			INVE	
		SB-13 (4-5)	4-5	8/4/2010	1417	Χ	Χ	Χ					INVE	
		SB-13 (9-10)	9-10	8/4/2010	1420	Χ		Χ				Hold	INVE	
		SB-13 (14-15)	14-15	8/4/2010	1430	Χ		Χ				Hold	INVE	Refusal at 15.5 feet bgs in weathered basalt
SB-14 824272.63	391339.71	SB-14 (0-1)	0-1	8/4/2010	1454	Χ	X	X					INVE	
		SB-14 (1-2)	1-2	8/4/2010	1455	X	Χ	X	Χ				INVE	
		SB-14 (4-5)	4-5	8/4/2010	1457	X	Χ	X					INVE	
		SB-14 (9-10)	9-10	8/4/2010	1500	Χ		Χ				Hold	INVE	
		SB-14 (14-15)	14-15	8/4/2010	1502	Х		Χ				Hold	INVE	Refusal at 16 feet bgs in weathered basalt
SB-15 824355.37	391310.14	SB-15 (0-1)	0-1	8/4/2010	1523	Χ	Χ	X					INVE	
		SB-15 (1-2)	1-2	8/4/2010	1525	Χ	X X	Χ	X				INVE	
		SB-15 (4-5)	4-5	8/4/2010	1527	X	Х	X					INVE	
		SB-15 (9-10)	9-10	8/4/2010	1529	X		X				Hold	INVE	
		SB-15 (14-15)	14-15	8/4/2010	1531	Χ		Χ				Hold	INVE	
		SB-15 (19-20)	19-20	8/4/2010	1535	X		X				Hold	INVE	
		SB-15 (24-25)	24-25	8/4/2010	1540	X		X				Hold	INVE	
		SB-15 (26-27)	26-27	8/4/2010	1610	Χ		Х				Hold		Refusal at 28.5 feet bgs in weathered basalt; wet from 27-28.5 feet bgs
PPA Plant														
Control room and tanks	200000 00	CD 40 (0.4)	0.4	0/0/0040	4.405	V	V	V					INI\/=	
SB-16 824390.96	390986.06	SB-16 (0-1)	0-1	8/2/2010	1405	X	X	X	V				INVE INVE	
		SB-16 (1-2)	1-2	8/2/2010 8/2/2010	1410 1415	X X	X X	X X	Χ				INVE	
		SB-16 (4-5)	4-5 8-9	8/2/2010	1415	X	^	X				Hold		Refusal at 9.5 feet bgs in weathered basalt
		SB-16 (8-9)	6-9	6/2/2010	1420	^		^				Holu	IINVE	Refusal at 9.5 feet bys in weathered basait
SB-17 824356.34	390743.45	SB-17 (0-1)	0-1	8/2/2010	1528	X	Χ	Χ					INVE	
		SB-17 (1-2)	1-2	8/2/2010	1529	X	Χ	Χ	X				INVE	
		SB-17 (4-5)	4-5	8/2/2010	1530	Χ	Χ	Χ					INVE	
		SB-17 (9-10)	9-10	8/2/2010	1535	X						Hold	INVE	
		SB-17 (13-14)	14-15	8/2/2010	1537	Х						Hold	INVE	Refusal at 14 feet bgs in weathered basalt
SB-18 824394.03	390581.64	SB-18 (0-1)	0-1	8/2/2010	1604	Χ	Χ	X					INVE	
		SB-118 (0-1)	0-1	8/2/2010	1550	X	Χ	Χ					DUP	
		SB-18 (1-2)	1-2	8/2/2010	1605	X	Χ	Χ	Χ				INVE	
		SB-18 (4-5)	4-5	8/2/2010	1606	Χ	Χ	X					INVE	
		SB-18 (8.5-9.5)	8.5-9.5	8/2/2010	1608	Χ						Hold	INVE	Refusal at 9.5 feet bgs in weathered basalt

Areas of Interest and Soil Boring Locations Sample Information								General	Chemical Rad Ch	Parameters nem ^(e)	GRO &	Analysis	Sample Type			
<u>Soi</u> South Ball Mill	il Boring	Easting	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	<u>Hq</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	Additional Comments
July 1991	SB-19 8	823892.81	392031.54		0-1	8/9/2010	1450	X	X	X	V	V			INVE INVE	textile liner at 1 foot bgs
				SB-19 (1-2) SB-19 (4-5)	1-2 4-5	8/9/2010 8/9/2010	1455 1457	X X	X X	X X	Х	Χ			INVE	
				SB-19 (9-10)	9-10	8/9/2010	1500	X	^	X				Hold	INVE	
				SB-19 (14-15)		8/9/2010	1505	Χ		Χ				Hold		Refusal at 16 feet bgs in weathered basalt
	SB-20 8	323971.94	392033.59			8/18/2010		Χ	Χ	X					INVE	
				SB-20 (1-2)	1-2	8/18/2010		X	X	X	X				INVE	
				SB-20 (4-5)	4-5	8/18/2010		X	Χ	X				11-1-1	INVE	
				SB-20 (9-10) SB-20 (13-14)	9-10	8/18/2010 8/18/2010		X X		X X				Hold Hold	INVE INVE	Refusal at 14 feet bgs in basalt
														Поіц		Netusal at 14 leet bys III basait
	SB-21 8	324070.17	391991.85		0-1	8/9/2010		X	X	X					INVE	
				SB-21 (1-2)	1-2	8/9/2010	1407	X	X X	X	X				INVE	
				SB-21 (4-5) SB-121 (4-5)	4-5 4-5		1410	X	X X	X					INVE DUP	
				SB-121 (4-5) SB-21 (9-10)	4-5 9-10	8/9/2010 8/9/2010	1345 1415	X X	^	X X				Hold	INVE	
				SB-21 (14-15)			1420	X		X				Hold	INVE	
				SB-21 (15.5-16.5)				X		X				Hold		Refusal at 16.5 feet bgs in weathered basalt
Phosphoric and North SPA Area	-	-		00.00 (0.4)	0.4	0/0/0040	4450	v	V	V					IAN /E	
	SB-22 8	324128.88	391851.56			8/9/2010		X	X	X	V				INVE	
				SB-22 (1-2) SB-22 (4-5)	1-2 4-5	8/9/2010 8/9/2010		X X	X X	X X	Х				INVE INVE	
				SB-22 (4-5)	9-10	8/9/2010		X	^	X				Hold	INVE	
				SB-22 (14-15)		8/9/2010		X		X				Hold	INVE	
				SB-22 (15.5-16.5)				X		X				Hold		Refusal at 18 feet bgs in weathered basalt
	SB-23 8	324082.66	391849.09	SB-23 (0-1)	0-1	8/9/2010	1320	Χ	Х	X					INVE	
				SB-23 (1-2)	1-2	8/9/2010		X	Χ	X	Χ				INVE	
				SB-23 (2.5-3.5)	2.5-3.5	8/9/2010	1325	Χ	Χ	X					INVE	Liner at 1.5 feet bgs; Refusal at 3.5 feet bgs in silt
	SB-24	824095.21	391786.97	SB-24 (0-1)	0-1	8/9/2010	1030	X	Х	X					INVE	textile liner at 1 foot bgs
	052.	JE 1000.E1	001700.07	SB-24 (1-2)	1-2	8/9/2010		X	X	X	Χ	X			INVE	toxus uner at 1 loct age
				SB-24 (4-5)		8/9/2010		Χ	Χ	X					INVE	
				SB-24 (6-7)		8/9/2010		Х		Χ				Hold	INVE	Refusal at 7 feet bgs in silt
South Car Wash S																Logotion amittade inconnecible
	SB-25	-	-	-	-	-	-								-	Location omitted; inaccessible
	SB-26 8	324161.97	391755.83		0-1	8/18/2010		Χ	Χ	Χ					INVE	
				SB-26 (1-2)		8/18/2010		Χ	Χ	Χ	X				INVE	
				SB-26 (4-5)		8/18/2010		X	X	X					INVE	D () 1 () () () ()
				SB-26 (7-8)	7-8	8/18/2010	1435	Χ	X	X					INVE	Refusal at 8 feet bgs in silt

Areas of	f Interest and S	Soil Boring L	ocations	San	nple Inforr	mation				General	Chemical Rad Ch	Parameters nem ^(e)	GRO &	Analysis	Sample Type	
	Soil Boring and Super Ph		<u>Northing</u> id Area (cont	Sample ID inued)	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	Нq	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold (b)	(INVE/DUP)	Additional Comments
SPA Ditch	CD 07	824075.64	204722.02	SB-27 (0-1)	0.4	8/9/2010	900	V	V	V					INVE	Textile Liner at 1 foot bgs
	3D-21	024075.04	391733.92	SB-127 (0-1)	0-1 0-1	8/9/2010		X X	X	X X					DUP	Textile Liner at 1 100t bgs
				SB-27 (1-2)	1-2	8/9/2010		X	X X	X	Χ				INVE	
				SB-127 (1-2)	1-2	8/9/2010		χ	^	χ	X				DUP	
				SB-27 (4-5)	4-5	8/9/2010		Χ	Х	X	Α				INVE	
				SB-27 (8-9)	8-9	8/9/2010		X	^	X				Hold		Refusal at 9 feet bgs in silt
	SB-28	824073.22	391676.55	SB-28 (0-1)	0-1	8/5/2010	1529	X	Х	Χ					INVE	Textile Liner at 1 foot bgs
	02 20	00. 0	00.0.00	SB-28 (1-2)	1-2	8/5/2010		X	X	X	X	X			INVE	
				SB-28 (3.5-4.5)		8/5/2010		X	X	X					INVE	Refusal at 4.5 feet bgs in silty sand
	SB-29	824070.80	391614 09	SB-29 (0-1)	0-1	8/5/2010	1445	X	Х	Χ					INVE	Textile Liner at 1 foot bgs
	OB 2 0	02 107 0.00	001011.00	SB-29 (1-2)	1-2	8/5/2010		X	X	X	Χ				INVE	Tokalo Elifor at 1 100t bgb
				SB-29 (4-5)	4-5	8/5/2010		X	X	X	,,				INVE	
				SB-29 (8-9)	8-9	8/5/2010		X		X				Hold		Refusal at 9 feet bgs in silty sand
	SB-30	824078.32	391534 12	SB-30 (0-1)	0-1	8/5/2010	1104	X	Х	Χ					INVE	
	02 00	02 101 0.02	001001112	SB-30 (1-2)	1-2	8/5/2010		X	X	X	Χ					Textile liner at 2 feet bgs
				SB-30 (4-5)	4-5	8/5/2010		X	X	X					INVE	Refusal at 5 feet bgs in dense fill material
SPA Building	α															
		824101.76	391550.45	SB-31 (0-1)	0-1	8/5/2010	1409	X	Χ	Χ					INVE	Textile Liner at 1 foot bgs
				SB-31 (1-2)	1-2	8/5/2010		Χ	Χ	Χ	Χ				INVE	S
				SB-131 (1-2)	1-2	8/5/2010		Χ	Χ	Χ					DUP	
				SB-31 (4-5)	4-5	8/5/2010	1411	X	Χ	Χ					INVE	
				SB-31 (9-10)	9-10	8/5/2010	1413	Χ		X				Hold	INVE	Refusal at 11 feet bgs in silt
	SB-32	824158.81	391548.21	SB-32 (0-1)	0-1	8/5/2010	1225	Χ	Х	X					INVE	Textile Liner at 1 foot bgs
				SB-32 (1-2)	1-2	8/5/2010	1227	Χ	X X	X	Χ	Χ			INVE	
				SB-32 (4-5)	4-5	8/5/2010	1229	Χ	Χ	Χ					INVE	
				SB-32 (7-8)	7-8	8/5/2010	1234	X		Χ				Hold	INVE	
				SB-32 (14-15)	14-15	8/5/2010	1250	Χ		X				Hold	INVE	Refusal at 16.5 feet bgs in weathered basalt
North Car W	ash Sump Area															
	SB-33	824182.99	391985.43	SB-33 (0-1)	0-1	8/18/2010		X	Х	X					INVE	
				SB-33 (1-2)	1-2	8/18/2010		X	X	X	X				INVE	
				SB-33 (3.5-4)	3.5-4	8/18/2010	1400	Χ	Χ	Х					INVE	Hand auger refusal at 4 feet bgs on gravel
	SB-34	824172.36	392031.72	SB-134 (0-1)	0-1	8/18/2010		X	Χ	Χ						Sample not analyzed (duplicate)
				SB-34 (0-1)	0-1	8/18/2010		Χ	Χ	X					INVE	
				SB-34 (1-2.5)	1-2.5	8/18/2010		X	X	X	X	X			INVE	
				SB-34 (4-5)	4-5	8/18/2010		X	X	X					INVE	
				SB-34 (9-10)	9-10	8/18/2010		X	Χ	X					INVE	
				SB-34 (14-15)	14-15	8/18/2010		X		X				Hold	INVE	
				SB-34 (19-20)	19-20	8/18/2010		X		X				Hold	INVE	
				SB-34 (24-25)	24-25	8/18/2010	1320	Χ		Х				Hold	INVE	Refusal at 29 feet bgs in basalt; wet at 26 feet bgs
																J,

Areas of Interest and Soil Bori	Sar	mple Infori	mation				General	Chemical Rad Ch	Parameters nem ^(e)	GRO &	Analysis	Sample Type		
Soil Boring Eastin Central Tank Farm Area	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals ^(c)	<u>рН</u>	Chemistry (d)	Gross a & B	<u>Ra-226/228</u>	BTEX (f)	on Hold ^(b)	(INVE/DUP)	<u>Additional Comments</u>
Central Tank Farm SB-35 823854.	58 391671.02	SB-35 (0-1) SB-35 (1-2) SB-35 (4-5) SB-135 (4-5) SB-35 (9-10) SB-35 (14-15)	1-2 4-5 4-5 9-10	8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010	1423 1425 1411 1430	X X X X X	X X X	X X X X X	X	X		Hold Hold	INVE INVE INVE DUP INVE INVE	Refusal at 15 feet bgs in silt
SB-36 823885.	84 391559.76	SB-36 (0-1) SB-36 (1-2) SB-36 (4-5)	1-2	8/17/2010 8/17/2010 8/17/2010	1145	X X X	X X X	X X X	Х				INVE INVE INVE	Refusal at 5 feet bgs in silt
SB-37 823931.	69 391549.23	SB-37 (0-1) SB-137 (0-1) SB-37 (1-2) SB-137 (1-2) SB-37 (4-5) SB-137 (4-5)	0-1 1-2 1-2 4-5	8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010	1031 1053 1033 1055	X X X X X	X X X X X	X X X X	X X				INVE DUP INVE DUP INVE DUP	Refusal at 5 feet bgs in silt
SB-38 824016.	53 391542.26	SB-38 (0-1) SB-38 (1-2) SB-38 (4-5)	1-2	8/17/2010 8/17/2010 8/17/2010	1020	X X X	X X X	X X X	Х				INVE INVE INVE	Refusal at 5.25 feet bgs in silt
Oxidation Tank Farm Area SB-39 823982.	56 391518.83	SB-39 (0-1) SB-39 (1-2) SB-39 (4-5)	1-2	8/17/2010 8/17/2010 8/17/2010	952	X X X	X X X	X X X	Х	X			INVE INVE INVE	Refusal at 5 feet bgs in silt; sample container not intact and thus not analyzed
SB-40 823887.	20 391491.88	SB-40 (0-1) SB-40 (1-2) SB-40 (4-5) SB-140 (4-5) SB-40 (9-10) SB-40 (11.5-12.5)	1-2 4-5 4-5 9-10	8/10/2010 8/10/2010 8/10/2010 8/10/2010 8/10/2010 8/10/2010	1602 1604 1730 1610	X X X X X	X X X X	X X X X X	Х			Hold Hold	INVE INVE INVE DUP INVE INVE	Refusal at 12.5 feet bgs in silt
SB-41 823984.	52 391365.73	SB-41 (0-1) SB-41 (1-2) SB-41 (4-5) SB-41 (9-10) SB-41 (14-15) SB-41 (17.5-18.5)	1-2 4-5 9-10 14-15	8/10/2010 8/10/2010 8/10/2010 8/10/2010 8/10/2010 8/10/2010	1521 1525 1530 1535	X X X X X	X X X	X X X X	Χ			Hold Hold Hold	INVE INVE INVE INVE INVE	Refusal at 18.5 feet bgs in silt
SB-42 824025.	60 391312.63	SB-42 (0-1) SB-42 (1-2) SB-42 (4-5) SB-42 (9-10) SB-42 (14-15) SB-42 (19-20) SB-42 (24-25)	1-2 4-5 9-10 14-15 19-20	8/13/2010 8/13/2010 8/13/2010 8/13/2010 8/13/2010 8/13/2010 8/13/2010	1653 1655 1659 1705 1710	X X X X X	X X X	X X X X X	Х			Hold Hold Hold Hold	INVE INVE INVE INVE INVE INVE	Refuesed at 25.5 feet in basalt

Areas of Interest and Soil Boring Locations		San	nple Inforn	nation				General	Chemical Rad Ch	Parameters	GRO &	Analysis	Sample Type			
Solution Solution Oxidation Tan	Soil Boring		Northing	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	Ηq	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	<u>Additional Comments</u>
Oxidation ran		824084.42		SB-43 (0-1)	0-1	8/5/2010	1019	Χ	Χ	X					INVE	
				SB-43 (1-2)	1-2	8/5/2010		X	Χ	X	Χ				INVE	
				SB-43 (4-5)		8/5/2010		X	Χ	X					INVE	
				SB-43 (9-10)		8/5/2010		X		X				Hold	INVE	
				SB-43 (14-15) SB-43 (17.5-18.5)		8/5/2010 8/5/2010		X X		X X				Hold Hold	INVE INVE	Refusal at 18.5 in weathered basalt
				30-43 (17.5-16.5)	17.5-10.5	0/3/2010	1033	^		^				riola	IIVL	iverusai at 10.5 iii weathered basait
West Sulfuric	Acid Plant															
West Shop	OD 44	000000 70	004000 4	00 44 (0.4)	0.4	0/40/0040	4.440	V		V					IN 17 /F	
	SB-44	823900.78	391392.4 <i>i</i>			8/10/2010 8/10/2010		X	X	X X	V				INVE INVE	
				SB-44 (1-2) SB-144 (1-2)		8/10/2010		X X	X	×	X				DUP	
				SB-44 (4-5)	4-5	8/10/2010		X	X X	X					INVE	
				SB-44 (9-10)		8/10/2010		X	•	X				Hold	INVE	
				SB-44 (14-15)		8/10/2010		X		Χ				Hold	INVE	
				SB-44 (16.5-17.5)	16.5-17.5	8/10/2010	1440	Χ		X				Hold	INVE	Refusal at 18 in weathered basalt
Granulation And Granulation Bu	ilding															
	SB-45	824138.42	391153.93		0-1	8/16/2010		X	X	X					INVE	
				SB-45 (1-2)		8/16/2010		X	X X	X	X	X			INVE	
				SB-45 (4-5) SB-45 (9-10)	4-5 9-10	8/16/2010 8/16/2010		X X	Χ.	X X				Hold	INVE INVE	
				SB-45 (14-15)	14-15	8/16/2010		X		X				Hold	INVE	
				SB-45 (19-20)	19-20	8/16/2010		X		X				Hold	INVE	
				SB-45 (23-24)		8/16/2010		Χ		X				Hold	INVE	Refusal at 24 feet bgs in basalt
	CD 46	824030.19	201222	SB-46 (0-1)	0-1	8/16/2010	1215	Χ	~	Χ					INVE	
	3D-40	024030.19	391222.32	SB-46 (1-2)		8/16/2010		X	X X	x	X				INVE	
				SB-46 (4-5)	4-5	8/16/2010		X	X	X	Λ				INVE	
				SB-146 (4-5)	4-5	8/16/2010		Χ	Χ	Χ					DUP	
				SB-46 (9-10)		8/16/2010		Χ		X				Hold	INVE	
				SB-46 (14-15)		8/16/2010		X		X				Hold	INVE	
				SB-46 (15-16)	15-16	8/16/2010	1340	Χ		Х				Hold	INVE	Refusal at 16 feet bgs in basalt
	SB-47	824003.71	391136.61	SB-47 (0-1)	0-1	8/12/2010	1435	Χ	Χ	X					INVE	
				SB-147 (0-1)		8/12/2010	1330	X	Χ	X					DUP	
				SB-47 (1-2)		8/12/2010		Χ	Χ	X	Χ	X			INVE	
				SB-47 (4-5)		8/12/2010		X	X	X					INVE	
				SB-47 (9-10)	9-10	8/12/2010	1444	Χ		X				Hold	INVE	Refusal at 13 feet bgs in basalt; no recovery 10 to 13 feet bgs
																13 leet bys
	SB-48	824026.77	391097.51			8/12/2010		X	X	X					INVE	
				SB-48 (1-2)		8/12/2010		X	X	X	Χ				INVE	
				SB-48 (4-5)		8/12/2010		X	Χ	X					INVE	
				SB-48 (9-10)		8/12/2010		X		X				Hold	INVE	Defugal at 45 feet has in becalt
				SB-48 (14-15)	14-15	8/12/2010	1530	Χ		Χ				Hold	INVE	Refusal at 15 feet bgs in basalt
	SB-49	824018.76	391044.32			8/12/2010		Χ	Χ	X					INVE	
				SB-49 (1-2)		8/12/2010		Χ	X X	X	Χ				INVE	
				SB-49 (4-5)		8/12/2010		X	X	X					INVE	
				SB-49 (9-10)		8/12/2010		X		X				Hold	INVE	Defined at 40.5 feet has in besself
				SB-49 (11-12)	11-12	8/12/2010	1605	Χ		X				Hold	INVE	Refusal at 12.5 feet bgs in basalt

Areas of Interest and Soil Boring Locations		San	nple Infor	mation				General		Parameters hem ^(e)	GRO &	Analysis	Sample Type	
Soil Boring Easting	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	<u>Hq</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold (b)	(INVE/DUP)	Additional Comments
Process Sewer Lagoon														
Area North of Lagoon														
SB-50 823644.	05 392077.4	SB-50 (0-1) SB-50 (1-2) SB-50 (4-5) SB-50 (9-10) SB-50 (13-14)	4-5 9-10	8/18/2010 8/18/2010 8/18/2010 8/18/2010 8/18/2010	1012 1015 1020	X X X X	X X X	X X X X	X			Hold Hold	INVE INVE INVE INVE	Refusal at 14 feet bgs in basalt
SB-51 823766.	392085.62	SB-51 (0-1) SB-51 (1-2) SB-51 (2-3)	0-1 1-2 2-3	8/18/2010 8/18/2010 8/18/2010	1057	X X X	X X X	X X X	X				INVE INVE INVE	Hand auger refusal at 3 feet bgs
SB-52 823812.	392012.60 392012.60	SB-52 (0-1) SB-52 (1-2) SB-52 (4-5) SB-52 (9-10) SB-52 (14-15) SB-52 (19-20) SB-52 (23-24)	0-1 1-2 4-5 9-10 14-15 19-20 23-24	8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010 8/17/2010	1554 1557 1600 1602 1605	X X X X X	X X X	X X X X X	X			Hold Hold Hold Hold	INVE INVE INVE INVE INVE INVE	Refusal at 24 feet bgs in silt
Process Sewer Lagoon Sump SB-53 823829.	23 391891.6	7 SB-53 (0-1) SB-53 (1-2) SB-53 (4-5) SB-53 (9-10) SB-53 (14-15) SB-153 (14-15) SB-53 (16-17)	1-2 4-5 9-10 14-15 14-15	8/18/2010 8/18/2010 8/18/2010 8/18/2010 8/18/2010 8/18/2010	934 937 940 945 939	X X X X X	X X X X X	X X X X X	X			Hold	INVE INVE INVE INVE DUP INVE	Refusal at 17 feet bgs in basalt
SB-54 823829.	19 391858.09	SB-54 (0-1) SB-54 (1-2) SB-54 (4-5) SB-54 (9-10) SB-154 (9-10) SB-54 (14-15) SB-54 (17-18)		8/18/2010 8/18/2010 8/18/2010 8/18/2010 8/18/2010 8/18/2010 8/18/2010	853 856 859 847 905	X X X X X	X X X X X	X X X X X	X	X		Hold	INVE INVE INVE INVE DUP INVE INVE	Refusal at 17 feet bgs in basalt
Process Sewer Lagoon SB-55 823678.	95 391972.88	SB-55 (0-1) SB-55 (1-2) SB-55 (4-5) SB-55 (9-10) SB-55 (14-15) SB-55 (19-20) SB-55 (24.5-25.5)	1-2 4-5 9-10 14-15 19-20	8/9/2010 8/9/2010 8/9/2010 8/9/2010 8/9/2010 8/9/2010 5 8/9/2010	1527 1530 1535 1540 1543	X X X X X	X X X X	X X X X X	X			Hold Hold	INVE INVE INVE INVE INVE INVE	Refusal at 25.5 feet bgs in basalt

Soil Sample Summary - 2010 Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Areas of Interest and Soil Boring Locations Sample Information Chemical Parameters General Rad Chem (e) GRO & Analysis Sample Type Metals (c) Chemistry (d) Ra-226/228 BTEX (f) on Hold (b) (INVE/DUP) Soil Boring Easting Sample ID <u>Time</u> <u>рН</u> Gross a & B **Additional Comments Depths** <u>Date</u> **Process Sewer Lagoon Area (continued)** INVE SB-56 823696.69 391871.27 SB-56 (0-1) 0-1 8/10/2010 Χ SB-56 (1-2) 1-2 8/10/2010 815 Χ Χ Χ INVE SB-56 (4-5) 4-5 8/10/2010 817 Χ Χ Χ INVE Χ SB-56 (9-10) 9-10 8/10/2010 821 Χ Χ INVE Χ SB-156 (9-10) 9-10 8/10/2010 730 Χ Χ DUP Χ Χ Χ INVE SB-56 (14-15) 14-15 8/10/2010 830 Χ SB-56 (18.5-19.5) 18.5-19.5 8/10/2010 835 Χ Hold INVE Refusal at 19.5 feet bgs in weathered basalt SB-57 823722.24 391810.93 SB-57 (0-1) 0-1 8/10/2010 Χ Χ Х INVE SB-57 (1-2) 8/10/2010 902 Χ Χ INVE 1-2 Χ Χ DUP SB-157 (1-2) 1-2 8/10/2010 845 SB-57 (4-5) 8/10/2010 Χ Χ Χ INVE 905 4-5 SB-57 (9-10) 8/10/2010 910 Χ Χ Χ INVE 9-10 Χ SB-57 (14-15) 8/10/2010 915 Χ Χ INVE 14-15 Χ SB-57 (19-20) 19-20 8/10/2010 920 Χ Hold INVE SB-57 (24-25) Χ 24-25 8/10/2010 925 Χ Hold INVE Refusal at 26 feet bgs in weathered basalt Χ INVE SB-58 823793.02 391813.73 SB-58 (0-1) 0-1 8/10/2010 1209 Χ SB-58 (1-2) 1-2 8/10/2010 1210 Χ Χ Χ INVE Χ Χ INVE SB-58 (4-5) 4-5 8/10/2010 1211 Χ SB-58 (9-10) Χ Χ INVE 9-10 8/10/2010 1212 Χ Refusal at 19 in weathered basalt; wet at 16 feet Χ Χ Χ INVE SB-58 (14-15) 14-15 8/10/2010 1213 SB-158 (14-15) 14-15 8/10/2010 1215 Χ Χ DUP **Dry Products Storage & East Ditch** East Ditch INVE Χ Χ Χ SB-59 823846.00 391737.11 SB-59 (0-1) 0-1 8/17/2010 1525 Χ Χ INVE SB-59 (1-2) 1-2 8/17/2010 1527 Χ Χ SB-159 (1-2) DUP 1-2 8/17/2010 1500 Χ Χ SB-59 (4-5) 4-5 8/17/2010 1530 Χ INVE Refusal at 5 feet bgs in slag SB-60 823851.98 391618.67 Χ INVE SB-60 (0-1) 0-1 8/17/2010 1355 Χ SB-60 (1-2) 1-2 8/17/2010 1358 Χ Χ Χ INVE Χ SB-60 (4-5) 4-5 8/17/2010 1400 Χ Χ Χ INVE SB-60 (6-7) 6-7 8/17/2010 1402 Χ Χ Hold INVE Refusal at 7 feet bgs in lithfied silt Χ INVE SB-61 823856.55 391527.31 SB-61 (0-1) 0-1 8/17/2010 1330 Χ Χ SB-61 (1-2) 1-2 8/17/2010 1335 Χ Χ Χ Χ INVE Χ Χ Χ INVE SB-61 (4-5) 4-5 8/17/2010 1400 Χ Χ SB-61 (5.5-6.5) 5.5-6.5 8/17/2010 1402 Hold INVE Refusal at 6.5 feet bgs in silt/tufa Χ INVE SB-62 823883.23 391428.02 Χ Χ SB-62 (0-1) 0-1 8/17/2010 900 INVE SB-62 (1-2) 902 Χ Χ Χ 1-2 8/17/2010 Χ SB-62 (4-5) 4-5 8/17/2010 905 Χ INVE SB-62 (9-10) 9-10 8/17/2010 910 Χ Χ Hold INVE SB-62 (14-15) 14-15 8/17/2010 920 Χ Χ Hold INVE Χ Χ SB-62 (19-20) 19-20 8/17/2010 925 Hold INVE Refusal at 20 feet bgs in basalt

Areas of Interest and Soil Bori	ng Locations	San	nple Infori	mation				General	Chemical Rad C	Parameters hem ^(e)	GRO &	Analysis	Sample Type	
Soil Boring Easting		Sample ID	Depths	<u>Date</u>	<u>Time</u>	Metals (c)	Нq	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	Additional Comments
Dry Products Storage & East Ditch	(continued)													
North End of Building	44 004707.00	OD 00 (0.4)	0.4	0/40/0040	055	V		V					IN 17 / E	
SB-63 823692.	41 391767.39	SB-63 (0-1)	0-1	8/10/2010		X	X X	X	V				INVE	
		SB-63 (1-2)	1-2	8/10/2010		X	X	X	X				INVE	
		SB-63 (4-5)	4-5	8/10/2010		X	Х	X				الملط	INVE	
		SB-63 (9-10)	9-10	8/10/2010		X		X				Hold	INVE INVE	
		SB-63 (14-15)	14-15	8/10/2010		X X		X X				Hold	INVE	Defugal at 22 in weathered begalt
		SB-63 (19-20)	19-20	8/10/2010	1130	Χ		Χ				Hold	IINVE	Refusal at 22 in weathered basalt
SB-64 823804.	17 391778.33	SB-64 (0-1)	0-1	8/10/2010	1330	Χ	Χ	Χ					INVE	
		SB-64 (1-2)	1-2	8/10/2010	1334	X	Χ	Χ	X	Χ			INVE	
		SB-64 (4-5)	4-5	8/10/2010		Χ	Χ	Χ					INVE	
		SB-64 (9-10)	9-10	8/10/2010		Χ		Χ				Hold	INVE	
		SB-64 (14-15)	14-15	8/10/2010		Χ		Χ				Hold	INVE	
		SB-64 (19-20)	16-17	8/10/2010	1350	Χ		X				Hold	INVE	Refusal at 17.5 in weathered basalt
Loading Area														
SB-65 823642.	34 391185.64	SB-65 (0-1)	0-1	8/13/2010	805	Χ	Χ	Χ					INVE	
		SB-65 (1-2)	1-2	8/13/2010		X	X	Χ	Χ				INVE	
		SB-65 (4-5)	4-5	8/13/2010		X	Χ	Χ					INVE	
		SB-65 (9-10)	9-10	8/13/2010		X		X				Hold	INVE	
		SB-65 (14-15)	14-15	8/13/2010		X		X				Hold	INVE	
		SB-65 (19-20)	19-20	8/13/2010		X		X				Hold	INVE	Refusal at 20.5 feet bgs in weathered basalt
		22 -2 /2 /)												
SB-66 823613.	02 391134.05	SB-66 (0-1)	0-1	8/13/2010		X	X	X	V				INVE	
		SB-66 (1-2)	1-2	8/13/2010		X	X	X	X				INVE	
		SB-66 (4-5)	4-5	8/13/2010		X	Χ	X				11-1-1	INVE	
		SB-66 (9-10)	9-10	8/13/2010		X		X				Hold	INVE	
		SB-66 (14-15)	14-15	8/13/2010		X		X				Hold	INVE	
		SB-66 (19-20)	19-20	8/13/2010		X		X				Hold	INVE	Defined at 24 feet has in weathered becalt
		SB-66 (20-21)	20-21	8/13/2010	1005	Х		Χ				Hold	INVE	Refusal at 21 feet bgs in weathered basalt
SB-67 823645.	82 391086.70	SB-67 (0-1)	0-1	8/13/2010		Χ	Χ	X					INVE	
		SB-67 (1-2)	1-2	8/13/2010		Χ	Χ	Χ	Χ	Χ			INVE	
		SB-67 (4-5)	4-5	8/13/2010		X	Χ	Χ					INVE	
		SB-67 (9-10)	9-10	8/13/2010		Χ		Χ				Hold	INVE	
		SB-67 (14-15)	14-15	8/13/2010		X		Χ				Hold	INVE	
		SB-67 (19-20)	19-20	8/13/2010		X		Χ				Hold	INVE	
		SB-67 (22-23)	22-23	8/13/2010	912	X		X				Hold	INVE	Refusal at 23 feet bgs in weathered basalt

Soil Sample Summary - 2010 Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Areas of Interest and Soil Boring Locations Sample Information Chemical Parameters General Rad Chem (e) GRO & Analysis Sample Type Metals (c) Chemistry (d) Ra-226/228 BTEX (f) on Hold (b) (INVE/DUP) Soil Boring Easting Northing Sample ID **Depths** <u>рН</u> Gross a & B **Additional Comments** <u>Date</u> <u>Time</u> **Dry Products Storage & East Ditch (continued)** East and South of Building SB-68 823846.78 391240.62 SB-68 (0-1) 0-1 8/16/2010 1010 Χ Χ INVE SB-68 (1-2) 1-2 8/16/2010 1012 Χ Χ Χ INVE Χ Χ Χ INVE SB-68 (4-5) 4-5 8/16/2010 1015 SB-68 (9-10) 9-10 8/16/2010 1017 Χ Χ INVE Hold Χ Χ INVE SB-68 (14-15) 14-15 8/16/2010 1020 Hold Χ Χ SB-68 (19-20) 19-20 INVE 8/16/2010 1023 Hold Χ Χ SB-68 (21.5-22.5) 21.5-22.5 8/16/2010 1027 INVE Hold Refusal at 23 feet bgs in basalt SB-69 823855.91 391044.90 SB-69 (0-1) 0-1 Χ INVE 8/16/2010 1120 Χ SB-69 (1-2) Χ Χ Χ Χ INVE 1-2 8/16/2010 1123 SB-69 (4-5) 8/16/2010 1126 Χ Χ INVE 4-5 SB-69 (9-10) 9-10 8/16/2010 1129 Χ Χ Hold INVE Χ INVE SB-69 (14-15) 14-15 8/16/2010 1131 Χ Hold Χ SB-69 (19-20) 19-20 8/16/2010 1135 Χ Hold INVE Χ Χ SB-69 (20-21) 20-21 8/16/2010 1140 Hold INVE Refusal at 21 feet bgs in basalt SB-70 823846.13 390749.19 Χ INVE SB-70 (0-1) 0-1 8/13/2010 1100 Χ Χ Χ INVE SB-70 (1-2) 1-2 8/13/2010 1101 Χ Χ Χ Χ Χ INVE SB-70 (4-5) 4-5 8/13/2010 1103 Χ SB-70 (6.5-7.5) 6.5-7.5 8/13/2010 1107 Χ Hold INVE Refusal at 7.5 feet bgs in weathered basalt SB-171 (0-1) Χ Χ DUP SB-71 823764.42 390666.61 0-1 8/16/2010 800 Χ SB-71 (0-1) 0-1 8/16/2010 900 Χ Χ Х INVE SB-71 (1-2) 1-2 8/16/2010 905 Χ Χ Χ Χ INVE SB-171 (1-2) 1-2 8/16/2010 906 Χ DUP Χ Χ Χ INVE SB-71 (4-5) 4-5 8/16/2010 907 Χ SB-71 (5-6) 910 Χ Refusal at 7.5 feet bgs in weathered basalt 5-6 8/16/2010 Hold INVE All locations moved to the west versus work plan **Production Well NW-9** drawing based on location of well NW-9. SB-72 823969.87 390013.37 SB-72 (0-1) 0-1 8/3/2010 930 Χ INVE Χ SB-72 (1-2) 1-2 8/3/2010 932 Χ Χ Χ INVE Χ SB-72 (4-5) 8/3/2010 934 Χ Χ Χ INVE 4-5 SB-72 (9-10) 9-10 8/3/2010 936 Χ Χ INVE Hold Χ SB-72 (14-15) 14-15 8/3/2010 938 Χ Hold INVE Χ Χ SB-72 (19-20) 19-20 8/3/2010 940 Hold INVE Χ Χ SB-72 (24-25) 24-25 8/3/2010 942 Hold INVE Χ SB-72 (29-30) 29-30 8/3/2010 948 Χ Hold INVE SB-72 (31-32) 31-32 Χ Χ Hold 8/3/2010 950 INVE Refusal at 32 feet bgs in weathered basalt

Areas of Interest and Soil Boring L	ocations	San	nple Inform	nation				General		Parameters hem ^(e)	GRO &	Analysis	Sample Type	
Soil Boring Easting Production Well NW-9 (continued)	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	<u>pH</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold (b)	(INVE/DUP)	Additional Comments
SB-73 824092.05	389877.34	SB-73 (0-1)	0-1	8/3/2010	1025	Χ	Χ	Χ					INVE	
		SB-73 (1-2)	1-2	8/3/2010		X	Х	Χ	X				INVE	
		SB-73 (4-5)	4-5	8/3/2010		X	Χ	Χ					INVE	
		SB-73 (9-10)	9-10	8/3/2010		X		Χ				Hold	INVE	
		SB-73 (14-15)	14-15	8/3/2010	1048	X		Χ				Hold	INVE	
		SB-73 (19-20)	19-20	8/3/2010	1052	X		Χ				Hold	INVE	
		SB-73 (24-25)	24-25	8/3/2010	1055	X		Χ				Hold	INVE	
		SB-73 (29-30)	29-30	8/3/2010	1057	X		Χ				Hold	INVE	
		SB-73 (31.5-32.5)	31.5-32.5	8/3/2010	1100	Χ		X				Hold	INVE	Refusal at 32.5 feet bgs in basalt
SB-74 823880.04	389922.60	SB-74 (0-1)	0-1	8/3/2010		Χ	Χ	X					INVE	
		SB-74 (1-2)	1-2	8/3/2010		X	X	X	X	Х			INVE	
		SB-174 (1-2)	1-2	8/3/2010		X	X	X					DUP	
		SB-74 (4-5)	4-5	8/3/2010		X	Χ	X					INVE	
		SB-74 (9-10)	9-10	8/3/2010		X		X				Hold	INVE	
		SB-74 (14-15)	14-15	8/3/2010		X		X				Hold	INVE	
		SB-74 (19-20)	19-20	8/3/2010		X		X				Hold	INVE	
		SB-74 (24-25)	24-25	8/3/2010 8/3/2010		X X		X X				Hold	INVE INVE	Refusal at 30 feet bgs in basalt
		SB-74 (29-30)	29-30	6/3/2010	1152	^		^				Hold	IINV⊏	Refusal at 30 feet bys III basait
QC Laboratory Leach Field														
	389968.12	SB-75 (0-1)	0-1	8/3/2010	1500	Χ	Χ	X					INVE	
OB 10 023114.50	000000.12	SB-75 (1-2)	1-2	8/3/2010		X	X	X	X				INVE	
		SB-75 (4-5)	4-5	8/3/2010		X	X	X	,,				INVE	
		SB-175 (4-5)	4-5	8/3/2010		X	X	X					DUP	
		SB-75 (9-10)	9-10	8/3/2010		X		X				Hold	INVE	
		SB-75 (14-15)	14-15	8/3/2010		X		Χ				Hold	INVE	
		SB-75 (19-20)	19-20	8/3/2010		Χ		X				Hold		Refusal at 20 feet bgs in basalt
SB-76 823753.31	389954.43	SB-76 (0-1)	0-1	8/3/2010	1416	Х	Х	Χ					INVE	
02 70 020 000		SB-76 (1-2)	1-2	8/3/2010		X	X	X	Χ				INVE	
		SB-76 (4-5)	4-5	8/3/2010		X	X	X					INVE	
		SB-76 (9-10)	9-10	8/3/2010		X		Χ				Hold	INVE	
		SB-76 (14-15)	14-15	8/3/2010		X		Χ				Hold	INVE	
		SB-76 (19-20)		8/3/2010		X		Χ				Hold	INVE	
		SB-76 (21.5-22.5)		8/3/2010		Χ	Χ	X				Hold		Refusal at 22.5 feet bgs in weathered basalt
SB-77 823778.32	389922.05	SB-77 (0-1)	0-1	8/3/2010	1329	X	Х	Χ					INVE	
		SB-77 (1-2)	1-2	8/3/2010		X	Χ	Χ	Χ	Χ			INVE	
		SB-77 (4-5)	4-5	8/3/2010		X	Χ	Χ					INVE	
		SB-77 (9-10)	9-10	8/3/2010		X		X				Hold	INVE	
		SB-77 (14-15)	14-15	8/3/2010		X		X				Hold	INVE	
		SB-77 (19-20)	19-20	8/3/2010		Χ		Χ				Hold	INVE	
		SB-77 (24-25)	24-25	8/3/2010	1347	Х		X				Hold	INVE	Refusal at 25 feet bgs in basalt

Areas of Interest a	nd Soil I	Boring L	ocations	San	nple Inforr	mation				General	Chemical Rad Cl	Parameters nem ^(e)	GRO &	Analysis	Sample Type	
Soil Bor Gypstack Area Decant ditch	ing <u>E</u> a	<u>asting</u>	Northing	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (c)	<u>рН</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	Additional Comments
	-78 820	0804.97	393922.18	SB-78 (0-1)	0-1	8/11/2010	950	Χ	Χ	X					INVE	
				SB-178 (0-1)	0-1	8/11/2010		Χ	Χ	X					DUP	
				SB-78 (1-2)	1-2	8/11/2010		Χ	Χ	Χ	X	Χ			INVE	
				SB-78 (4-5)	4-5	8/11/2010		Χ	Х	X					INVE	
				SB-78 (9-10)	9-10	8/11/2010				X				Hold	INVE	
				SB-78 (14-15)	14-15	8/11/2010				X				Hold	INVE	D (1 100 (11 1 1 1 1 1 1 1 1 1 1 1 1 1
				SB-78 (19-20)	19-20	8/11/2010	1005			Χ				Hold	INVE	Refusal at 20 feet bgs in basalt
SB	-79 820	0806.68	392908.01	SB-79 (0-1)	0-1	8/11/2010		Χ	Χ	Χ					INVE	
				SB-79 (1-2)	1-2	8/11/2010		Χ	Х	X	X				INVE	
				SB-79 (4-5)	4-5	8/11/2010		X	Χ	X					INVE	
				SB-79 (9-10)	9-10	8/11/2010				X				Hold	INVE	
				SB-79 (14-15)	14-15	8/11/2010				X				Hold	INVE	Defined at 47 feet has in heart
				SB-79 (16-17)	16-17	8/11/2010	1051			Χ				Hold	INVE	Refusal at 17 feet bgs in basalt
SB	-80 820	0812.97	392149.01	SB-80 (0-1)	0-1	8/11/2010	1102	Χ	Χ	X					INVE	
				SB-80 (1-2)	1-2	8/11/2010		Χ	Χ	Χ	X				INVE	
				SB-80 (4-5)	4-5	8/11/2010		Χ	Χ	X					INVE	
				SB-80 (9-10)	9-10	8/11/2010		X		X				Hold	INVE	_ ,
				SB-80 (13-14)	13-14	8/11/2010	1115	Χ		Χ				Hold	INVE	Refusal at 14.5 feet bgs in Tufa
SB	-81 820	0818.39	390900.38	SB-81 (0-1)	0-1	8/11/2010	1138	X	Χ	Χ					INVE	
				SB-81 (1-2)	1-2	8/11/2010	1139	Χ	Χ	Χ	Χ				INVE	
				SB-81 (4-5)	4-5	8/11/2010		Χ	Χ	Χ					INVE	
				SB-81 (9-10)	9-10	8/11/2010		X		X				Hold	INVE	
				SB-81 (14-15)	14-15	8/11/2010		X		X				Hold	INVE	
				SB-81 (19-20)	19-20	8/11/2010		X		X X				Hold	INVE	Defined at 00 feet has in heart
				SB-81 (22-23)	22-23	8/11/2010	1200	Χ		*				Hold	INVE	Refusal at 23 feet bgs in basalt
East Gypstack Area	00 000	2000 10	004050 40	05 00 (0.4)	0.4	0/44/0040	1010		v	V					IN 175	
SB	-82 823	3203.19	394658.49	SB-82 (0-1)	0-1	8/11/2010		X	X	X	V	V			INVE	
				SB-82 (1-2) SB-182 (1-2)	1-2 1-2	8/11/2010 8/11/2010		X X	X X	X X	Х	X			INVE DUP	
				SB-82 (4-5)	4-5	8/11/2010		X	X	X					INVE	
				SB-82 (9-10)	9-10	8/11/2010		X	^	X				Hold	INVE	
				SB-82 (14-15)	14-15	8/11/2010		X		X				Hold	INVE	
				SB-82 (19-20)	19-20	8/11/2010		X		X				Hold	INVE	Refusal at 21 feet bgs in basalt
Q.P.	-83 823	3194 87	393362.79	SB-83 (0-1)	0-1	8/11/2010	1429	Х	Х	Х					INVE	
OL	-00 020	J13 4 .01	333302.73	SB-83 (1-2)	1-2	8/11/2010		X	X	X	Χ				INVE	
				SB-83 (4-5)	4-5	8/11/2010		X	X	X	Α				INVE	
				SB-83 (5-6)	5-6	8/11/2010								Hold	INVE	Wet from 6 to 9.5 feet bgs, Refusal at 10 feet bgs in tufa
SB	-84 822	2367.63	390463.51	SB-84 (0-1)	0-1	8/11/2010	1505	Х	Х	Χ					INVE	
	J. ULL		200.00.01	SB-84 (1-2)	1-2	8/11/2010		X	X	X	Χ				INVE	
				SB-84 (4-5)	4-5	8/11/2010		X	X	X	-				INVE	
				SB-84 (9-10)		8/11/2010		Χ		Χ				Hold	INVE	Refusal at 10 feet bgs in basalt

Soil Sample Summary - 2010 Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Areas of Interest and Soil Boring I	Locations	San	nple Inforn	nation				General	Chemical Rad Ch	Parameters nem ^(e)	GRO &	Analysis	Sample Type	
<u>Soil Boring</u> <u>Easting</u> Cooling Pond Area	<u>Northing</u>	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals ^(c)	<u>Hq</u>	Chemistry (d)	Gross a & B	Ra-226/228	BTEX (f)	on Hold ^(b)	(INVE/DUP)	Additional Comments
SB-85 821855.03	389809.78	SB-85 (0-1) SB-85 (1-2) SB-85 (4-5) SB-85 (8-9)	0-1 1-2 4-5 8-9	8/11/2010 8/11/2010 8/11/2010	1542 1544 1547	X X X	X X X	X X X	Х			Hold	INVE INVE INVE INVE	Refusal at 9 feet bgs in tufa
SB-86 820700.00	388900.00	SB-186 (0-1) SB-186 (1-2) SB-186 (4-5) SB-86 (0-1) SB-86 (1-2) SB-86 (4-5) SB-86 (9-10) SB-86 (14-15)	0-1 1-2 4-5 0-1 1-2 4-5 9-10 14-15	8/12/2010 8/12/2010 8/12/2010 8/12/2010 8/12/2010 8/12/2010 8/12/2010 8/12/2010	732 735 815 817 819 825 830	X X X X X X	X X X X X	X X X X X X	x x	x x		Hold Hold	DUP DUP DUP INVE INVE INVE INVE	
SB-87 820985.27	388229.81	SB-86 (18-19) SB-87 (0-1) SB-87 (1-2) SB-87 (4-5) SB-87 (9-10) SB-87 (12-13)	0-1 1-2 4-5 9-10 10-11	8/12/2010 8/12/2010 8/12/2010 8/12/2010 8/12/2010 8/12/2010	930 931 932 933	X X X X X	X X X	X X X X X	Х			Hold Hold Hold Hold	INVE INVE INVE INVE INVE	Refusal at 19 feet bgs in weathered basalt Refusal at 11 feet bgs in weathered basalt
Additional QC Samples Trip Blanks	- - -	TB080410 TB080510 TB080510	- - -	8/4/2010 8/5/2010 8/6/2010	-						X X X		INVE INVE INVE	
Equipment Blanks	- - - - - - - -	EB080410 EB080510 EB080610 EB080910 EB081010 EB081110 EB081210 EB081310 EB081610 EB081810	-	8/4/2010 8/5/2010 8/6/2010 8/9/2010 8/10/2010 8/11/2010 8/13/2010 8/16/2010 8/17/2010 8/18/2010	915 1100 1615 1050 1300 1045 935 1750 1115	X X X X X X X X	X X X X X X X X X X	X X X X X X X X	X X X X X X X	x x	X X X		INVE INVE INVE INVE INVE INVE INVE INVE	Collected from deconned Geoprobe cutting shoe

a/ Oxidation Tank Farm includes Tank #11, Tank #11A, Tank #25, Tank #25A, Tank #38, Tank #41, and Tank #50. Central Tank Farm includes Tank #12, Tank #13, Tank #18, Tank #20, Tank #21, Tank #23, Tank #24, Tank #24A, and Tank #47.

Bold = Above criteria

WSP Environment & Energy 14 of 14

b/ Samples were collected from 0-1 ft bgs; 1-2 ft bgs, and 4-5 ft bgs; followed by every 5 feet until saturation or refusal at bedrock. In general, samples deeper than 5 feet were submitted to the laboratory on hold pending the initial analysis. Deeper samples were subsequently analyzed for analytes requiring vertical delineation. Borings adjacent to sumps, lagoons, or ponds had deeper samples submitted initially.

c/ Metals include list presented in Table 2.

d/ General chemistry parameters include total phosphorus, nitrate, Total Kjeldahl Nitrogen, ammonia, and fluoride.

e/ Radiological analysis includes gross alpha and gross beta radiation for samples from 0-1 ft bgs. 25% of these samples were analyzed for radium-226 and radium-228.

f/ BTEX = Benzene, toluene, ethylbenzene, and xylenes.

Table D-2

Analyte Test Methods and Regulatory Criteria Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

		Method	Laboratory	EPA Soil RSLs ^(c)	Idaho Dep of Environmen		EPA Region 9 RSL Protection of	Background				Sample	Criteria ^(e)	
	Test		Reporting	Industrial	Risk Based	Critical	Groundwater (e)	Concentrations	Comparative	Comparative		Quantity		Holding
<u>Analytes</u>	Method (b)	<u>Limit</u>	<u>Limit</u>	<u>Soil</u>	IDTLs	Pathway	Risk-Based/MCL-Based	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	<u>Basis</u>	Container	(grams)	<u>Preservative</u>	_
Metals (mg/kg)														
Aluminum	SW-846 6010C	1.2	10	99,000	- (g)	NS	23,000/NS	15,041	23,000	GW RSL	G	100	none	180 days
Antimony	SW-846 6010C	0.1	1	41	4.8	GWP	0.27/0.27	0.50	0.50	background	G	100	none	180 days
Arsenic	SW-846 6010C	0.1	0.5	1.6	0.39	Surficial Soil	0.0013/0.29	4.7	4.7	background	G	100	none	180 days
Barium	SW-846 6010C	0.5	10	19,000	896	GWP	1,200/82	170.2	170.2	background	G	100	none	180 days
Beryllium	SW-846 6010C	0.05	0.25	200	1.6	GWP	13/3.2	0.89	1.63	IDTL	G	100	none	180 days
Cadmium	SW-846 6010C	0.05	0.2	80	1.4	GWP	NS/NS	0.869	1.4	IDTL	G	100	none	180 days
Calcium	SW-846 6010C	5	250	(j)	NS	NS	NS/NS	38,270 (I)	38,270	background	G	100	none	180 days
Chromium, Total	SW-846 6010C	0.05	0.5	15,0000	2,135	GWP	NS/180,000	18.61	2,135	IDTL	G	100	none	180 days
Iron	SW-846 6010C	1.7	15	72,000	5.76	GWP	270/NS	14,811	14,811	background	G	100	none	180 days
Lead	SW-846 6010C	0.05	1	80	50	GWP	NS/14	13.59	14	GW RSL	G	100	none	180 days
Magnesium	SW-846 6010C	5	250	(j)	NS	NS	NS/NS	7,146	'7,146	background	G	100	none	180 days
Manganese	SW-846 6010C	0.05	0.75	2,300	223	G	NS/NS	742 (I)	742	background	G	100	none	180 days
Nickel	SW-846 6010C	0.05	2	2,000	59	GWP	20/NS	15.71	20	GW RSL	G	100	none	180 days
Potassium	SW-846 6010C	25	500	(j)	NS	NS	NS/NS	3,393 (I)	3,393	background	G	100	none	180 days
Selenium	SW-846 6010C	0.2	1	510	2.0	GWP	0.4/0.26	1.04	1.04	background	G	100	none	180 days
Sodium	SW-846 6010C	55	500	(j)	NS	NS	NS/NS	289 (I)	289	background	G	100	none	180 days
Thallium	SW-846 6010C	0.13	0.5	1	1.6	GWP	0.0011/0.14	NS	0.5	RĽ (k)	G	100	none	180 days
Vanadium	SW-846 6010C	0.05	2.5	520	NS	NS	78/NS	22.68	78	GW RSL	G	100	none	180 days
General Chemistry (mg/kg))													
Fluoride	EPA 9056A	1	0.5	4,100	7.4	GWP	NS	3.95	7	IDTL	G	100	4°C	28 days
pH (S.U.)	SW-846 9045	0.01	0.01	NS	NS	NS	NS	NS	NS	NS	G	100	4°C	ASAP
Total Phosphorous	EPA 365.3	6.7	3.35	NS	NS	NS	NS	NS	NS	NS	G	100	4°C	28 days
Total Kjeldahl Nitrogen	EPA 351.2	11	4.2	NS	NS	NS	NS	NS	NS	NS	G	100	4°C	28 days
Nitrate	EPA 9056A	1	0.5	1,600,000	18.4	GWP	NS	NS	NS	NS	G	100	4°C	48 hours
Ammonia	EPA 350.1	1.2	0.6	NS	4.1	Subsurface Soil		NS	NS	NS	G	100	4°C	28 days
Radiological Parameters (p	oCi/g) ^(c)													
Gross alpha	EPA 900	3 (m)	NS	NS	NS	NS	NS	6.21	NS	NS	G or Poly	30	4°C	28 days
Gross beta	EPA 900	4 (m)	NS	NS	NS	NS	NS	5.52	NS	NS	G or Poly	30	4°C	28 days
Radium-226 - surface E	PA 901/EPA 903.1	1 (m)	NS	5	NS	NS	NS	1.958	1.958	background	G or Poly	250	4°C	28 days
Radium-226 - subsurface E	PA 901/EPA 903.1	1 (m)	NS	15	NS	NS	NS	1.958	1.958	background	G or Poly	250	4°C	28 days
	EPA 901/EPA 904.0	1 (m)	NS	15	NS	NS	NS	1.756	1.756	background	G or Poly	250	4°C	28 days

Analyte Test Methods and Regulatory Criteria Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

"m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

"NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85. Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

c/ Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/

- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ Background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV
- h/ Provided by Accutest Laboratories.
- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects

Soil Sampling Analytical Results - 2010 Ore Processing Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve		epartment ental Quality ^(b)	Dealerraind		SB-01 INVE	SB-01 INVE	SB-01 INVE	SB-02 INVE	SB-02 INVE	SB-02 DUP	SB-02 INVE	SB-02 INVE	SB-02 INVE
Sample Type: Sampling Date:	Industrial	Protection of Groundwater ^(e)	Risk Based	Critical	Background Concentrations	Comparative		08/13/10	08/13/10	08/12/10	08/12/10	08/12/10	08/12/10	1NVE 08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	0-1	1-2	1-2	4-5	9-10	11.5-12.5
Dopun.	con (mg/ng)	Mon Bacca, mon Bacca		. atmay	(0.2000)	valuoo									1110 1210
Metals (mg/kg)															
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1.4 U	3 U	1.9 ∪	3.2 ∪	2 U	-	4.4 U	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	5.3 J	6 J	1.9 U	5.5 J	5.8 J	-	9.8 J	4.7	1.7
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	121 J	132 J	98.7 J	174 J	172 J	-	71.1 J	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.76 J	1.5 U	0.95 U	1.6 ∪	1 U	-	2.2 U	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	16.3	42.3	1.5 J	43.6	10.3	-	132	0.33 J	0.93 J
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	64,300	110,000	12,500	119,000	37,400	-	263,000	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	87.7	226	42.9	190	79.4	-	642	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	9.7	10.7	11	8.3	15.2	-	13.4	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	7,260	4,900 J	4,010 J	7,040 J	5,500	-	1,670 J	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	35.1	57.2	26.4	46.2	29.1	-	116	23.2	14.7
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,610 J	3,180 J	2,690 J	2,650 J	3,670 J	-	2,500 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	2.9 U	6 U	3.8 U	6.3 ∪	4.1 U	-	8.7 U	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	540 U	1,100 U	710 U	1,740 J	770 U	-	5,210 J	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.25 J	0.86	0.095 U	0.32 J	0.1 U	-	3.3	0.13 L	J 0.13 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	156	402	32.9 J	370	130	-	1,370	28	21.6
Wet Chemistry (mg/kg)	NS	NC	4.4	Subsurface Soil	NS	NC	0	7.4	0.00 1	0.0	0.6 U		0.0	2.0	4.7
Ammonia (as N)	4,100	NS NS			3.95	NS 7	8 19.4	7.1 12.9	0.99 J	2.6		-	8.2	3.9	1.7
Fluoride	4,100 1,600,000	NS NS	7.4 18.4	GWP GWP		, NS			0.58 U	20.8	403	-	1.6	-	-
Nitrate as N				_	NS NC	NS NS	22.8 780	6 994	6.7 700	6.2 771	4.7 911	-	2.5	-	-
Total Kjeldahl Nitrogen	NS	NS NS	NS	NS	NS NC	NS NS					• • •	-	922	-	-
Total Phosphorus	NS		NS	NS	NS		519	2,810	1,000	2,560	7,060	-	8,410	-	-
рН	NS	NS	NS	NS	NS	NS	7.2	7.28	6.68	7.43	7.5	-	7.73	-	-
Radiological (pCi/g)															
Gross alpha	NS	NS	NS	NS	NS	NS	-	27.4	-	-	17.1	22.4	-	-	-
Gross beta	NS	NS	NS	NS	NS	NS	-	20.7	-	-	13.4	17.8	-	-	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	6.3	-	-	-	-	-	-	-
Radium 228	15	NS	NS	NS	NS	15	-	0.85	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Soil Sampling Analytical Results - 2010 Ore Processing Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-03 INVE	SB-03 INVE	SB-03 INVE	SB-03 INVE	SB-03 INVE	SB-04 INVE	SB-04 INVE	SB-04 INVE	SB-04 DUP
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base		Pathway	(UTL 95-95) ^(f)	<u>Values (g)</u>	0-1	1-2	4-5	9-10	12-13	0-1	1-2	4-5	4-5
	(<u>,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,</u>										
Metals (mg/kg)															
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1 U	0.89 ∪	0.9 ∪	-	-	3 U	3.3 U	0.96 ∪	1.1 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	6.3	5.8	2.7 J	3	12.5	6.4 J	12.9 J	2.6 J	2.9
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	46 J	54.6 J	130	-	-	143 J	99 J	142	143
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.51 U	0.48 J	0.73 J	-	-	1.5 U	1.6 U	0.63 J	0.71
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	5.6	4.6	1.9	0.17 U	0.73	19.2	77.6	1.5 J	2.3
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	175,000	23,000	25,400	-	-	95,700	198,000	61,000	35,200
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	48.3	115	27.4	-	-	104	464	20.6	26.3
Lead	80	NS/14	50	GWP	NS/14	13.59	3.2	4.7	9.8	-	-	6.9	128	7.9	9.5
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	3,130	2,990	5,910	-	-	8,790	3,790 J	8,540	5,590
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	18.7	41.4	30.3	-	-	30.4	101	17.4	22.2
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	935 J	1,490 J	2,950 J	-	-	2,840 J	3,940 J	2,210 J	2,830
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	2 U	4.8	1.8 U	-	-	6 U	9 J	1.9 ∪	2.2 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	380 U	330 U	351 J	-	-	1,570 J	2,220 J	360 U	410 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.1 U	0.23 J	0.099 J	-	-	0.074 U	1.7	0.096 U	0.11 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	51.6	51.3	49.7	-	-	165	777	25	29.6
Mot Chamiatry (malka)															
Wet Chemistry (mg/kg) Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	5.8	15.4	54.7	2.5	3	11.6	8.6	18.3	46.2
Fluoride	4,100	NS	7.4	GWP	3.95	7	31.5	41.3	21.3	1.1 J	0.62 U	87.7	21.4	2.7	4.8
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	2.5	2.4	4.4	-	-	404	965	18.9	17.4
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	171	711	1,190	_	_	7.76	7.65	650	1,030
Total Phosphorus	NS	NS	NS	NS	NS	NS	859	519	883			7.70	7.00	1,280	5,160
рН	NS	NS	NS	NS	NS	NS	8	7.82	7.36	-	-	-	-	7.65	7.63
•															
Radiological (pCi/g)															
Gross alpha	NS	NS	NS	NS	NS	NS	-	88	-	-	-	-	88	-	-
Gross beta	NS	NS	NS	NS	NS	NS	-	63	-	-	-	-	54.1	-	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	-	-	-	-	-	-	-	-
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Soil Sampling Analytical Results - 2010 Ore Processing Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-04 INVE	SB-04 INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative Values ^(g)	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	a iDTES (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>values ""</u>	9-10	12-13
Metals (mg/kg)								
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.1	2.9
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.32 J	0.9
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	-	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	-
Wet Chemistry (mg/kg)								
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	1.5	1.7
Fluoride	4,100	NS	7.4	GWP	3.95	7	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	5	4.5
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	-	-
рН	NS	NS	NS	NS	NS	NS	-	-
Radiological (pCi/g)								
Gross alpha	NS	NS	NS	NS	NS	NS	-	-
Gross beta	NS	NS	NS	NS	NS	NS	-	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	-
Radium 228	15	NS	NS	NS	NS	15	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-05 INVE	SB-06 INVE	SB-06 INVE	SB-06 INVE						
Sample Type: Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative		08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	14-15	19-20	21-22	0-1	1-2	4-5
Volatile Organic Compur	nds (ma/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	0.0012 U	0.00094 U	0.00093 U	-	-	-	-	0.0014 U	0.0011 U	0.0013 U
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	0.0013 U	0.001 U	0.001 U	-	-	-	-	0.0015 U	0.0013 U	0.0014 U
Toluene	45,000	0.69	4.89	GWP	NS	NS	0.0014 U	0.0011 U	0.0011 U	-	-	-	-	0.0017 U	0.0014 U	0.0015 U
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	3.1 U	2.8 U	2.6 U	-	-	-	-	3.9 U	3.4 U	3.8 U
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	0.0025 U	0.0021 U	0.002 U	-	-	-	-	0.003 U	0.0025 U	0.0028 U
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.48 U	0.44 U	0.43 U	_	-	-	-	0.54 U	2 U	0.11 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3	4.8	4.8	4.7	3.2 J	4.6	10.8	3.6	7.7 J	1.9
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	71 J	118	115	-	-	-	-	108 J	146 J	195
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.48 U	0.74 J	0.71 J	-	-	-	-	0.54 U	1 J	0.56 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	3.6	3.8	4.6	3.9	3.6	0.73 J	1.1 J	9.4	48.2	21.3
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	40,100	45,000	40,100	-	-	-	-	86,900	115,000	2,200 J
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	29	45.2	30.9	-	-	-	-	94.4	282	132
Lead	80	NS/14	50	GWP	NS/14	13.59	5.5	9.5	9	-	-	-	-	11.9	14.9	15.1
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	8,960	6,950	7,680	-	-	-	-	2,320 J	2,100 J	156 J
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	16.9	30	28.3	-	-	-	-	17.3	29.9	1 J
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,450 J	2,430 J	2,400 J	-	-	-	-	1,940 J	4,420 J	4,540
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.9 ∪	4 J	2.4 J	-	-	-	-	2.2 U	4.1 J	0.59 J
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	360 U	330 U	461 J	-	-	-	-	400 U	1,180 J	420 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.096 U	0.88 ∪	0.87 ∪	-	-	-	-	0.11 U	0.77 J	0.79
Vanadium	520	78/NS	NS	NS	78/NS	22.68	44.7	56.1	43.1	-	-	-	-	109	416	163
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	35.4	19.1	27.1	73.4	9.6	17.1	11.2	104	148	450
Fluoride	4,100	NS	7.4	GWP	3.95	7	18	11	6	-	-	-	-	14.2	38.9	4.8
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	3.8	4.9	2.4	-	-	-	-	2.4	0.57 U	0.62 U
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	812	448	280	-	-	-	-	753	947	1,870
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,180	2,740	4,670	-	-	-	-	3,230	7,260	2,880
рН	NS	NS	NS	NS	NS	NS	7.84	7.75	7.83	-	-	-	-	6.94	6.55	3.49
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	10.4	-	-	-	-	-	-	53	-
Gross beta	NS	NS	NS	NS	NS	NS	-	7.3	-	-	-	-	-	-	35.5	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	-	-	-	-	-	-	-	-	-
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-06 INVE	SB-06 INVE	SB-06 INVE	SB-07 INVE	SB-07 DUP	SB-07 INVE	SB-07 INVE	SB-07 INVE	SB-07 INVE	SB-07 INVE
Sample Type: Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative		08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10	08/06/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	9-10	14-15	18-19	0-1	0-1	1-2	4-5	9-10	14-15	18-19
	(.															
Volatile Organic Compur	nds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	-	-	-	0.0014 U	0.0011 U	0.0013 U	0.001 U	-	-	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	-	-	-	0.0015 U	0.0012 U	0.0014 U	0.0011 U	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	-	-	-	0.0016 U	0.0013 U	0.0015 U	0.0012 U	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	-	-	-	3.5 U	3.6 U	4 U	2.9 U	-	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	-	-	-	0.003 U	0.0024 U	0.0028 U	0.0022 U	-	-	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	_	_	_	1.6 U	0.44 ∪	0.9 U	0.44 U	_	_	_
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.2	3.3 J	4.1	5	4.2	3.1 J	3.5	2.5	2.9	4.3
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2		-	-	130	128	130 J	105			-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	_	_	_	0.38 J	0.65	0.9 U	0.68 J	_	_	_
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.46	1.8	0.56	2.5	4.3	10.7	1.3	_	_	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	-	-	103,000	19,000	76,000	18,600	_	_	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	_	-	-	340	51.9	139	22.6	-	_	-
Lead	80	NS/14	50	GWP	NS/14	13.59	6.9	12.7	10.4	8.9	11.3	40.3	10.5	_	_	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	-	-	8,220	4,170	2,150 J	5,140	_	_	_
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	_	-	-	26.4	25.9	9.4	21.2	_	_	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	_	-	-	1,340 J	2,800	2,230 J	2,960 J	_	_	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	_	-	-	1.8 J	1	1.8 ∪	0.88 J	-	_	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	_	-	-	1,100 J	330 U	680 U	330 U	-	_	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.1 U	0.079 U	0.12 U	0.31 ∪	0.088 U	0.14 J	0.87 ∪	-	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	-	-	72.7	51.1	133	25.5	-	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	17.5	8.1	10.7	38.2	44.8	294	119	6.7	6	28.1
Fluoride	4,100	NS	7.4	GWP	3.95	7	-	-	-	14.1	17.6	11.8	15.2	1.5	5.7	11.2
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	_	_	_	12.2	3.3	0.6 U	0.59 U	-	-	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	-	-	295	518	673	1,070	_	_	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	_	-	-	2,430	5,370	6,470	995	_	_	-
рН	NS	NS	NS	NS	NS	NS	-	-	-	6.97	7.00	4.94	7.15	-	-	-
Padialogical (nCi/a)																
Radiological (pCi/g) Gross alpha	NS	NS	NC	NC	NS	NS						36				
Gross alpha Gross beta	NS NS	NS NS	NS NS	NS NS	NS NS	NS NS	-	-	-	-	-	21.3	-	-	-	-
Radium 226	5 ^(surface) . 15	NS NS	NS NS	NS NS	NS NS	5 ^(surface) . 15	-	-	-	-	-	۷۱.3	-	-	-	-
	5 ^(surface) , 15	NS NS	NS NS	NS NS	NS NS	5 (surface), 15	-	_	-	-	-	-	-	-		-
Radium 228	15	INO	1/10	ONI	INO	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Soil Sampling Analytical Results - 2010 Sulfuric Acid and PPA Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type: Sampling Date:	Region 9 Regi	onal Screening Leve Protection of Groundwater ^(e)		epartment ental Quality ^(b) Critical	Background Concentrations	Comparative	SB-08 INVE 08/04/10	SB-08 INVE 08/04/10	SB-08 DUP 08/04/10	SB-08 INVE 08/04/10	SB-08 INVE 08/04/10	SB-08 INVE 08/04/10	SB-08 INVE 08/04/10	SB-09 INVE 08/05/10	SB-09 INVE 08/05/10	SB-09 DUP 08/05/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base			(UTL 95-95) ^(f)	Values (g)	0-1	1-2	1-2	4-5	9-10	14-15	19-20	0-1	1-2	1-2
Volatile Organic Compur	nds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	0.001 U	0.0011 U	-	0.00067 U	-	-	-	0.00094 U	0.0011 U	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	0.0011 U	0.0012 U	-	0.00074 U	-	-	-	0.001 U	0.0012 U	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	0.0012 U	0.0013 U	-	0.00081 U	-	-	-	0.0011 U	0.0013 U	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	2.7 U	2.9 U	-	2.1 U	-	-	-	2.5 U	2.5 U	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	0.0022 U	0.0023 U	-	0.0015 U	-	-	-	0.0021 U	0.0023 U	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.74 ∪	0.44 U	-	0.37 U	-	-	-	1.1 U	0.99 ∪	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.6 J	5	-	3.2	3.3 J	6.4	4.3	5.1 J	3.7 J	-
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	74.8	88.9	-	66.9	-	-	-	102 J	75.7 J	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.37 U	0.52 J	-	0.33 J	-	-	-	0.64 J	0.49 U	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	4.3	4.2	-	3.8	6	3.4	1.6	4.6	2.6	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	28,700	46,500	-	59,500	-	-	-	39,400	29,000	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	29.6	33.6	-	29.5	-	-	-	41.5	22.6	-
Lead	80	NS/14	50	GWP	NS/14	13.59	7	8	-	6.9	-	-	-	10.6	5	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	7,140	11,400	-	25,000	-	-	-	11,700	7,640	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	21.3	26.6	-	19.2	-	-	-	30	17.8	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,660 J	1,870 J	-	1,310 J	-	-	-	2,110 J	1,330 J	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.5 ∪	2 J	-	1.3 J	-	-	-	2.1 U	2 U	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	280 U	170 U	-	140 U	-	-	-	400 U	370 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.074 U	0.088 U	-	0.074 U	-	-	-	0.11 U	0.099 U	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	75.7	48.3	-	37.2	-	-	-	53.8	37.4	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	9.4	9.7	-	35.3	39.6	399	771	20	21.4	-
Fluoride	4,100	NS	7.4	GWP	3.95	7	7.8	4.5	-	2.6	-	-	-	6.1	12.5	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	13.9	7.5	-	0.56 U	-	-	-	0.55 U	0.56 U	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	84.7	111	-	357	-	-	-	400	323	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	416	2,160	-	3,280	-	-	-	2,000	2,130	-
рН	NS	NS	NS	NS	NS	NS			-	7.63	-	-	-	8.01	8.06	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	10.5	11.9	-	-	-	-	-	6.7	12.9
Gross beta	NS	NS	NS	NS	NS	NS	-	7.2	7	-	-	-	-	-	5.7	8.1
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	2.75	3.68	-	-	-	-	-	3.14	2.82
Radium 228	15	NS	NS	NS	NS	15	-	1.23	1.04	-	-	-	-	-	1.03	0.75

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho D	epartment	Dooleanound		SB-09 INVE	SB-09	SB-09	SB-09	SB-09 INVE	SB-10 INVE	SB-10 INVE	SB-10 INVE	SB-10 INVE	SB-10 INVE
Sample Type: Sampling Date:	Industrial	Protection of Groundwater ^(e)	Of Environm	ental Quality (b) Critical	Background Concentrations	Comparative	1NVE 08/05/10	DUP 08/05/10	INVE 08/05/10	INVE 08/05/10	08/05/10	1NVE 08/04/10	08/04/10	08/04/10	08/04/10	1NVE 08/04/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based			(UTL 95-95) ^(f)	Values (g)	4-5	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10	14-15
2004	con (mg/ng/				(0:=0000)											
Volatile Organic Compur	nds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	0.00082 U	U 8000.0	-	-	-	-	-	-	-	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	0.0009 U	0.00088 U	-	-	-	-	-	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	0.00099 U	0.00096 U	-	-	-	-	-	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	2.5 U	2.6 U	-	-	-	-	-	-	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	0.0018 U	0.0018 U	-	-	-	-	-	-	-	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.97 ∪	0.91 ∪	_	_	_	1 U	0.8 U	0.84 ∪	_	_
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.8 J	2.5	5.1	6.1	4.3	3.4 J	4.1	3.4 J	4.4 J	4.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	96.8 J	91.9	-	-	-	57.2 J	152	126	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.83 J	0.74	_	_	_	0.52 U	0.4 U	0.42 U	_	_
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	1.4 J	0.99	_	_	-	1.5 J	6.7	6.6	1.4 J	0.85
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	6,630	5,800	_	_	-	254,000	24,300	21,800	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	30.9	29.1	-	-	-	76.5	61.5	79.2	_	_
Lead	80	NS/14	50	GWP	NS/14	13.59	11.8	11.4	-	-	-	3.5	13.6	20.9	13.8	10.5
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	3,970	3,330	-	-	-	5,730	7,530	5,240	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	19.3	16.5	-	-	-	15	37.4	23.6	-	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,630 J	2,560	-	-	-	710 J	3,080 J	2,390 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.9 U	1.8 U	-	-	-	2.1 U	1.6 U	1.7 U	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	360 U	340 U	-	-	-	585 J	1,530 J	1,210 J	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.097 U	0.091 U	-	-	-	2.1 U	0.8 U	0.084 U	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	26.3	25.4	-	-	-	39.3	101	90.5	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	59.8	50.6	263	66.9	89.5	33.6	49.5	75	10.9	4.5
Fluoride	4,100	NS	7.4	GWP	3.95	7	11.7	17.7	2.1	1.4	1.4	24.6	23.7	18.9	8.5	3.7
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	9.4	9.6	-	-	-	14.8	4.8	10	-	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	765	782	-	-	-	223	512	485	_	_
Total Phosphorus	NS	NS	NS	NS	NS	NS	1,060	801	-	-	-	6,240	2,370	1,500	_	_
рН	NS	NS	NS	NS	NS	NS	6.86	7.06	-	-	-	7.36	7	6.98	-	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS							8.3			
Gross alpha Gross beta	NS NS	NS	NS NS	NS NS	NS NS	NS NS	-	-	-	-	-	-	o.s 5	-	-	-
Radium 226	5 ^(surface) . 15	NS	NS NS	NS	NS NS	5 ^(surface) . 15	_	_	-	_	-	_	J -	_	_	-
Radium 228	5 (********, 15	NS NS	NS NS	NS	NS NS	5 (, 15 15	_		-	-	-	-	-	-	-	-
radium ZZU	10	140	140	140	140	10	-	-	=	_	_	_		-	=	_

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID:	Region 9 Regi	onal Screening Leve Protection of		epartment	Dooleanound		SB-11	SB-11	SB-11	SB-11	SB-11	SB-11	SB-12	SB-12	SB-12	SB-12
Sample Type: Sampling Date:	Industrial	Groundwater (e)	Risk Based	ental Quality ^(b) Critical	Background Concentrations	Comparative	INVE 08/04/10	DUP 08/04/10	INVE 08/04/10							
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	0-1	1-2	4-5	9-10	12.5-13.5	00/04/10	1-2	4-5	9-10
	(3,3)				,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,											
Volatile Organic Compur	nds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	-	-	-	-	-	-	-	-	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	-	-	-	-	-	-	-	-	-	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.69 ∪	0.87 ∪	0.84 ∪	0.81 U	_	_	0.43 U	0.42 U	0.44 U	_
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	6	3.7	4.7	3.2 J	5.4	2.4 J	7.2	2 J	2.7	3
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	149	197	153	168	-	-	203	220	92.7	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.42 J	0.46	0.45 J	1.4 J	_	_	0.3 J	0.21 U	0.66 J	_
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	5.9	5.7	8.6	5.2	11.1	0.92 J	5.4	0.97	0.32 J	_
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	36,500	41,400	41,700	13,200	-	-	30,100	14,300	5,300	_
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	61.4	67.7	102	248	-	-	115	27.3	25.5	-
Lead	80	NS/14	50	GWP	NS/14	13.59	27.5	19.3	35.3	9.9	-	-	66.7	4.7	11.2	_
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	8,740	12,800	7,270	3,300	_	_	9,450	14,100	3,840	_
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	24.8	34.7	27	9.4	-	-	91.1	32.1	13.7	_
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,020 J	2,620	2,320 J	4,930	-	-	2,640	3,350	2,390	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	2 J	1.7 U	2.3 J	1.6 ∪	-	-	2.2 J	1.2 J	0.88 ∪	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	2,280 J	3,140	2110 J	871 J	-	-	1,530 J	2,770	170 U	-
Thallium	ĺ	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.69 ∪	0.87 ∪	0.84 ∪	0.081 U	-	-	0.087 U	0.85 ∪	0.088 U	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	103	114	135	424	36.2	29.6	198	213	24.3	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	66.6	75.8	32.1	333	25.2	23.1	49.7	10.7	93.1	9.1
Fluoride	4,100	NS	7.4	GWP	3.95	7	39.7	41	41.7	295	4.3	46.7	15	4.5	2	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	53.4	88	19.6	78.6	43.9	34.1	8.7	2.8	4.3	_
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	405	387	355	1,280	-	-	576	165	843	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,030	464	6,530	1,620	-	-	6,540	2,090	499	_
pH	NS	NS	NS	NS	NS	NS	5.14	5.09	5.19	4.7	-	-	7.29	7.29	7.32	-
Padialogical (nCi/a)																
Radiological (pCi/g) Gross alpha	NS	NS	NS	NS	NS	NS		_	17.7					1.04		
Gross alpha Gross beta	NS NS	NS NS	NS NS	NS NS	NS NS	NS NS	-	-	17.7 17.7	-	-	-	-	1.04	-	-
Radium 226	5 ^(surface) . 15	NS	NS NS	NS NS	NS NS	5 ^(surface) . 15	-	-	17.7	-	-	-	-	1.90	-	-
Radium 228	5 (sunass), 15 15	NS NS	NS NS	NS NS	NS NS	5 (surface), 15	-	-	-	-	-	-	-	-	-	-
Naululli 220	19	INO	INO	INO	INO	10	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-12 INVE	SB-13 INVE	SB-13 INVE	SB-13 INVE	SB-13 INVE	SB-13 INVE	SB-14 INVE	SB-14 INVE	SB-14 INVE	SB-14 INVE
Sampling Date: Depth:	Industrial Soil (mg/kg)	Groundwater ^(e) Risk-Based/MCL-Base	Risk Based d IDTLs (mg/kg)	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative Values (g)	08/04/10 14-15	08/04/10 <u>0-1</u>	08/04/10 1-2	08/04/10 4-5	08/04/10 9-10	08/04/10 14-15	08/04/10 <u>0-1</u>	08/04/10 1-2	08/04/10 4-5	08/04/10 9-10
Volatile Organic Compur	nds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	-	-	-	-	-	-	-	-	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	-	-	-	-	-	-	-	-	-	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	0.91 ∪	0.37 U	0.082 U	-	-	0.77 ∪	1 U	0.57 U	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.6	5.9	3.6	3.2	4.6	2.2	3.3 J	7.9	9.4	4.4
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	110	273	192	-	-	75.5 J	225	197	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	0.45 U	0.19 U	0.041 U	-	-	0.38 U	0.52 U	0.66 J	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	-	6.6	1.7	0.31	-	-	5.4	4.9	0.42 J	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	89,700	38,200	33,200	-	-	58,900	30,100	10,600	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	76.6	31.3	9.4	-	-	108	86	452	-
Lead	80	NS/14	50	GWP	NS/14	13.59	-	14.8	183	9.8	-	-	5.1 J	66.9	12.4	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	41,700	4,980	204	-	-	26,900	8,570	5,180	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	-	35.9	10.5	0.57 J	-	-	40	26.6	17.7	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	1,550 J	1,700 J	1,820	-	-	992 J	2,090 J	3,280	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	1.8 U	1 J	0.66 ∪	-	-	1.5 ∪	2.1 U	1.1 U	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	2,400 J	2,410	31 U	-	-	1,680 J	2,790 J	220 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	1.4 J	0.074 U	0.082 U	-	-	0.77 ∪	0.1 U	0.65	0.32 J
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	83.8	50.6	13.3	-	-	69.3	87.1	115	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	4.2	23.4	21.7	27.4	231	510	23.2	10.4	19.6	588
Fluoride	4,100	NS	7.4	GWP	3.95	7	-	43.2	17.9	6.7	-	-	20.9	19.3	10.8	5.3
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	-	27.3	22.9	17.2	-	-	17.9	5.1	3.7	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	318	283	311	-	-	282	268	607	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	-	853	4,960	682	-	-	3,390	3,320	2,500	-
рН	NS	NS	NS	NS	NS	NS	-	7.09	4.84	5.57	-	-	7.1	7.81	3.82	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	-	8.6	-	-	-	-	6.6	-	-
Gross beta	NS	NS	NS	NS	NS	NS	-	-	8.4	-	-	-	-	6.3	-	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	-	2.34	-	-	-	-	-	-	-
Radium 228	15	NS	NS	NS	NS	15	-	-	0.67 U	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID:	Region 9 Regi	ional Screening Leve		epartment ental Quality ^(b)	Daalamaa d		SB-14	SB-15	SB-15	SB-15	SB-15	SB-15	SB-15	SB-15	SB-15	SB-16
Sample Type:	ا ماد، مدیناما	Protection of Groundwater ^(e)			Background	Commonative	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial Soil (mg/kg)	Risk-Based/MCL-Base	Risk Based	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative Values (g)	08/04/10	08/04/10	08/04/10	08/04/10	08/04/10	08/04/10	08/04/10	08/04/10	08/04/10	08/02/10
Depth:	Soli (ilig/kg)	RISK-Based/WCL-Base	u ibits (ilig/kg)	Patriway	(UIL 95-95)**	<u>values " </u>	14-15	0-1	1-2	4-5	9-10	14-15	19-20	24-25	26-27	<u>0-1</u>
Volatile Organic Compu	nds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	-	-	-	-	-	-	-	-	_	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	-	-	-	-	-	-	-	_	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	-	-	-	-	-	-	-	-	-	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	0.82 ∪	0.98 ∪	0.97 ∪	-	-	-	_	_	0.51 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	6.0	4.6	5.8	7.6	4.4	5.1 J	3.2	3.5	3.3	5.3
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	98.6	326	200	-	-	-	-	-	96.6
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	0.41 U	0.49 U	1.3 J	-	-	-	_	_	0.43 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	-	6.1	4	3.9	0.52 J	0.8 U	1.1	1.2	1.2	4.2
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	92,100	50,000	41,400	-	-	-	-	_	73,900
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	54.3	452	25.3	-	-	-	-	-	33.7
Lead	80	NS/14	50	GWP	NS/14	13.59	-	4.8 J	21.7	11.7	-	-	-	-	_	6.3
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	36,400	9,730	5,120	-	-	-	-	_	8,390
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	-	28.8	25.6	39.8	-	-	-	-	-	20.8
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	1,930 J	3,240 J	2,530 J	-	-	-	-	-	2,550 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	1.6 U	2 U	1.9 ∪	-	-	-	_	-	1 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	1,070 J	2,570 J	360 U	-	-	-	_	-	264 J
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.24	0.82 ∪	1.6 J	0.97 ∪	-	-	-	-	-	1 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	71.9	139	43.7	-	-	-	-	-	41.9
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	184	11.7	15.3	24.1	271	105	12.4	7	2.6	34.9
Fluoride	4,100	NS	7.4	GWP	3.95	7	1.8	20.7	9.8	58.1	4.9	9.4	188	682	546	139
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	-	15.7	10	12.6	-	-	-	-	-	8.5
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	308	209	585	-	-	-	-	-	378
Total Phosphorus	NS	NS	NS	NS	NS	NS	-	2,650	2,720	1,150	-	-	-	-	-	2,160
рН	NS	NS	NS	NS	NS	NS	-	7.55	6.51	7.06	-	-	-	-	-	6.51
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	-	6.5	-	-	-	-	-	-	-
Gross beta	NS	NS	NS	NS	NS	NS	-	-	17.8	-	-	-	-	-	-	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	-	-	-	-	-	-	-	-	-
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID: Sample Type:	Region 9 Reg	ional Screening Leve Protection of	Idaho D	epartment ental Quality ^(b)	Background		SB-16 INVE	SB-16 INVE	SB-16 INVE	SB-17 INVE	SB-17 INVE	SB-17 INVE	SB-17 INVE	SB-17 INVE	SB-18 INVE	SB-18 DUP
Sampling Date: Depth:	Industrial Soil (mg/kg)	Groundwater ^(e) Risk-Based/MCL-Base	Risk Based	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative Values (g)		08/02/10 4-5	08/02/10 8-9	08/02/10 0-1	08/02/10 1-2	08/02/10 4-5	08/02/10 9-10	08/02/10 13-14	08/02/10 0-1	08/02/10 0-1
	- (g/g/		···· ··· ··· ··· ··· ··· ··· ··· ··· ·	.	10:20007								<u> </u>			
Volatile Organic Compu	ınds (mg/l)															
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	-	-	-	-	-	-	-	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	-	-	-	-	-	-	-	-	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	-	-	-	-	-	-	-	-	-	-
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	18 U	0.46 U	_	0.44 U	0.072 U	0.087 U	_	_	0.53 U	0.96 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	32.4	3.7	1.8 J	4.1	4.7	3.3	4.7	2.8	2.9	3.5
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	71.5	125	-	107	12.8	9.6	_	-	59	70.2
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.35 J	0.63 J	_	0.7 J	0.075 J	0.074 J	_	_	0.39 J	0.48
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	2.3	1	7.2	2.4	0.24	0.044 U	-	_	2.1	4.3
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	27,700	18,700	-	22,300	4,490	2,750	_	_	67,100	70,900
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	18.5	52.8	-	32.4	5.3	4.6	-	-	24.8	34.3
Lead	80	NS/14	50	GWP	NS/14	13.59	4.6	9.8	-	10.2	1.1	1	-	_	4.8	5.1
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	9,090	5,050	-	4,800	761	624	-	_	11,900	17,500
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	12.4	17.1	-	21.4	4.3	1.9	-	-	16.6	19.9
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,780 J	2,550	-	2,200	234 J	215 J	-	-	1,290 J	1,530
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	23.9	0.92 ∪	-	0.87 ∪	0.14 U	0.17 U	-	-	1.1 U	1.9 ∪
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	170 U	170 U	-	160 U	27 U	33 U	-	-	200 U	360 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.091 U	0.092 U	-	0.087 U	0.072 U	0.087 U	-	-	0.11 U	0.096 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	25.2	38	-	41.4	8.1	4.1	-	-	32.1	43.7
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	12.6	63.7	220	41.7	9.7	2.5	-	_	3.9	5.6
Fluoride	4,100	NS	7.4	GWP	3.95	7	10.2	23.2	178	9.5	2.4	1.6	_	-	5.2	1.9
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	7.2	4.1	-	9	3.2	1.7	-	_	1.6	1.3
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	199	841	-	651	221	29.4	-	-	193	219
Total Phosphorus	NS	NS	NS	NS	NS	NS	639	269	-	577	1,080	194	-	_	1,210	1,920
рН	NS	NS	NS	NS	NS	NS	6.96	6.65	-	6.98	7.04	7.57			7.45	7.36
Radiological (pCi/g)														_		
Gross alpha	NS	NS	NS	NS	NS	NS	6	_	_	_	2.99	_	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	4.6	_	_	_	2.4				_	_
Radium 226	5 ^(surface) , 15	NS	NS	NS	NS	5 ^(surface) , 15	-	_	_	_		_	_	_	_	_
Radium 228	15	NS	NS	NS	NS	15	_	-	-	-	_	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Soil Sampling Analytical Results - 2010 Sulfuric Acid and PPA Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Region	onal Screening Leve Protection of	Idaho De	epartment ental Quality ^(b)	Background		SB-18 INVE	SB-18 INVE	SB-18 INVE
Sampling Date: Depth:	Industrial Soil (mg/kg)	Groundwater ^(e) Risk-Based/MCL-Based	Risk Based	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative Values (g)		08/02/10 4-5	08/02/10 8.5-9.5
•	•		, ,	•					
Volatile Organic Compune									
Benzene	5.4	0.00021	0.0178	GWP	NS	NS	-	-	-
Ethylbenzene	27	0.0017	10.2	GWP	NS	NS	-	-	-
Toluene	45,000	0.69	4.89	GWP	NS	NS	-	-	-
TPH-GRO (C6-C10)	NS	NS	NS	NS	NS	NS	-	-	-
Xylene, (total)	2,700	0.2	1.67	Subsurface Soil	NS	NS	-	-	-
Metals (mg/kg)									
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.52 U	0.58 U	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	4.3	4.1	6.7
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	116	129	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.71 J	0.75 J	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	3.4	0.31 J	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	31,400	8,500	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	38	20.6	-
Lead	80	NS/14	50	GWP	NS/14	13.59	9.8	10.5	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	8,300	5,520	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	22.4	18.5	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,370 J	2,230 J	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.2 J	1.2 U	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	190 U	220 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.1 U	0.12 U	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	50.9	28.6	-
Wet Chemistry (mg/kg)									
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	37.6	6.2	-
Fluoride	4,100	NS	7.4	GWP	3.95	7	8.1	0.92	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	1.2	7.6	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	585	218	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,390	2,170	-
рН	NS	NS	NS	NS	NS	NS	7.29	7.51	-
Radiological (pCi/g)									
Gross alpha	NS	NS	NS	NS	NS	NS	7.7	-	-
Gross beta	NS	NS	NS	NS	NS	NS	5.7	-	-
Radium 226	5 (surface), 15	NS	NS	NS	NS	5 (surface), 15	-	-	-
Radium 228	15	NS	NS	NS	NS	15	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve		epartment ental Quality ^(b)	Da alama and		SB-19 INVE	SB-19 INVE	SB-19 INVE	SB-19 INVE	SB-19 INVE	SB-20 INVE	SB-20 INVE	SB-20 INVE	SB-20 INVE	SB-20 INVE
Sampling Date:	Industrial	Protection of Groundwater ^(e)	of Environme Risk Based	Critical	Background Concentrations	Comparative		08/09/10	08/09/10	08/09/10	08/09/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base		Pathway	(UTL 95-95) ^(f)	Comparative Values (g)	0-1	1-2	4-5	9-10	14-15	0-1	1-2	4-5	9-10	13-14
Берин.	Soli (Ilig/kg)	RISK-Daseu/IVICL-Dase	ed IDTES (IIIg/kg)	Falliway	(OIL 93-93)**	values	<u> </u>	1-2		9-10	14-13		1-2	 3	3-10	13-14
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1.8 U	2 U	0.88 ∪	-	-	0.82 ∪	1.9 ∪	0.46 U	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	4.9 J	3.2 J	5.2	0.85	2.2	4.9	6.6 J	3.4	5	4.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	114 J	85.2 J	199	-	-	147	149 J	141	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.91 U	1 U	0.97 J	-	-	0.82 U	2.4 U	0.69 J	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	10.7	21.6	0.44 U	-	-	14.4	29.4	0.64 J	-	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	92,500	125,000	31,200	-	-	95,800	164,000	11,000	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	73.1	114	23.1	-	-	63.8	173	18.2	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	9.6	7.7	11.1	-	-	11.6	12.6 J	10	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	11,100	9,900	7,700	-	-	15,000	8,930 J	4,320	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	24	34.3	16.3	-	-	33.3	46.5	14.9	-	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,460 J	2,100 J	4,020 J	-	-	2,580	3,850 J	3,520 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	3.6 ∪	4 U	1.8 ∪	-	-	4.2	8.4 J	0.92 ∪	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,050 J	753 J	330 U	-	-	2,280	1,800 U	430 U	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.091 U	0.1 U	0.088 U	-	-	0.082 U	1.9 ∪	0.12 U	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	116	182	32.6	-	-	118	259	21.9	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	15.5	10.6	5.5	35	5.2	253	22.7	44	4.8	2.5
Fluoride	4,100	NS	7.4	GWP	3.95	7	99.8	26.2	2.3	-	-	28.2	44.9	1.7	_	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	7.7	8.2	16.2	-	_	19.7	11.2	13.3	-	_
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	322	348	283	-	_	1,080	561	1,130	-	_
Total Phosphorus	NS	NS	NS	NS	NS	NS	6,620	7,900	934	-	_	1,890	3,130	632	_	-
рН	NS	NS	NS	NS	NS	NS	6.99	7.25	7.53	-	-	6.34	6.93	7.53	-	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	28.6	_	_	_	_	38.8	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	25	_	_	_	_	27.9	_	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	_	10.6	_	_	_	_		_	_	_
Tagiani ZZO	(subsurface)	NS	NS	NS	NS	(subsurface)		10.0								
Radium 228	15	NS	NS	NS	NS	15	-	0.59 U	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

 $a/\ mg/kg = milligrams\ per\ kilogram;\ s.u. = standard\ units;\ pCi/g = picocuries\ per\ gram;\ "n"\ indicates\ RSL\ based\ on\ non-carcinogenic\ toxicity;$

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve	<u> </u>				SB-19	SB-19	SB-19	SB-19	SB-19	SB-20	SB-20	SB-20	SB-20	SB-20
Sample Type:		Protection of	of Environmental Quality (b)		Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	14-15	0-1	1-2	4-5	9-10	13-14

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects. The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Table D-5

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Deelemannd		SB-21 INVE	SB-21 INVE	SB-21 INVE	SB-21 DUP	SB-21 INVE	SB-21 INVE	SB-21 INVE	SB-22 INVE	SB-22 INVE	SB-22 INVE
Sampling Date:	Industrial	Groundwater ^(e)	of Environme Risk Based	Critical	Background Concentrations	Comparative		08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	4-5	9-10	14-15	15.5-16.5	0-1	1-2	4-5
Берин.	Son (mg/kg)	RISK-Daseu/WICL-Dase	ed IDTES (IIIg/kg)	Falliway	(OIL 93-93)**	values		1-2			9-10	14-13	13.3-10.3		<u> </u>	
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	3.2 J	1.9 ∪	1.1 U	0.54 ∪	-	-	-	1.6 ∪	0.94 ∪	1 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	7.4 J	6.7 J	4.4 J	5.7	4.7	3.4	3.1	4.7 J	10.7	7.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	145 J	113 J	143	180	-	-	-	60.7 J	12.9 J	118
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.94 U	0.96 U	0.74 J	0.73	-	-	-	0.8 U	0.47 U	0.52 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	28.6	41.2	0.57 U	0.44	-	-	-	9.6	15.6	12.8
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	118,000	124,000	18,800	35,000	-	-	-	69,700	153,000	55,500
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	218	205	25.8	19.9	-	-	-	90	66.3	72.1
Lead	80	NS/14	50	GWP	NS/14	13.59	72.4	12.8	10.9	10.1	-	-	-	6.5	2.3	11.5
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	5,300	6,680	5,620	6,890	-	-	-	9,820	85,500	14,700
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	23	56.8	16	15.8	-	-	-	56.9	33.4	53.7
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	5,280 J	3,320 J	3,140 J	2,720	-	-	-	1,420 J	642 J	2,410 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	4.4 J	3.9 J	2.3 U	1.1 U	-	-	-	3.7 J	2.8 J	2.1 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	2,840 J	1,670 J	430 U	200 U	-	-	-	600 U	382 J	390 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.094 U	0.096 U	0.11 U	0.11 U	-	-	-	0.08 U	0.094 U	1.3 J
Vanadium	520	78/NS	NS	NS	78/NS	22.68	238	306	32.2	29.1	-	-	-	69.1	71.8	79.6
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	112	17.2	67.9	59.1	4.5	4	2.4	23.1	3.4	68.1
Fluoride	4,100	NS	7.4	GWP	3.95	7	144	26.3	0.78 J	2.7	-	-	-	20.2	111	20.5
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	5	16.6	19.2	34.8	5.3	7	6.2	9.6	3.7	0.59 U
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	349	318	793	686	-	-	-	804	176	599
Total Phosphorus	NS	NS	NS	NS	NS	NS	13,100	5,980	602	1110	-	-	_	5,000	626	3,010
рН	NS	NS	NS	NS	NS	NS	6.49	6.77	7.44	7.40	-	-	-	4.70	5.09	4.79
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	39.4	_	_	_	_	_	_	13.8	_
Gross beta	NS	NS	NS	NS	NS	NS	_	34.1	_	_	_	_	_	_	9.7	_
Radium 226	(f)	140	140	110	110	5 ^(surface) , 15	_	34.1	-	_	_	_	_	_	9.7	_
Nadium 220	5 ^(surface) , 15 (subsurface)	NC	NC	NC	NC	5 ', 15 (subsurface)	-	-	-	-	_	_	_	_	-	_
B. II. 200	,	NS	NS	NS	NS	,										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathwa GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve Idaho Department				SB-21	SB-21	SB-21	SB-21	SB-21	SB-21	SB-21	SB-22	SB-22	SB-22	
Sample Type:		Protection of	of Environmen	ital Quality (b)	Background		INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	4-5	9-10	14-15	15.5-16.5	0-1	1-2	4-5

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluories

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-5

Sample ID:	Region 9 Regi	ional Screening Leve		epartment			SB-22	SB-22	SB-22	SB-23	SB-23	SB-23	SB-24	SB-24	SB-24	SB-24
Sample Type:		Protection of		ental Quality (b)	Background	_	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative		08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	9-10	14-15	15.5-16.5	0-1	1-2	2.5-3.5	0-1	1-2	4-5	6-7
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	-	-	1.7 U	2 U	0.66 ∪	1 U	2 U	2.6 ∪	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.4	0.89	2.1	7.1 J	6.3 J	1.5 J	2.9 J	3 J	2.6 ∪	-
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	-	-	189	125 J	97.8	72.9 J	32.8 J	462	203 J
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	-	-	0.86 U	0.98 U	0.33 U	0.52 U	1 U	1.3 U	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	1.5	1.6	4.9	17.5	20.3	1.4	15.5	17.1	5.9	19.4
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	-	-	94,400	144,000	130,000	72,000	27,600	2,690 J	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	-	-	162	138	40.4	51.9	108	550	-
Lead	80	NS/14	50	GWP	NS/14	13.59	-	-	-	66.1	28.6	3.8	6.1	3.4	19.2	10.9
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	-	-	5,810	4,600 J	77.8 J	24,600	11,000	307 J	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	23.7	4.1	22.2	22.1	15.7	1.7 J	35.2	15.8	6	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	-	-	4,960 J	2,890 J	386 J	1,640 J	1,120 J	5,110 J	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	-	-	3.4 U	6.1 J	3.8 J	2.1 U	4 U	5.1 U	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	-	-	3,220 J	1,350 J	250 U	390 U	750 U	960 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	-	-	0.086 U	0.098 U	0.14 J	0.1 U	0.31 J	3.3	0.64
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	-	-	165	168	27	49.7	173	130	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	3.2	6.3	12	80.2	97.1	59.2	8.8	13.3	175	2.5
Fluoride	4,100	NS	7.4	GWP	3.95	7	3.3	817	161	435	147	388	168	56.9	1,690	245
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	_	_	_	1.8	0.6 U	0.7 U	13.5	4.2	0.7 U	
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	_	-	_	309	385	119	353	82.3	1,520	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	_	-	_	6,350	2,870	3,680	1,750	1,850	2,680	-
рН	NS	NS	NS	NS	NS	NS	-	-	-	4.82	5.32	2.72	5.04	5.73	2.67	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	_	62	_	_	8.3	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	_	_	_	36.4	_	_	5.37	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	_	_	_	_	-	_	_	7.2	_	_
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	_	_	-	-	_	_	4.52	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve					SB-22	SB-22	SB-22	SB-23	SB-23	SB-23	SB-24	SB-24	SB-24	SB-24
Sample Type:		Protection of	of Environmen	ital Quality (b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	9-10	14-15	15.5-16.5	0-1	1-2	2.5-3.5	0-1	1-2	4-5	6-7

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorion

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- i/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

WSP Environment & Energy
Page 6 of 16

Table D-5

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve		epartment ental Quality ^(b)	Dooleanound		SB-26 INVE	SB-26 INVE	SB-26 INVE	SB-26 INVE	SB-27 INVE	SB-27 DUP	SB-27 INVE	SB-27 DUP	SB-27 INVE	SB-27 INVE
Sampling Date:	Industrial	Protection of	Risk Based	Critical	Background Concentrations	Commonative	08/18/10	08/18/10	08/18/10	08/18/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10
Depth:		Groundwater (e)			(UTL 95-95) ^(f)	Comparative Values (g)	0-1	1-2	4-5	7-8	0-1	06/09/10	1-2	1-2	4-5	8-9
Бериі.	Soil (mg/kg)	Risk-Based/MCL-Based	IDILS (mg/kg)	Pathway	(UIL 95-95)\^	<u>values " </u>		1-2	4-5	<u> </u>			1-2	1-2	4-5	0-9
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.73 ∪	2.4 J	7.6	1.7 J	2 U	1.5 ∪	3.3 U	-	2.5 U	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	4.7	3.5	6.2	3.7 J	6.8 J	8.7	3.5 J	-	5.5 J	2.0 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	84.6 J	63.6 J	140	315	173 J	164	99.2 J	-	375	204
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.73 U	0.49 J	1.2 J	1.2 U	1 U	0.76 U	1.6 U	-	1.2 U	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	22.1	3.2	2.3	186	35.8	82	20.7	-	3.5 J	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	101,000	1,510 J	1,830 J	3,310 J	107,000	93,500	128,000	-	12,300	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	89	222	555	524	233	173	514	-	260	-
Lead	80	NS/14	50	GWP	NS/14	13.59	6.7 J	3.9	7.2	15.6	52	87.4	3.5 J	-	34	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	40,000	68.1 J	504 J	864 J	5,690	5,240	3,160 J	-	185 J	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	30.7	0.18 J	1.2 J	3.1	33.4 J	33.6	16.3 J	-	4.8	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,640 J	2,030	4,440 J	5,430 J	7,090 J	7,130	8,040 J	-	3,270 J	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.5 ∪	1.1 J	2 J	2.3 ∪	6 J	7.7	6.5 ∪	-	5.8 J	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	894 J	340 U	460 U	870 U	5,490	4,700	2,390 J	-	930 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.73 ∪	10.9	24	3	3.3 J	1.9	2.8 J	-	1.6	1.4
Vanadium	520	78/NS	NS	NS	78/NS	22.68	130	420	2,410	1,210	279	250	605	-	51 J	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	30.2	430	1,110	313	114	160	104	_	117	_
Fluoride	4,100	NS	7.4	GWP	3.95	7	21.6	12.7	34.8	40.3	136	113	154	_	991	285
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	13.4	30.5	93.4	18.2	17.7	2.6	18.1	_	11.4	
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	222	1,810	1,280	1,400	518	438	420	_	1,190	_
Total Phosphorus	NS	NS	NS	NS	NS	NS	10,300	4,060	4,920	14,400	7,300	4,700	7,100	_	9,490	_
рН	NS	NS	NS	NS	NS	NS	6.53	4.78	3.32	3.90	5.88	5.57	6.34	-	3.44	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS		19.6	_		_	_	85	86		_
Gross alpha Gross beta	NS NS	NS NS	NS NS	NS NS	NS	NS NS	-	7.7	-	_	-	-	68	63	-	-
Radium 226	5 ^(surface) , 15	NO	INO	INO	INO	(f)	-	7.7	-	-	-	-	00	-	-	-
Naululli 220	5 (subsurface)	NO	NO	NO	110	5 (subsurface), 15	-	-	-	-	-	-	-	-	-	-
		NS	NS	NS	NS											
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathwa GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	gional Screening Leve Idaho Department		artment			SB-26	SB-26	SB-26	SB-26	SB-27	SB-27	SB-27	SB-27	SB-27	SB-27
Sample Type:		Protection of	of Environmen	ıtal Quality ^(b)	Background		INVE	INVE	INVE	INVE	INVE	DUP	INVE	DUP	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/18/10	08/18/10	08/18/10	08/18/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	7-8	0-1	0-1	1-2	1-2	4-5	8-9

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorion

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-5

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Daakanaanad		SB-28 INVE	SB-28 INVE	SB-28 INVE	SB-28 DUP	SB-29 INVE	SB-29 INVE	SB-29 INVE	SB-29 INVE	SB-30 INVE	SB-30 INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Background Concentrations	Comparative		08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	3.5-4.5	1-2	0-1	1-2	4-5	8-9	0-1	1-2
	Jon (mg/kg)	Nisk-Dased/WCL-Dase	ed IDTES (IIIg/kg)	1 attiway	(OTE 33-33)	<u>values</u>										<u></u>
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.58 U	1.8 U	2.4 U	-	0.4 U	1.6 ∪	0.4 U	-	1.7 U	1.6 ∪
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.3 J	3.3 J	3.3 J	-	1.5 J	4.3 J	6.8	1.7	4.4 J	4.8 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	298	256	350	-	275	157 J	152	-	140 J	137 J
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.58 U	0.96 J	1.2 U	-	0.4 U	1.3 J	2.4	1.5	0.84 U	0.8 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	7.3	20.1	8.9	-	1.3	48.1	104	168	9.2	12.6
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	72,400	82,700	6,490	-	36,900	99,000	16,000	-	127,000	44,300
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	84.8	183	265	-	3.1 J	188	65.5	-	90.6	97.8
Lead	80	NS/14	50	GWP	NS/14	13.59	8.1	8.5	21.5	-	5.3	38.3	9	-	20.5	13.4
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	11,800	2,000 J	304 J	-	12,200	4,700	2,730	-	28,900	4,680
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	8.7 J	12.3	11.2	-	6.1 J	18.4	26.6	-	17.8 J	19.4
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	5,000 J	8,300 J	3,510 J	-	3,950 J	5,170 J	7,740	-	1,620 J	3,020 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	3.6 ∪	4.8 ∪	-	1.4 J	3.6 J	1.6 ∪	-	3.4 U	3.2 ∪
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	5,780 J	2,890 J	900 U	-	3,930 J	3,180 J	1,450 J	-	630 U	600 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.58 ∪	0.49 J	2	-	0.4 U	0.79 ∪	0.81 ∪	-	1.7 U	0.08 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	170	366	79.8	-	97	344	241	177	90.7	137
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	66.9	118	192	-	2.6	172	256	170	34.9	124
Fluoride	4,100	NS	7.4	GWP	3.95	7	493	463	1,220	-	42.8	272	322	43.5	18.5	11.6
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	7.5	14.1	0.69 U	-	2.4	7.7	1.2	-	9.8	6.8
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	276	624	1,380	-	63.8	321	1,330	_	324	486
Total Phosphorus	NS	NS	NS	NS	NS	NS	3,040	3,080	1,020	-	2,170	6,550	4,680	-	1,090	2,130
рН	NS	NS	NS	NS	NS	NS	5.12	6.46	2.86	-	7.11	5.15	4.71	-	6.26	6.13
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	47.7	_	50.4	_	23.4	_	_	_	1.7 U
Gross beta	NS	NS	NS	NS	NS	NS	_	59.5	_	61.1	_	15.6	_	_	_	2.4 U
Radium 226	5 ^(surface) . 15					5 ^(surface) , 15	_	3.47	_	-	_	10.0	_	_	_	2.40
	(subsurface)	NS	NS	NS	NS	(subsurface)		J								
Radium 228	15	NS	NS NS	NS NS	NS	15	-	0.62 U	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-28	SB-28	SB-28	SB-28	SB-29	SB-29	SB-29	SB-29	SB-30	SB-30
Sample Type:		Protection of	of Environmen	ital Quality (b)	Background		INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	3.5-4.5	1-2	0-1	1-2	4-5	8-9	0-1	1-2

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluories

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- i/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-5

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve		epartment ental Quality ^(b)	Daalamaa a		SB-30 INVE	SB-31 INVE	SB-31 INVE	SB-31 DUP	SB-31 INVE	SB-31 INVE	SB-32 INVE	SB-32 INVE	SB-32 INVE	SB-32 INVE
Sampling Date:	Industrial	Protection of Groundwater ^(e)	or Environme Risk Based	Critical	Background Concentrations	Comparativa		08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10
Depth:		Risk-Based/MCL-Base			(UTL 95-95) ^(f)	Comparative Values ^(g)	4-5	0-1	1-2	1-2	4-5	9-10	0-1	1-2	4-5	7-8
Берин.	Soil (mg/kg)	RISK-Based/IVICL-Base	a ibits (mg/kg)	Pathway	(UIL 95-95)	values		<u> </u>	1-2	1-2	4-5	9-10		1-2	4-5	7-0
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	2.2 ∪	0.081 U	1.1 U	1.1 U	1 J	-	1.7 U	2 U	1.6 ∪	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	5.6 J	2.5	4.6 J	4.9	6	4.3	4.7 J	3.2 J	2.3 J	2.3 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	113 J	76.4	234	133	67.8 J	-	135 J	94.1 J	68.8 J	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	1.1 J	0.23 J	0.93 J	0.97	0.4 U	-	0.85 U	1 U	1.1 J	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	10.9	2.4	32.3	26.8	9.1	24.8	10	40.7	51.2	56.5
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	15,300	182,000	35,200	65,200	102,000	-	180,000	53,700	111,000	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	110	23.2	89.4	108	36.5	-	44	186	228	-
Lead	80	NS/14	50	GWP	NS/14	13.59	9.8	3.4 J	7.7	21.1	65.2	9.4	5.9 J	5.4	23.4	7.2
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	3,060 J	13,600	9,250	11,900	7,210	-	10,200	4,500 J	7,410	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	11.5	16	23.7	30.6	25.7	-	15.2 J	26.7	23.7 J	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	4,700 J	1,000 J	3,240 J	2,230	1,780 J	-	1,930 J	1,410 J	824 J	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	4.4 U	1.6 U	2.2 U	2.5	1.7 J	-	3.4 U	4 U	3.2 ∪	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	830 U	835 J	3,310 J	1,920	599 J	-	1,570 J	750 U	610 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.11 U	0.4 U	0.11 U	0.11 U	0.08 U	-	1.7 U	0.57	1.6 ∪	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	202	37.8	257	247	52.7	-	91.1	301	464	570
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	358	19	74.2	71.5	61	95.1	12.4	91	4.1	5,160
Fluoride	4,100	NS	7.4	GWP	3.95	7	84.1	29	4.8	4.9	19.7	303	13.3	12.9	5.3	<i>,</i> -
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	3.2	6.8	10.4	14.6	2.6	-	4	17.4	5.8	_
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	899	183	123	224	1,170	-	211	325	149	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	1,600	2,740	2,940	6,610	5,610	-	2,790	5,220	10,600	_
рН	NS	NS	NS	NS	NS	NS	4.89	7.21	5.31	5.45	7.41	-	8.03	5.64	6.05	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	25.1	_	_	_	_	56.9	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	_	15.9	_	_	_	_	36.5	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	_	_	-	_	_	_	_	3.01	_	_
	(subsurface)	NS	NS	NS	NS	(subsurface)								0.0.		
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	1.14	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-30	SB-31	SB-31	SB-31	SB-31	SB-31	SB-32	SB-32	SB-32	SB-32
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values ^(g)	4-5	0-1	1-2	1-2	4-5	9-10	0-1	1-2	4-5	7-8

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorion

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-5

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve		epartment			SB-32	SB-33	SB-33	SB-33	SB-34	SB-34	SB-34	SB-34	SB-34	SB-34
Sample Type:		Protection of		ental Quality (b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative		08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	14-15	0-1	1-2	3.5-4	0-1	1-2.5	4-5	9-10	14-15	19-20
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	2.4 J	2.4 J	1 J	0.46 U	4.1 U	0.21 J	0.081 U	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	6.9	5.2	9.8	4.9 J	4.6	7.9 J	3.7	3	5.6	5
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	47.4 J	214	149	152	126 J	166	117	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	0.54 U	1.1 J	1.2 J	0.83 J	2.3 J	0.75 J	0.54 J	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	1.5	19.8	6.8	3.4	3.4	38.3	2.1	0.74	4	0.48 J
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	21,300	6,310	1,440 J	113,000	208,000	46,800	91,900	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	181	718	506	26.5	456	16.8	17.5	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	9.9	71.2	58.1	35.4	8.7	1.8 J	9.1	7.3	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	949 J	888 J	611 J	5,970	5,700 J	5,590	8,150	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	-	5.7	1.9 J	1.3 J	22.7	24.9 J	103	16.4	23.7	17.2
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	2,620 J	7,940	6,490	2,960 J	5,660 J	2,820	1,240 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	2 J	3.6 J	2.1 J	0.92 ∪	8.3 ∪	0.99 J	0.81 ∪	0.94 ∪	1 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	518 J	1,230 J	1,050 J	340 U	1,600 U	180 U	300 U	-	-
Thallium	ĩ	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	3.5	8.4	13.6	0.092 U	1.7 U	0.12 U	0.081 U	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	38	240	641	366	32.7	673	25.6	28.6	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	9.3	495	815	479	46.6	5.4	6.8	12.5	3.3	6.6
Fluoride	4,100	NS	7.4	GWP	3.95	7	-	22.3	43.8	122	6.1	39.5	5.3	1.3	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	_	14.3	3.6	0.68 U	26.4	20.8	59.7	5.8	7.5	7.8
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	_	1,380	1,410	1,280	1,860	180	403	393	-	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	_	3,850	13,600	12,500	421	5,610	19,700	1,270	-	-
рН	NS	NS	NS	NS	NS	NS	-	•	,	,	7.52	7.32	6.62	8.22	-	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	72	_	_	59	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	_	29.8	_	_	37.7	_	_	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	_	_	-	_	_	12.9	_	_	_	_
Townsell LLV	(subsurface)	NS	NS	NS	NS	(subsurface)						12.0				
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	0.42 U	-	-	_	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathwa GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-32	SB-33	SB-33	SB-33	SB-34	SB-34	SB-34	SB-34	SB-34	SB-34
Sample Type:		Protection of	of Environmen	ital Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative	08/05/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	14-15	0-1	1-2	3.5-4	0-1	1-2.5	4-5	9-10	14-15	19-20

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluories

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-34 INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	24-25
	(g,g)				(0.2.00)		
Metals (mg/kg)							
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	1
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-
Lead	80	NS/14	50	GWP	NS/14	13.59	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	23.8
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	0.48 J
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-
Thallium	ĩ	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	_
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-
Wet Chemistry (mg/kg)							
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	3.1
Fluoride	4.100	NS	7.4	GWP	3.95	7	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	12.2
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	_
рН	NS	NS	NS	NS	NS	NS	-
Radiological (pCi/g)							
Gross alpha	NS	NS	NS	NS	NS	NS	_
Gross beta	NS	NS	NS	NS	NS	NS	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	_
	(subsurface)	NS	NS	NS	NS	(subsurface)	
Radium 228	15	NS NS	NS NS	NS NS	NS NS	15	_
							

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Phosphoric Acid Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-34
Sample Type:		Protection of	of Environmen	tal Quality (b)	Background		INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	l IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	24-25

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment	David was seed to		SB-35	SB-35	SB-35	SB-35	SB-35	SB-35	SB-36	SB-36	SB-36	SB-37
Sample Type:	In Investment	Protection of		ental Quality (b)	Background	0	INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	DILS (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	<u>0-1</u>	1-2	4-5	4-5	9-10	13-14	0-1	1-2	4-5	0-1
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1.7 U	0.13 J	0.99 ∪	1.7 ∪	-	-	1.6 ∪	4.1 U	1 U	1.9 ∪
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	5.6 J	2.3	6.6	5.1	2.9	3.6	7 J	12.7 J	3.5 J	3 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	210	71.3	162 J	155	-	-	140 J	158 J	683	155 J
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.87 U	0.19 J	0.99 U	1.1	-	-	2 U	2.1 U	1 U	0.94 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	13.4	0.28	15.8	24.5	0.69	2.9	22.2	23.8	2.5	11.8
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	59,700	38,600	83,900	72,100	-	-	66,900	61,700	3,720 J	79,400
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	46.8	7.2	95.6	185	-	-	84.5	127	149	159
Lead	80	NS/14	50	GWP	NS/14	13.59	8.9	3.5	15.5	18.4	9.3	9.1	15.1	44.1	29.9	11.4
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	11,400	3,810	5,900	5,140	-	-	10,300	5,300 J	809 J	3,770 J
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	24.1	7.2	36.1	30.9	-	-	13.6	24.2	16.8	8.7
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	5,030 J	1,190 J	4,520 J	4,900	-	-	8,680 J	5,180 J	1,340 J	4,680 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	3.5 ∪	0.42 J	2.3 J	3.5 ∪	-	-	3.2 ∪	8.3 ∪	2.2 J	4.3 J
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	2,640 J	110 U	859 J	1850	-	-	3,640 J	3,160 J	760 U	4,500 J
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.087 U	0.076 U	0.16 J	0.18	0.11 U	0.11 U	0.47	0.26 J	2.7	0.94 ∪
Vanadium	520	78/NS	NS	NS	78/NS	22.68	101	12.3	134	234	21.4	23.3	125	166	58.6	147
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	13.5	6.7	35.3	142	138	186	146	147	107	184
Fluoride	4,100	NS	7.4	GWP	3.95	7	154	11.2	32.9	40.5	0.63 L		338	175	1,050	215
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	3.4	2.1	5	5.2	-	_	37.1	55.5	50.2	28.8
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	265	204	735	671	_	-	318	379	1,860	561
Total Phosphorus	NS	NS	NS	NS	NS	NS	5,190	980	1,210	7.740	_	_	3,320	2,980	6,970	5,270
рН	NS	NS	NS	NS	NS	NS	7.07	8.19	6.4	6.55	-	-	5.52	5.67	2.97	7.2
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	14.9	_	_	_	_	_	19.2	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	9.6	-	-	_	-	_	13.8	-	_
Radium 226	5 ^(surface) . 15		.,,	110	110	5 ^(surface) , 15	_	5.89	_	_	_	_	_	10.0	_	_
radiani 220	O , IO (subsurface)	NS	NO	NC	NS	o ` , Io (subsurface)		0.00								
Radium 228	15	NS NS	NS NS	NS NS	NS NS	15	_	0.49 U	_	_	_	_	_	_	_	_
	. •					. •		55 0								

U = Analyte not detected above method detection limit.

WSP Environment & Energy
Page 1 of 8

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-35	SB-35	SB-35	SB-35	SB-35	SB-35	SB-36	SB-36	SB-36	SB-37
Sample Type:		Protection of	of Environmen	tal Quality (b)	Background		INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	c IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	0-1	1-2	4-5	4-5	9-10	13-14	0-1	1-2	4-5	0-1

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Table D-6

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

SB-37 DUP	SB-37 INVE	SB-37 DUP	SB-37 INVE	SB-37 DUP	SB-38 INVE	SB-38 INVE	SB-38 INVE	SB-39 INVE	SB-39 INVE	SB-40 INVE	SB-40 INVE	SB-40 INVE	SB-40 DUP	SB-40 INVE	SB-40 INVE	SB-41 INVE	SB-41 INVE	SB-41 INVE	SB-41 INVE
08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
0-1	1-2	1-2	4-5	4-5	0-1	1-2	4-5	<u>0-1</u>	<u>1-2</u>	0-1	1-2	4-5	4-5	9-10	11.5-12.5	<u>0-1</u>	1-2	4-5	9-10
2.1 U	3.8 ∪	4 U	1.2 U	1.9 ∪	3.8 ∪	7.2 U	0.45 U	0.88	0.9 ∪	3.8 J	2.2 U	1.1 U	1 U	_	_	1.4 U	1.1 U	0.58 J	_
4.7	9.1 J	6.2	7.2	10.8	7.7 J	7.2 U	2.4	2.8	1.5 J	17.8	5.1 J	4 J	4	2.9	5.1	7.6	4.4 J	2.1	3.2
179	298 J	178	399	391	123 J	158 J	92.3	31.4	69.2 J	100	159 J	160	160	-	-	78.4 J	53.5 J	41.4	-
2.7 U	1.9 U	2 U	1.3 J	1.8	1.9 U	3.6 ∪	0.67 J	0.24	0.45 U	0.46 U	1.1 U	0.82 J	0.73	_	-	0.69 U	0.53 U	0.32 J	_
12.8	8.3	8.3	49.6	22.2	12.7	11.7 J	0.58 J	1.4	1.2 J	5.7	24.5	4.9	0.94	1.6	2.4	10.9	7.1	0.55	_
141,000	52,600	192,000	1,890 J	2,580	62,200	191,000	4,700	38,800	22,300	21,500	71,800	18,700	8,540	-	_	73,900	107,000	19,000	-
278	173	119	662	659	257	218	22.4	10.2	29	22.4	159	42.1	22.7	-	-	72	57.3	7.6	-
13.2	14.6	5.9	20.6	22.3	25.8	4.6 J	10.2	10.6	8.7	38.2	21.9	11.8	11.7	_	-	7.6	4.7	6.8	-
2,380	2,770 J	7,910	316 J	214	5,090 J	4,520 J	3,260	2,690	1,710 J	2,050 J	4,860 J	4,950	4,300	-	-	16,400	34,600	2,800	-
3.5	9.1	21.3	6.6	6.3	15.2	47	13	11.6	18	7.4	31.4	25.5	16.9	-	-	35.1	22	6.6	-
4,560	5,750 J	6,610	5,420 J	5,290	5,380 J	5,870 J	3,050 J	648 J	2,010 J	1,260 J	4,300 J	3,840 J	3,650	-	-	1,620 J	1,290 J	378 J	-
4.3 ∪	11.8 J	8.1	5.5 J	5.4	7.6 ∪	14 U	1.3 J	0.15 U	1.8 U	1.9 ∪	4.4 U	2.2 ∪	2 U	-	-	3 J	3 J	0.34 J	-
6,910	3,120 J	3,220	450 U	700 U	2,020 J	2,700 U	170 U	229 J	1,220 J	685 J	1,410 J	410 U	370 U	-	-	650 J	418 J	329 J	-
2.3	0.95 ∪	2 U	3.6	5.1	0.95 ∪	0.72 ∪	0.89 ∪	0.74 ∪	0.9 ∪	0.14 J	0.11 U	0.11 U	0.1 U	-	-	0.069 U	0.11 U	0.09 U	-
194	229	337	411	322	335	268	21.2	14.2	30.4	37.1	297	68	31.4	-	-	93.4	72.7	8.5	-
181	90	117	689	828	199	187	126	10.3	13.6	26.1	41.2	73.5	79.2	6.5	9.1	13.4	6.3	3.5	
218	627	235	1,160	1,310	585	52	0.59 U	51.9	38	211	56.1	11.2	3	45.6	612	35.4	13	13.5	7.4
13.4	24	6.4	3.4	6	7	31	30.3	12.3	22.5	9.4	20	41.4	11.4	4.8	3.7	7.9	8.8	11.5	
569	562	369	2,310	2,420	227	436	1,030	209	268	174	669	2,110	1,800		-	241	401	72.2	_
5,890	6,640	5,610	4,900	12,700	6,910	12,800	1,160	1,040	1,020	1,100	2,890	800	1110	_	_	2,050	2,250	918	_
6.96	5.8	6.83	2.28	2.33	5.93	7.04	7.1	6.74	6.06	6.03	5.6	6.6	6.75	-	-	7.45	7.68	7.3	-
-	53.5	40.8	-	-	-	67	-	-	6	-	46	-	-	-	-	-	10.8	-	-
-	44.2	25.8	-	-	-	44.3	-	-	5.4	-	35.8	-	-	-	-	-	8.3	-	-
-	-	-	-	-	-	-	-	-	9.8	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	4.19	-	-	-	-	-	-	-	-	-	-

1Wi

∌d.

вĿ

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

SB-37	SB-37	SB-37	SB-37	SB-37	SB-38	SB-38	SB-38	SB-39	SB-39	SB-40	SB-40	SB-40	SB-40	SB-40	SB-40	SB-41	SB-41	SB-41	SB-41
DUP	INVE	DUP	INVE	DUP	INVE	DUP	INVE	INVE	INVE	INVE	INVE	INVE							
08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
0-1	1-2	1-2	4-5	4-5	0-1	1-2	4-5	0-1	1-2	0-1	1-2	4-5	4-5	9-10	11.5-12.5	0-1	1-2	4-5	9-10

orid

nar

um

um

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

SB-41 INVE 08/10/10	SB-41 INVE 08/10/10	SB-42 INVE 08/16/10	SB-43 INVE 08/05/10	SB-43 INVE 08/05/10	SB-43 INVE 08/05/10	SB-43 INVE 08/05/10	SB-43 INVE 08/05/10	SB-43 INVE 08/05/10	SB-44 INVE 08/10/10	SB-44 INVE 08/10/10	SB-44 DUP 08/10/10	SB-44 INVE 08/10/10	SB-44 INVE 08/10/10						
14-15	17.5-18.5	0-1	1-2	4-5	9-10	14-15	19-20	24-25	0-1	1-2	4-5	9-10	14-15	17.5-18.5	0-1	1-2	1-2	4-5	9-10
		<u> </u>							·										
_	_	1.8 U	3.3 ∪	2 U	_	_	_	_	1.6 ∪	1.1 U	3.7 ∪	_	_	_	0.5 J	2 U	2.3 ∪	1.7 U	_
7.1	2.6	6.2 J	4.9 J	7.1 J	2.7	4.5	4.6	3.5	5.1 J	6.3	6.3 J	3.7	5.8	3.8	8.1	6.5 J	4.5	4.8 J	4.2
-	-	108 J	118 J	88.1 J		-	-	-	129 J	119	158 J	-	-	-	9.9	144 J	149	132 J	-
_	-	0.91 U	1.7 U	1 U	_	-	_	-	0.79 U	0.88 J	1.9 ∪	-	-	_	0.24	1 U	1.1 U	0.86 U	_
-	-	22.6	19.6	18.8	0.84	0.53 J	0.55 J	0.54	14.2	17.3	25.9	1	1.6 J	1.1	1	8.1	9.9	8.1	13.1
-	-	161,000	192,000	93,700	-	-	-	-	73,200	46,600	185,000	-	-	-	351,000	64,600	49,700	47,000	-
-	-	217	199	136	-	-	-	-	126	61.4	204	-	-	-	10.6	83.2	98.2	110	-
-	-	43.2	8.6	11.6	-	-	-	-	41.1	10.6	10 J	-	-	-	2 J	25.5	13.7	13.9	-
-	-	5,780	3,740 J	3,860 J	-	-	-	-	6,790	7,550	7,470 J	-	-	-	2,600	6,460	6,780	6,740	-
-	-	43.7	34.2	34.9	-	-	-	-	23.8	50.7	46.1 J	-	-	-	8.2	28.7	27.9	28.4	-
-	-	2,110 J	4,260 J	2,470 J	-	-	-	-	2,650 J	2,760 J	5,700 J	-	-	-	289 J	3,250 J	3,370	3,390 J	-
-	-	5 J	6.6 ∪	4 U	-	-	-	-	3.2 ∪	2.3 J	7.5 ∪	-	-	-	3 U	4.1 U	4.5 ∪	3.4 ∪	-
-	-	2,610 J	3,080 J	1,350 J	-	-	-	-	3,300 J	663 J	2,430 J	-	-	-	104 J	1,060 J	850 U	715 J	-
-	-	1.8 U	0.83 ∪	0.1 U	-	-	-	-	0.79 ∪	1.1 U	3.9 J	-	-	-	0.3 ∪	0.1 U	0.11 U	0.086 U	-
-	-	282	259	225	10.9	29.8	32.7	39.3	167	89.2	266	26.6	31.4	26.1	10.1	88.6	102	89.9	-
_	_	66.1	77.2	109	314	431	52.2	54.1	123	56.2	15.6	53.9	4,600	2,180	134	110	29.1	215	246
785	301	38.6	8.4	14.3	1.3	1.8	112	31.5	31.3	6.9	27.4	33.2	50.9	8.9	4.5	27.6	35.8	44.5	44.2
-	-	182	51.7	124	13.4	50.1	32	21.8	27	26.7	24.4	17	20.5	16.8	10.5	58.6	157	4.9	
-	-	543	368	628	-	-	-	-	526	479	392	-	-	-	101	892	1130	877	_
_	-	6,880	6,600	10,800	_	-	-	-	3,230	2,560	813	-	-	-	2,090	2,390	1,220	5,320	_
-	-	7.42	7.57	5.97	-	-	-	-	5.78	6.72	7.91	-	-	-	7.53	7.03	6.8	7.03	-
-	-	-	52.6	-	-	-	-	-	-	22.6	-	-	-	-	-	14.3	-	-	-
-	-	-	36.4	-	-	-	-	-	-	15	-	-	-	-	-	12.9	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
_	_	-	-	-	_	_	_	-	-	-	-	_	-	-	-	_	-	-	-

1Wi

∌d.

вĿ

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

SB-41	SB-41	SB-42	SB-43	SB-43	SB-43	SB-43	SB-43	SB-43	SB-44	SB-44	SB-44	SB-44	SB-44						
INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	DUP	INVE	INVE
08/10/10	08/10/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/05/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
14-15	17.5-18.5	0-1	1-2	4-5	9-10	14-15	19-20	24-25	0-1	1-2	4-5	9-10	14-15	17.5-18.5	0-1	1-2	1-2	4-5	9-10

orid

nar

um

um

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

SB-44 INVE 08/10/10 14-15	SB-44 INVE 08/10/10 16.5-17.5
- 5.2	- 4.6
-	
-	-
1.3 J	3
-	-
-	-
-	-
-	-
-	-
-	- 3 - - - - - -
-	-
-	-
-	-
119 510	154 825
510	
_	- - -
-	-
-	-
<u>-</u>	-
-	-

ìWi

∌d.

вĿ

Soil Sampling Analytical Results - 2010 Central and Oxidation Tank Farm Areas
Nu-West Industries, Inc.
Conda Phosphate Operations Facility
Soda Springs, Idaho

SB-44	SB-44
INVE	INVE
08/10/10	08/10/10
14-15	16.5-17.5

orid

ıs. ıar

um

um

Soil Sampling Analytical Results - 2010 Granulation Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		partment ntal Quality ^(b)	Background		SB-45 INVE	SB-45 INVE	SB-45 INVE	SB-45 INVE	SB-45 INVE	SB-45 INVE	SB-45 INVE	SB-46 INVE	SB-46 INVE	SB-46 INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	14-15	19-20	23-24	0-1	1-2	4-5
Берин.	Soli (Hig/kg)	KISK-Daseu/IVICL-Dase	ed IDTES (IIIg/kg)	Falliway	(012 93-93)	values				3-10	14-13	13-20				
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.48 U	1.7 U	0.4 U	-	-	-	-	0.47 U	2 U	0.87 ∪
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.2	3.9 J	4.9	3.6	6	3.9	3.8	2.2 J	4.3 J	6.4
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	95.5	95 J	165	-	-	-	-	29 J	141 J	150
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.5 J	1.1 J	0.57 J	-	-	-	-	0.26 J	1.7 J	0.74 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	19.9	58.6	0.57 J	-	-	-	-	7.2	42.1	0.52 J
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	99,600	143,000	50,500	-	-	-	-	18,200	50,200	24,300
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	69.8	128	15.3	-	-	-	-	38.7	331	19.9
Lead	80	NS/14	50	GWP	NS/14	13.59	3.8	3.5	8.2	-	-	-	-	2.2	19.7	10.4
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	15,200	4,180 J	6,390	-	-	-	-	1,860	2,590 J	5,120
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	31	50.9	13.5	-	-	-	-	15.5	19.5	14.4
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,650 J	3,290 J	2,250	-	-	-	-	483 J	5,080 J	2,530 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.6 J	3.4 U	1.9 J	0.67 ∪	0.77 J	0.69 ∪	0.46 J	1.2 J	4 J	1.9 J
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,090 J	1,600 J	150 U	-	-	-	-	180 U	1,190 J	330 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.97 ∪	0.86 ∪	0.81 ∪	-	-	-	-	0.095 U	0.099 U	0.87 ∪
Vanadium	520	78/NS	NS	NS	78/NS	22.68	152	163	20.9	-	-	-	-	56	497	29.5
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	1,270	4,170	2,900	4,220	3,110	2,370	1,910	1,040	2,100	873
Fluoride	4,100	NS	7.4	GWP	3.95	7	15.9	16.4	4.6	-,	-	_,=====================================	-	22.3	45.5	2
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	369	70.6	12.3	_	_	_	_	108	58.3	14.8
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,550	3.720	4,180	_	-	-	-	1.120	3.850	1,700
Total Phosphorus	NS	NS	NS	NS	NS	NS	8,570	9.450	5,900	_	_	_	_	14,100	10,900	13,900
рН	NS	NS	NS	NS	NS	NS	11.13	7.51	7.88	-	-	-	-	5.93	5.07	7.59
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS		53.7							34.3	
Gross alpha Gross beta	NS NS	NS NS	NS NS	NS NS	NS NS	NS NS	-	39.6	-	-	<u>-</u>	<u>-</u>	- -	-	34.3 26	-
Radium 226	5 ^(surface) , 15	INO	INO	INO	INO		_	20.7	-	-	<u>-</u>	<u>-</u>	-	_	20	-
Nadiuiii 220	5 (subsurface)	NO		110		5 ^(surface) , 15 (subsurface)	-	20.1	-	-	-	-	-	-		-
B. II	,	NS	NS	NS	NS	,		0.55								
Radium 228	15	NS	NS	NS	NS	15	-	0.52 U	-	-	-	-	-	-		-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Granulation Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Region	onal Screening Leve	Idaho Dep	partment			SB-45	SB-46	SB-46	SB-46						
Sample Type:		Protection of	of Environmen	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	14-15	19-20	23-24	0-1	1-2	4-5

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects. The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

WSP Environment & Energy
Page 2 of 6

Table D-7

Soil Sampling Analytical Results - 2010 Granulation Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-46 DUP	SB-46 INVE	SB-46 INVE	SB-46 INVE	SB-47 INVE	SB-47 DUP	SB-47 INVE	SB-47 INVE	SB-47 INVE	SB-48 INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative	_	08/16/10	08/16/10	08/16/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	4-5	9-10	14-15	15-16	0-1	0-1	1-2	4-5	9-10	0-1
	con (mg/kg)	Misk Basca/MOL Basca	u ibits (ilig/kg)	Tatilway	(012 00 00)	<u>values</u>										
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.78 ∪	-	-	-	2 U	4.2 U		0.63 U	-	1.5 ∪
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	5.5	1.8	3.9	2.8	4.6 J	5.6	25.1 J	3.3	4.6	4.9 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	160	-	-	-	97.3 J	95.8	134 J	150	-	98.5 J
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.76	-	-	-	1.2 J	3.2	6.5 ∪	0.81 J	-	0.79 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.62	-	-	-	51.8	105	13.4 J	0.31 U	-	24.4
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	21,200	-	-	-	35,200	46,200	153,000	16,400	-	27,000
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	20.2	-	-	-	252	810	130	22.4	-	114
Lead	80	NS/14	50	GWP	NS/14	13.59	10.9	-	-	-	12.9	19.3	8.8	9.7	-	9.7
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	5,040	-	-	-	7,680	12,600	2,500 J	11,000	-	6,020
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	15.4	-	-	-	64.2	138	41.8	70.3	14.2	36.1
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,510	-	-	-	1,250 J	2,030	5,220 J	2,900 J	-	1,240 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	2	2 U	0.96 ∪	0.21 J	3.9 ∪	8.5 ∪	26 U	1.3 ∪	-	3 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	290 U	-	-	-	730 U	1920	4,900 U	240 U	-	798 J
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.78 ∪	-	-	-	1.1	2.2	1.3 U	0.13 U	-	0.66
Vanadium	520	78/NS	NS	NS	78/NS	22.68	27.3	-	-	-	401	1190	84.5 J	33.3	-	175
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	766	134	177	425	3,520	4,290	4,350	6,420	2,620	7,400
Fluoride	4,100	NS	7.4	GWP	3.95	7	3	-	-	-	15.4	16.1	5.4	1.4	-	6.7
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	14.7	-	_	_	228	195	23.6	62.6	7.8	787
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,680	-	-	-	5,070	6,700	70,000	9,020	-	6,450
Total Phosphorus	NS	NS	NS	NS	NS	NS	65	-	_	_	4,590	1,620	1,460	5,810	-	5,920
рН	NS	NS	NS	NS	NS	NS	7.64	-	-	-	5.6	5.43	6.18	6.57	-	5.49
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	_	_	_	15.8	_	_	_
Gross alpha Gross beta	NS NS	NS NS	NS	NS NS	NS	NS	-	-	_	-	-	-	14.5	-	-	-
Radium 226	5 ^(surface) , 15	140	INO	INO	INO	5 ^(surface) , 15	-	_	_	_	-	_	14.5	-	_	_
Nadium ZZU	5 (subsurface)	NO	NO	NO	NO	5 (subsurface)	-	_	_	-	-	-	4.6	_	_	-
D = 1' 200	,	NS	NS	NS	NS	,							4.6			
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	0.76 U	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revise EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated

Soil Sampling Analytical Results - 2010 Granulation Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-46	SB-46	SB-46	SB-46	SB-47	SB-47	SB-47	SB-47	SB-47	SB-48
Sample Type:		Protection of	of Environmen	ital Quality (b)	Background		DUP	INVE	INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	4-5	9-10	14-15	15-16	0-1	0-1	1-2	4-5	9-10	0-1

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluc

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentration
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the hurr
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimi

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for contraction to the EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to the publishe

WSP Environment & Energy
Page 4 of 6

Table D-7

Soil Sampling Analytical Results - 2010 Granulation Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Dookground		SB-48 INVE	SB-48 INVE	SB-48 INVE	SB-48 INVE	SB-49 INVE	SB-49 INVE	SB-49 INVE	SB-49 INVE	SB-49 INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Background Concentrations	Comparative	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base		Pathway	(UTL 95-95) ^(f)	Values (g)	1-2	4-5	9-10	14-15	0-1	1-2	4-5	9-10	11-12
Борин.	Son (mg/kg)	NISK-Dased/WCL-Dase	u ibits (ilig/kg)	Falliway	(OTE 33-33)	values									
Metals (mg/kg)															
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	3.2 ∪	0.98 ∪	-	-	1.4 U	2.1 U	1.2 U	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.2 ∪	4 J	3.2	3.3	4.6 J	2.6 J	4.2 J	2.4	3.6
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	96.1 J	128	-	-	102 J	137 J	146	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	1.6 U	0.8 J	-	-	1.5 J	1.1 U	0.72 J	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	29.6	0.49 U	-	-	17.2	16.9	0.59 U	-	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	115,000	3,510	-	-	43,500	52,700	4,630	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	234	25.6	-	-	125	118	24.9	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	25.2	11.3	-	-	25.4	11.4	10.7	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	4,680 J	4,700	-	-	7,280	5,700	4,720	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	36.7	17.6	-	-	38	31.7	16.7	-	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	3,500 J	2,610 J	-	-	1,900 J	3,550 J	2,560 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	6.4 U	2 U	-	-	2.9 ∪	4.2 U	2.4 U	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,270 J	384 J	-	-	1,520 J	790 U	440 U	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.35 J	0.47 J	-	-	0.43	0.29 J	0.29 J	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	277	30.9	-	-	185	176	32.2	-	-
Wet Chemistry (mg/kg)	NS	NS	4.4	Subsurface Soil	NS	NC	2 020	4 700	2 270	2.550	2 220	4 200	2.050	24.6	200
Ammonia (as N)	4,100	NS NS	4.1 7.4	GWP	3.95	NS 7	3,920	1,780	2,370	2,550	3,330	1,200	2,950	24.6	398
Fluoride		NS NS				/ NC	6.4	1.1 J	- 0.64.11	- E4 0	9.5	14.9	1.1 J	464	- -
Nitrate as N	1,600,000		18.4	GWP	NS	NS NS	238	18.5	0.61 U		477	448	314	164	50.6
Total Kjeldahl Nitrogen	NS	NS	NS	NS NC	NS NC	NS NS	5,220	2,140	-	-	2,740	2,370	3,180	-	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	3,310	377	-	-	2,280	1,620	2,360	-	-
рН	NS	NS	NS	NS	NS	NS	6.98	7.82	-	-	5.94	5.86	7.46	-	-
Radiological (pCi/g)															
Gross alpha	NS	NS	NS	NS	NS	NS	57.5	_	_	_	_	14.8	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	34	_	_	_	_	16.3	_	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	-	-	-	_	-	-	-	-	-
-	(subsurface)	NS	NS	NS	NS	(subsurface)									
Radium 228	15	NS NS	NS NS	NS NS	NS NS	15	_	_	_	_	_	_	_	_	_
Naululli 220	10	INO	INO	INO	INO	10	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical path GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revise EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated

Soil Sampling Analytical Results - 2010 Granulation Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Depa	artment			SB-48	SB-48	SB-48	SB-48	SB-49	SB-49	SB-49	SB-49	SB-49
Sample Type:		Protection of	of Environment	tal Quality ^(b)	Background		INVE								
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	1-2	4-5	9-10	14-15	0-1	1-2	4-5	9-10	11-12

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluc

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentration
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the hum
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimi

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for contraction to the EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to account for contraction to the published screening levels to the publishe

WSP Environment & Energy
Page 6 of 6

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment			SB-50	SB-50	SB-50	SB-50	SB-50	SB-51	SB-51	SB-51	SB-52	SB-52 INVE
Sample Type:		Protection of		ental Quality (b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative		08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/17/10	08/17/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	13-14	0-1	1-2	2-3	0-1	1-2
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	4 U	1.9 ∪	0.45 U	_	_	0.86 ∪	2.1 J	0.86 ∪	0.76 ∪	2.3 ∪
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	12.3 J	7.7 J	3.8	0.62 J	2	16	10.9	6	17.9	21.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	120 J	141 J	130	-	-	76.6 J	105 J	135 J	104	185 J
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	2 U	2.4 U	0.68 J	-	-	0.86 U	2 U	0.86 U	0.38 U	1.2 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	109	36.2	0.52 J	-	-	27.5	50.7	13.6	9.6	5.3
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	199,000	134,000	5,180	-	-	110,000	155,000	86,700	61,800	88,300
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	590	220	18.9	-	-	154	303	125	52.4	74.8
Lead	80	NS/14	50	GWP	NS/14	13.59	18.5	11	10.2	-	-	8.2	11.2 J	9.2	10.3	61
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	2,380 J	4,590 J	3,740	-	-	4,890	6,510 J	4,520	11,700	6,720
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	114	51.2	14.7	-	-	43.9	97	33.1	18.1	19.8
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	4,280 J	4,280 J	2,510	-	-	2,510 J	3,250 J	3,250 J	2,170 J	3,910 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	11.1 J	6.5 J	0.91 ∪	-	-	5 J	7.3 J	2.8 J	1.5 ∪	4.6 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,500 U	1,940 J	170 U	-	-	640 U	1,500 U	650 U	1,370 J	860 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	1.6 U	0.58	0.15 J	0.52 ∪	0.14 ∪	0.23 J	1.6 U	0.34 J	0.11 J	0.12 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	636	307	21.7	-	-	182	346	223	90.9	90.4
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	67.3	16.8	39.8	17.5	8.3	17.3	12.1	21.9	51.5	9.6
Fluoride	4,100	NS	7.4	GWP	3.95	7	43.5	24.7	1.1 J	-	-	21.2	28.4	29.9	52.9	11.4
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	12.1	6.1	2.4	_	_	12.3	19.8	5	0.54 U	24.2
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	974	684	1,240	_	_	496	522	921	342	578
Total Phosphorus	NS	NS	NS	NS	NS	NS	6,480	3,790	2,020	_	_	1,170	6,020	2,800	2,220	6,500
рН	NS	NS	NS	NS	NS	NS	6.58	8.09	7.59	-	-	6.86	7.45	7.65	6.98	8.05
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	41.6					56.2			41.9
Gross beta	NS NS	NS NS	NS NS	NS NS	NS	NS NS	-	37.2	-	-	-	-	37.1	-	-	33.7
Radium 226		INO	INO	INO	INO	5 ^(surface) , 15	-	31.2	-	-	-	-	31.1	-	-	JJ.1
Nadium 220	5 ^(surface) , 15 (subsurface)				NO	5 (subsurface)	-	-	-	-	-	-	-	-	-	-
D !' 000	,	NS	NS	NS	NS	, ,										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve	Idaho Dep	partment			SB-50	SB-50	SB-50	SB-50	SB-50	SB-51	SB-51	SB-51	SB-52	SB-52
Sample Type:		Protection of	of Environmen	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/17/10	08/17/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	13-14	0-1	1-2	2-3	0-1	1-2

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects. The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Table D-8

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment			SB-52	SB-52	SB-52	SB-52	SB-52	SB-53	SB-53	SB-53	SB-53	SB-53
Sample Type:		Protection of		ental Quality (b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative		08/17/10	08/17/10	08/17/10	08/17/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	4-5	9-10	14-15	19-20	23-24	0-1	1-2	4-5	9-10	14-15
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.52 ∪	-	-	-	_	2.2 J	0.93 ∪	3.2 U	0.34 J	0.53 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	4.6	0.45 J	4.5	4.6	2.8	7.2	4.3 J	7.3 J	2	3.3
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	151	-	-	-	-	272	130	214 J	107	110
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.63 J	-	-	-	-	1 U	0.75 J	1.6 ∪	0.32	0.79 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.31 J	0.43	0.94	0.81 J	0.81	34.5	8.5	29	3	3.3
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	29,000	-	-	-	-	81,900	101,000	170,000	248,000	14,600
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	17.5	-	-	-	-	160	48.3	167	7.9 J	31.8
Lead	80	NS/14	50	GWP	NS/14	13.59	10.1	0.47 J	9.1	14.8	8.7	125	8.7	27.9	4.7 J	10
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	6,950	-	-	-	-	5,720	6,550	5,160 J	3,440	7,460
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	16.2	-	-	-	-	44.2	20.4	31.8	88.1	24
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,980	-	-	-	-	3,810 J	2,720 J	5,690 J	1,380	2,940 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1 U	-	-	-	-	2.1 U	1.9 ∪	6.3 ∪	3.4 ∪	1.1 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	352 J	-	-	-	-	1,100 J	525 J	1730 J	148 J	400 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.18 J	0.63 ∪	0.41 J	0.11 U	0.62	0.44 J	0.093 U	0.79 ∪	0.85 ∪	0.15 J
Vanadium	520	78/NS	NS	NS	78/NS	22.68	27.1	-	-	-	-	226	114	206	8.5	40.4
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	44.2	46	7.9	14.3	165	245	31.2	28.6	130	33.7
Fluoride	4,100	NS	7.4	GWP	3.95	7	0.64 J	-	-	-	-	173	13.7	122	2.5	313
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	17.5	-	-	-	-	43.8	29.5	22.4	140	20
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	304	-	-	-	-	783	462	575	1,440	343
Total Phosphorus	NS	NS	NS	NS	NS	NS	1,410	-	-	-	-	4,090	2,390	7,090	2,620	1,060
pH .	NS	NS	NS	NS	NS	NS	7.7	-	-	-	-	6.08	7.38	6.75	6.94	5.66
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	_	_	_	19.5	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	-	-	_	-	_	-	12.1	-	_	-
Radium 226	5 ^(surface) , 15	-	-	-	-	5 ^(surface) , 15	-	_	-	-	_	-	_	-	_	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathwa GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-52	SB-52	SB-52	SB-52	SB-52	SB-53	SB-53	SB-53	SB-53	SB-53
Sample Type:		Protection of	of Environmen	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	4-5	9-10	14-15	19-20	23-24	0-1	1-2	4-5	9-10	14-15

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorion

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-8

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve		epartment			SB-53	SB-53	SB-54	SB-54	SB-54	SB-54	SB-54	SB-54	SB-54	SB-55
Sample Type:		Protection of		ental Quality (b)	Background		DUP	INVE	INVE	INVE	INVE	INVE	DUP	INVE	INVE	INVE
Sampling Date: Depth:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations (UTL 95-95) ^(f)	Comparative Values (g)	08/18/10 14-15	08/18/10 16-17	08/18/10 0-1	08/18/10 1-2	08/18/10 4-5	08/18/10 9-10	08/18/10 9-10	08/18/10 14-15	08/18/10 17-18	08/09/10 0-1
рерии.	Soil (mg/kg)	Risk-Based/MCL-Based	וטונג (mg/kg)	Pathway	(UIL 95-95)``	<u>values ""</u>	14-15	10-17		1-2	4-5	9-10	9-10	14-15	17-10	
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1 U	-	0.77 ∪	1.7 ∪	0.51 U	0.46 U	0.47 U	0.42 U	-	0.93 ∪
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.6	2.8	3.4 J	3.5 J	5.2	5.4	5.4	4.2	5.5	10
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	95.7	-	185	93.4 J	150	166	132	92.3	-	64.2 J
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.68	-	0.79 J	1.9 J	0.78 J	0.69 J	0.57	0.55 J	-	0.51 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	4.2	2.7	14.4	60.8	1.6	0.49 J	0.35	1.4	6.9	8.9
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	17,200	-	60,500	180,000	14,000	50,200	35,500	38,500	-	118,000
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	32.7	-	72.7	245	20.9	18.2	18.2	16.8	-	56
Lead	80	NS/14	50	GWP	NS/14	13.59	9.9	-	10.2	5 J	9.8	9	9.2	8.6	-	8
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	8,380	-	6,310	1,640 J	2,350	6,170	5,190	9,760	-	19,900
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	24.6	-	19.9	35	51.8	15.6	16.4	15.6	35.6	19
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,340	-	7,350 J	4,290 J	3,760	2,140 J	2,300	1,780 J	-	2,100 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	2 U	-	1.5 ∪	3.4 ∪	1.4 J	0.93 ∪	0.94 ∪	0.84 ∪	1 U	1.9 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	380 U	-	2,220 J	1,300 U	414 J	170 U	180 U	320 U	-	350 U
Thallium	í	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.1 U	_	0.19 J	1.7 U	0.51	0.38 J	0.19	0.13 J	0.13 U	0.093 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	42.3	34.4	123	405	25.8	23.8	24.5	22.2	-	73.5
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	44	59.5	50.5	1,260	160	1,140	1,610	1,020	1,230	288
Fluoride	4,100	NS	7.4	GWP	3.95	7	193	48.1	700	37.7	0.6 U	0.98 J	0.71	4.4	-,	34.4
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	26.3	15.4	11.2	71.1	165	151	153	41.1	30.8	116
Total Kieldahl Nitrogen	NS	NS	NS	NS	NS	NS	327	-	376	1,030	521	1,540	2,610	1,730	-	556
Total Phosphorus	NS	NS	NS	NS	NS	NS	707	-	13,900	14,900	6,700	2,490	6,960	8,610	-	5,730
рН	NS	NS	NS	NS	NS	NS	5.65	-	5.26	6.45	4.6	7.23	6.98	7.65	-	6.91
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	156	_	_	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	_	_	89	_	_	_	_	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	_	_	_	32.5	_	_	_	_	_	_
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS NS	NS NS	NS NS	15	-	-	-	0.54 U	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-53	SB-53	SB-54	SB-55						
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		DUP	INVE	INVE	INVE	INVE	INVE	DUP	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/18/10	08/09/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values ^(g)	14-15	16-17	0-1	1-2	4-5	9-10	9-10	14-15	17-18	0-1

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorion

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-8

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment			SB-55 INVE	SB-55 INVE	SB-55	SB-55	SB-55 INVE	SB-55	SB-56 INVE	SB-56 INVE	SB-56 INVE	SB-56 INVE
Sample Type: Sampling Date:	la decatal al	Protection of Groundwater ^(e)		ental Quality (b)	Background	0	08/09/10	08/09/10	INVE 08/09/10	INVE 08/09/10	1NVE 08/09/10	INVE 08/09/10	1NVE 08/10/10	08/10/10	08/10/10	08/10/10
Depth:	Industrial Soil (mg/kg)	Risk-Based/MCL-Base	Risk Based	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative Values (g)	1-2	4-5	9-10	14-15	19-20	24.5-25.5	0-1	1-2	4-5	9-10
'	(g/g/				101-00-00/											
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1.7 U	0.82 ∪	0.22 J	0.6 U	-	-	0.099 U	0.42 U	0.52 U	0.54 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	7.6 J	4.8	0.88	3	4.3	4	2.4	3.9	5.1	3
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	181	159	74.1	69.6	-	-	122	126	144	107
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	1 J	0.77 J	0.2	0.63 J	-	-	0.4	0.57 J	1.5	0.52 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	45.7	0.41 U	0.65	2.4	-	-	4.9	10.9	37.8	2.9
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	128,000	10,600	168,000	32,500	-	-	150,000	181,000	30,300	86,300
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	223	22	4.5	20.2	-	-	46.6	66.7	28.1	18
Lead	80	NS/14	50	GWP	NS/14	13.59	43.9	11.1	2.8 J	9.9	-	-	14.6	21.2	9.3	7.4
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	6,570	4,810	2,000	7,740	-	-	2,450	3,920	4,030	5,710
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	55.4	15.9	4 J	19.2	-	-	37.2	27.7	18.5	51
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	4,580 J	2,860 J	857	1,610 J	-	-	1,340	1,970 J	3,550	1,510 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	3.4 J	1.6 ∪	1.6 ∪	1.2 ∪	-	-	2 U	2.1 U	1 U	1.1 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,140 J	310 U	86 J	230 U	-	-	137 J	160 U	190 U	200 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	1.7 U	0.082 U	0.32 ∪	0.12 U	-	-	0.39 ∪	0.42 ∪	0.1 U	0.11 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	323	29.1	4.8	25.2	-	-	54.5	70.3	73.7	19.2
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	1,920	34.6	74	138	368	329	60.9	49.9	36.3	179
Fluoride	4,100	NS	7.4	GWP	3.95	7	111	4.3	1.5	0.8 J	-	-	5.1	14.1	29.5	3.1 U
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	75.8	53.2	25.7	24.6	33.2	36.4	120	16.6	35.9	224
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,080	492	1,810	298	-	-	893	1,060	500	1,720
Total Phosphorus	NS	NS	NS	NS	NS	NS	6,670	674	1,050	6,350	_	-	17,200	4,910	6,600	12,000
рН	NS	NS	NS	NS	NS	NS	6.01	7.16	7.28	7.26	-	-	7.03	7.1	6.04	6.15
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	57.7	_	_	_	_	_	_	14	_	_
Gross beta	NS	NS	NS	NS	NS	NS	41.9	_	_	_	_	_	_	8.9	_	_
Radium 226	5 ^(surface) , 15	1.0				5 ^(surface) , 15	-	_	_	_	_	_	_	-	_	_
	(subsurface)	NS	NS	NC	NS	(subsurface)										
Radium 228	15	NS NS	NS NS	NS NS	NS NS	15	_	_	_	_	_	_	_	_	_	_
radium ZZO	10	140	140	140	140	10		· ·	=	=	_	=	_	-	-	

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathwa GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-55	SB-55	SB-55	SB-55	SB-55	SB-55	SB-56	SB-56	SB-56	SB-56
Sample Type:		Protection of	of Environmen	ntal Quality ^(b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/09/10	08/10/10	08/10/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	1-2	4-5	9-10	14-15	19-20	24.5-25.5	0-1	1-2	4-5	9-10

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorio

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Table D-8

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment	Daalamaaaad		SB-56 DUP	SB-56 INVE	SB-56 INVE	SB-57 INVE	SB-57 INVE	SB-57 DUP	SB-57 INVE	SB-57 INVE	SB-57 INVE	SB-57 INVE
Sample Type: Sampling Date:	Industrial	Protection of Groundwater ^(e)	of Environme	ental Quality ^(b) Critical	Background Concentrations	Commonative	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
Depth:					(UTL 95-95) ^(f)	Comparative Values (g)	9-10	14-15	18.5-19.5	0-1	1-2	1-2	4-5	9-10	14-15	19-20
Берин.	Soil (mg/kg)	Risk-Based/MCL-Based	i iDTES (mg/kg)	Pathway	(UIL 95-95)**	values **	9-10	14-13	10.5-19.5	<u> </u>	1-2			9-10	14-13	13-20
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.47 U	0.4 U	-	2.1 U	1.7 U	-	2 U	0.32 J	0.59 U	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.7	4.9	4.9	8.1 J	8.7	-	3.7 J	2.8	4.8	6.2
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	110	132	-	156 J	101 J	-	127	116	152	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.58	0.71 J	-	1.5 J	0.85 U	-	1.1	0.44 J	0.81 J	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	5.7	0.88	1.9	54.7	17.5	-	37.8	0.74	1.1 J	0.49 J
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	89,500	31,000	-	74,800	46,800	-	74,800	159,000	34,600	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	14.9	14.9	-	297	160	-	177	14.7	19.3	-
Lead	80	NS/14	50	GWP	NS/14	13.59	7.3	9.9	-	35.5	9.3	-	17.3	6	10.5	11.6
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	3,900	6,770	-	7,460	2,180 J	-	5,410	3,890	5,960	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	77.8	19.7	-	81.6	58.2	-	67.5	12.5	19.1	19.5
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,780	1,640 J	-	3,840 J	2,380 J	-	3,630 J	1,370	2,130 J	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	0.93 ∪	0.8 ∪	-	5.6 J	6.4 J	-	4.1 U	2.1 ∪	1.2 U	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	170 U	150 U	-	800 U	640 U	-	760 U	79 U	220 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.093 U	0.08 U	-	1.4	0.51	-	0.34 J	1.1 U	0.12 U	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	15.8	22	-	456	123	-	265	10.9	22.6	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	190	827	803	2,200	1,270	-	5,020	3,560	4,280	3,730
Fluoride	4,100	NS	7.4	GWP	3.95	7	1.4	2.4	5.2	135	88.1	-	239	2	2.1	1.5
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	52.9	91	34.5	1,680	813	-	346	282	141	29.7
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,930	51.7	-	3,900	1,730	-	8,660	5,700	3,320	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	8,900	2,470	-	31,100	3,730	-	6,460	2,550	5,380	-
рН	NS	NS	NS	NS	NS	NS	6	6.84	-	5.55	5.36	-	6.87	8.01	8.15	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	_	16.3	17.2	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	_	_	_	10.8	13.5	_	_	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) . 15	_	-	-	_	-	-	-	_	_	-
<u> </u>	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS NS	NS NS	NS NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-56	SB-56	SB-56	SB-57						
Sample Type:		Protection of	of Environmen	ital Quality ^(b)	Background		DUP	INVE	INVE	INVE	INVE	DUP	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	9-10	14-15	18.5-19.5	0-1	1-2	1-2	4-5	9-10	14-15	19-20

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluorio

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

WSP Environment & Energy
Page 10 of 12

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		partment ntal Quality ^(b)	Background		SB-57 INVE	SB-58 INVE	SB-58 INVE	SB-58 INVE	SB-58 INVE	SB-58 INVE	SB-58 DUP
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	24-25	0-1	1-2	4-5	9-10	14-15	14-15
•		= = = = = = = = = = = = = = = = =			(
Metals (mg/kg)													
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	1.6 ∪	2 U	0.37 U	0.29 J	0.41 U	0.48 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	8.7	4.6 J	7.1 J	4.4	3.6	2.7	3.2
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	48.3	141	128	124	93.9	94.3
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	0.81 U	1.1	0.87 J	0.56	0.63 J	0.6
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.42 U	10.9	36	38.7	1.6	0.85	0.85
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	120,000	58,500	24,100	212,000	24,400	26,500
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	88.4	253	19.5	12.7	17.9	18
Lead	80	NS/14	50	GWP	NS/14	13.59	11.2	2.7 J	13.8	9.2	5.5	9.2	9
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	62,000	16,100	2,610	5,030	9,850	9,430
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	20	28.3 J	62.3	57.2	10.6	16.2	16.2
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	1,240 J	3,690 J	3,450	1,940	1,970 J	1,880
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	3.2 ∪	3.9 ∪	0.74 ∪	2.3 J	0.83 ∪	0.97 ∪
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	610 U	801 J	564 J	204	159 J	222
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	1.6 ∪	0.71	0.092 U	0.11 U	0.1 U	0.097 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	97.2	275	27.2	15.5	23.2	28.4
Wet Chemistry (mg/kg)													
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	4,240	24.4	1,740	2,060	4,770	2,630	2,790
Fluoride	4,100	NS	7.4	GWP	3.95	7	0.6 U	32.7	50.3	1.5	2.5	1.4	1.3
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	43	182	511	348	141	166	186
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	372	1,470	4,200	9,380	3,730	3,710
Total Phosphorus	NS	NS	NS	NS	NS	NS	-	1,460	5,450	8,880	11,800	2,170	1,230
рН	NS	NS	NS	NS	NS	NS	-	7.31	5.67	5.27	7.78	7.64	7.63
Radiological (pCi/g)													
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	15.6	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	_	15.8	_	_	_	_
Radium 226	5 ^(surface) , 15	110	110	110	110	5 ^(surface) , 15	_	_	-	_	_	_	_
	(subsurface)	NS	NS	NS	NS	(subsurface)							
Dodium 220		NS NS	NS NS	NS NS	NS NS								
Radium 228	15	ONI	IND	CNI	INO	15	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathwa GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Process Sewer Lagoon Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-57	SB-58	SB-58	SB-58	SB-58	SB-58	SB-58
Sample Type:		Protection of	of Environmen	tal Quality (b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	DUP
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	24-25	0-1	1-2	4-5	9-10	14-15	14-15

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluories

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment	Deed was and		SB-59	SB-59 INVE	SB-59	SB-59 INVE	SB-60 INVE	SB-60 INVE	SB-60 INVE	SB-60 INVE	SB-61	SB-61
Sample Type:	to to atilat	Protection of		ental Quality (b)	Background	0	INVE		DUP 08/17/10					1NVE 08/17/10	INVE 08/17/10	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10		08/17/10	08/17/10	08/17/10	08/17/10			08/17/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	ed IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	1-2	4-5	0-1	1-2	4-5	6-7	0-1	1-2
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	2 U	0.75 ∪	-	8.9 ∪	1.7 ∪	0.8 ∪	1 U	-	1.8 ∪	
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	8.6 J	3.1 J	-	10.8 J	9.2	3.5 J	6	3.9	5.4 J	5.7 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	189	114	-	318 J	173	112	114	-	146	209
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	2 U	0.37 U	-	4.5 U	0.86 U	0.43 J	0.52 U	-	0.9 U	0.87 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	20.9	13.3	-	11.3 J	25.5	8.3	23.4	1.1	15.1	5.6
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	122,000	110,000	-	49,200	89,600	88,000	18,700	-	62,200	41,300
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	124	74.3	-	228	143	56.5	113	-	89.4	42.8
Lead	80	NS/14	50	GWP	NS/14	13.59	17.3	10.4	-	21.4	40.3	6.7	56.8	10.8	12.5	10.6
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	15,100	13,000	-	3,900 J	11,800	11,500	2,340 J	-	9,030	4,330
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	30	23.3	-	10.2	37.7	18.5	36.9	-	26.6	18.8
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	5,330 J	3,890	-	6,740 J	4,890 J	2,930 J	1,910 J	-	3,940 J	1,850 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	4 U	1.5 ∪	-	18 U	3.5 ∪	1.6 ∪	5.2 J	0.91 ∪	3.6 ∪	3.5 ∪
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,500 J	2,180 J	-	3,300 U	2,380 J	708 J	390 U	-	1,050 J	650 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.2 J	0.75 ∪	-	0.37 J	0.099 J	0.12	0.42 J	0.12 U	0.22	0.087 J
Vanadium	520	78/NS	NS	NS	78/NS	22.68	173	107	-	240	236	81	136	-	164	63.3
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	148	83.2	-	367	68.3	9.9	451	772	168	26
Fluoride	4,100	NS	7.4	GWP	3.95	7	248	266	-	621	92.2	183	16.8	1.1 J	53.7	338
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	27.2	18.9	-	17.9	18.1	11.6	22	0.67 U	75.6	30.4
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	434	196	-	1,010	588	3,290	1,400	-	365	321
Total Phosphorus	NS	NS	NS	NS	NS	NS	11,500	3,150	-	15,200	3,140	4,260	7,710	-	5,300	5,590
рН	NS	NS	NS	NS	NS	NS	6.16	6.18	-	4.01	5.59	6.34	5.85	-	5.22	6.61
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	13.6	14.4	_	_	15.1	_	_	_	54
Gross beta	NS	NS	NS	NS	NS	NS	-	10.9	16.2	-	-	11.5	-	-	_	37.7
Radium 226	5 ^(surface) . 15	-	-	-	-	5 ^(surface) , 15	-	-	-	-	-	-	_	-	-	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS NS	NS NS	NS NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve	Idaho Dep	partment			SB-59	SB-59	SB-59	SB-59	SB-60	SB-60	SB-60	SB-60	SB-61	SB-61
Sample Type:		Protection of	of Environmen	ntal Quality ^(b)	Background		INVE	INVE	DUP	INVE						
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	0-1	1-2	1-2	4-5	0-1	1-2	4-5	6-7	0-1	1-2

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects. The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment			SB-61	SB-61	SB-62	SB-62	SB-62	SB-62	SB-62	SB-62	SB-63	SB-63
Sample Type:		Protection of		ental Quality (b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	4-5	5.5-6.5	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	10 U	-	1.4 ∪	3.7 U	3.1 U	-	-	-	0.41 U	2.3 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	24 J	5.9	4.4 J	4.1 J	6.5 J	3	7.3	5.3	1.8 J	3.2 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	402 J	521	88 J	158	81.4	-	-	-	85.9	150
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	10 U	0.65 J	0.69 U	1.8 U	1.6 ∪	-	-	-	0.21 J	1.1 U
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	5.2 ∪	-	15.2	26.1	17.1	0.71	7.4	5.7	3.6	39.9
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	73,500	-	82,900	117,000	217,000	-	-	-	161,000	149,000
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	100	-	92.4	120	307	-	-	-	25.2	179
Lead	80	NS/14	50	GWP	NS/14	13.59	24.7	23	9.7	3.3	11.8	-	-	-	10.3	19.4
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	1,190 J	-	9,290	7,260 J	6,640 J	-	-	-	7,370	7,090
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	4.1	-	28.7	21	25	-	-	-	18.7	46.6
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	5,200 U	-	1,770 J	3,350 J	5,610 J	-	-	-	937 J	2,360 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	21 U	-	4.1	7.4 U	8.5	0.84 ∪	0.66 ∪	0.9 ∪	1.6 ∪	4.5 ∪
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	7,800 U	-	1,950	3,040 J	2,690	-	-	-	803 J	840 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	1.1	3.3	0.69 ∪	1.8 U	1.6	0.14 ∪	0.11 U	0.26 J	0.082 U	0.47 J
Vanadium	520	78/NS	NS	NS	78/NS	22.68	140 J	-	143	217	322	15	54.8	65.8	49.4	243
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	673	872	48.7	106	16.7	655	692	194	30.4	2,070
Fluoride	4,100	NS	7.4	GWP	3.95	7	579	74.5	33.5	33	20.2	1.7	642	838	85.9	188
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	6.5	-	38	49.2	54.1	0.6 U	0.61 U	0.61 U	106	103
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,690	-	611	307	373	-	_	-	279	1,930
Total Phosphorus	NS	NS	NS	NS	NS	NS	6,200	-	4,690	5,740	6,040	-	_	-	1,060	4,520
рН	NS	NS	NS	NS	NS	NS	5.17	-	6.85	7.03	8.78	-	-	-	6.65	6.4
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	47.8	_	_	_	_	_	29.1
Gross beta	NS	NS	NS	NS	NS	NS	_	-	-	31.4	-	_	_	-	-	17.8
Radium 226	5 ^(surface) . 15					5 ^(surface) . 15	_	-	_	-	-	-	_	_	_	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS NS	NS NS	NS	15	-	-	-	_	-	-	_	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-61	SB-61	SB-62	SB-62	SB-62	SB-62	SB-62	SB-62	SB-63	SB-63
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/17/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	4-5	5.5-6.5	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type: Sampling Date:	Region 9 Regi	ional Screening Leve Protection of Groundwater ^(e)	of Environme	epartment ental Quality ^(b) Critical	Background Concentrations	Commonative	SB-63 INVE 08/10/10	SB-63 INVE 08/10/10	SB-63 INVE 08/10/10	SB-63 INVE 08/10/10	SB-64 INVE 08/10/10	SB-64 INVE 08/10/10	SB-64 INVE 08/10/10	SB-64 INVE 08/10/10	SB-64 INVE 08/10/10	SB-64 INVE 08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	Risk Based IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Comparative <u>Values ^(g)</u>	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10	14-15	16-17
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.81 ∪	-	-	-	2.4	2.4 U	0.44 J	-	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3 J	2.9	6.1	5.4	6.4 J	4.9 J	2.6	3.1	3.9	4.5
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	159	-	-	-	144 J	130 J	153	-	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.78 J	-	-	-	1.2 J	1.5	0.81 J	-	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	22.4	0.4 J	0.23 U	0.27 U	57	143	0.51	-	-	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	26,900	-	-	-	118,000	83,700	25,600	-	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	21.1	-	-	-	242	471	22.4	-	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	10.8	-	-	-	12.4	18.5	9.6	-	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	4,610	-	-	-	7,870	5,680 J	6,400	-	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	68.1	14.9	20.9	22.8	68.2	63.7	57.1	13.5	16.3	25.4
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	3,410 J	-	-	-	4,270 J	3,840 J	3,000 J	-	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.6 ∪	-	-	-	4.7 ∪	4.8 U	0.69 J	-	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	300 U	-	-	-	2,140 J	1,480 J	230 U	-	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.27 J	-	-	-	0.98	1.5 J	0.14 J	-	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	49.9	-	-	-	374	961	81.5	-	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	3,390	2,340	4,280	4,190	8.9	6	5.8	5,990	3,290	2,340
Fluoride	4,100	NS	7.4	GWP	3.95	7	17.7	1.7	1.8	1.4	197	47.3	0.63 U	-	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	35.8	13.8	2.7	7	378	61.8	48	12.7	20.2	5.1
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	5,410	-	-	-	5,870	7,880	6,140	-	-	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,980	-	-	-	32,100	12,000	2,480	-	-	-
рН	NS	NS	NS	NS	NS	NS	6.79	-	-	-	5.75	5.43	6.87	-	-	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	-	-	-	-	86	-	-	-	_
Gross beta	NS	NS	NS	NS	NS	NS	-	-	-	-	-	62	-	-	-	-
Radium 226	5 (surface), 15					5 (surface), 15	-	-	-	-	-	12.7	-	-	-	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	0.8 U	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-63	SB-63	SB-63	SB-63	SB-64	SB-64	SB-64	SB-64	SB-64	SB-64
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10	08/10/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10	14-15	16-17

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Pookaround		SB-65 INVE	SB-65 INVE	SB-65 INVE	SB-65 INVE	SB-65 INVE	SB-65 INVE	SB-66 INVE	SB-66 INVE	SB-66 INVE	SB-66 INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Background Concentrations	Comparative		08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	3.5 ∪	4.1 U	0.8 U	_	_	_	3.3 ∪	3.4 U	0.42 U	_
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	10.6 J	4.1 U	3.9 J	6	3.2	13.5	6.7 J	4.9 J	3.9	1.6
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	49.9 J	128 J	154	-	-	-	173 J	145	155	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	1.8 ∪	2 U	0.77 J	_	-	_	1.6 ∪	1.7 U	0.68 J	_
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	3.9 J	6.3 J	0.4 U	-	-	-	29.7	21.6	0.35 J	_
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	87,700	169,000	3,290	-	-	-	144,000	207,000	32,800	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	98.2	170	24.5	-	-	-	652	241	17.6	-
Lead	80	NS/14	50	GWP	NS/14	13.59	1.7	3.9	10.8	-	-	-	19.2	15.3 J	8.8	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	1,200 J	3,920 J	4,030	-	-	_	7,870 J	2,680 J	6,570	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	37.8	26.7	16.3	-	-	_	51.7	53.4 J	15.1	_
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,290 J	4,790 J	3,080 J	-	-	-	2,350 J	5,130 J	2,920	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	7.1 U	8.1 U	1.6 ∪	-	-	-	6.6 ∪	6.7 ∪	0.84 ∪	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,300 U	1,500 U	300 U	-	-	-	2,310 J	2,640 J	160 U	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.088 U	0.1 U	0.21 J	-	-	-	0.72	3.4 U	0.084 U	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	133	194	28.2	-	-	-	276	302	25.2	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	2,570	1,300	1,280	7,630	4,720	183	807	791	1,240	5,030
Fluoride	4,100	NS	7.4	GWP	3.95	7	5.2	9.8	0.8 J	, -	, <u> </u>	-	2.5	10.8	2.3	· -
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	74.5	14.3	16.3	-	-	-	47.1	137	101	244
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,410	1,880	6,310	-	-	_	3,760	2,190	7,220	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,610	1,280	581	-	-	-	4,870	2,620	7,060	-
рН	NS	NS	NS	NS	NS	NS	7.52	8.45	6.61	-	-	-	7.69	8.25	6.46	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	89	_	_	_	_	_	78	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	63	_	_	_	_	_	49.8	_	_
Radium 226	5 ^(surface) . 15					5 ^(surface) , 15	-	-	-	_	-	-	_	-	-	_
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-65	SB-65	SB-65	SB-65	SB-65	SB-65	SB-66	SB-66	SB-66	SB-66
Sample Type:		Protection of	of Environmen	tal Quality (b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-66 INVE	SB-66 INVE	SB-66 INVE	SB-67 INVE	SB-67 INVE	SB-67 INVE	SB-67 INVE	SB-67 INVE	SB-67 INVE	SB-67 INVE
Sampling Date: Depth:	Industrial Soil (mg/kg)	Groundwater ^(e) Risk-Based/MCL-Based	Risk Based	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative <u>Values ^(g)</u>	08/13/10 14-15	08/13/10 19-20	08/13/10 20-21	08/13/10 0-1	08/13/10 1-2	08/13/10 4-5	08/13/10 9-10	08/13/10 14-15	08/13/10 19-20	08/13/10 22-23
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	_	_	_	1.9 ∪	0.089 U	0.8 U	_	_	_	_
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.6	5.5 J	5.9	6.4 J	0.54	3.3 J	7.2	7.9	5.6	6.6
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	-	-	113	12.8	166	-	-	_	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	-	-	0.96	0.17 J	0.74 J	-	-	_	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	-	-	-	31.1	1.1	0.48 J	-	-	-	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	-	-	87,900	128,000	14,400	-	-	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	-	-	244	145	20.4	-	-	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	-	-	-	19.6	2.1 J	10.9	-	-	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	-	-	3,900 J	227	4,160	-	-	-	_
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	-	-	-	57.10	44.4	14.9	-	-	-	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	-	-	1,980 J	517	2,660 J	-	-	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	-	-	3.7 ∪	1.8 ∪	1.6 U	-	-	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	-	-	700 U	237 J	300 U	-	-	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	-	-	0.12 J	1.8 ∪	0.13 J	-	-	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	-	-	247	36.2	25.9	-	-	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	2,540	5,020	2,990	535	307	1,510	5,390	4,060	3,700	2,980
Fluoride	4,100	NS	7.4	GWP	3.95	7	· -	-	-	8.6	4	2.7	· -	-	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	211	24.3	1.5	42.5	46.5	70.6	11.9	62.2	37.4	39.8
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	_	-	4,660	880	3,870	-	-	-	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	-	_	-	1,260	12,900	4,930	-	-	-	-
рН	NS	NS	NS	NS	NS	NS	-	-	-	7.94	8.73	7.36	-	-	-	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	-	-	-	95	-	-	-	-	-
Gross beta	NS	NS	NS	NS	NS	NS	-	-	-	-	63	-	-	-	-	-
Radium 226	5 (surface), 15					5 ^(surface) , 15	-	-	-	-	45.5	-	-	-	-	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	0.51 U	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-66	SB-66	SB-66	SB-67						
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10	08/13/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values ^(g)	14-15	19-20	20-21	0-1	1-2	4-5	9-10	14-15	19-20	22-23

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- i/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Dooleanound		SB-68 INVE	SB-69 INVE	SB-69 INVE	SB-69 INVE						
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Background Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Comparative <u>Values ^(g)</u>	0-1	1-2	4-5	9-10	14-15	19-20	21.5-22.5	0-1	1-2	4-5
Берин.	Soli (mg/kg)	RISK-Based/WCL-Based	IDILS (mg/kg)	Pathway	(UIL 95-95)**	<u>values</u>	<u> </u>	1-2	4-3	9-10	14-13	19-20	21.5-22.5		1-2	
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	2 U	0.81 ∪	0.87 ∪	-	-	-	-	0.83 ∪	1 U	0.5 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	5.1 J	6.3	5.1	2.1	5.2	5.1	4.9	5	4.5 J	4.4
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	128 J	181	153	-	-	-	-	92.7	124	154
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	1.4 J	0.9 J	0.68 J	-	-	-	-	1.1 J	0.81 J	0.74 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	62.9	6	0.46 J	-	-	-	-	37.9	18.6	0.47 J
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	83,300	14,800	44,000	-	-	-	-	50,400	31,400	28,200
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	338	24.1	18.8	-	-	-	-	182	57.1	16.4
Lead	80	NS/14	50	GWP	NS/14	13.59	28.3	11.1	9.4	-	-	-	-	16.9	8.8	9.4
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	5,860	4,110	6,610	-	-	-	-	4,510	3,250	6,450
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	57.6	30.4	15.4	-	-	-	-	53.2	54.6	15
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,740 J	2,790 J	1,960 J	-	-	-	-	1,880 J	2,270 J	2,440 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	3.9 ∪	2.2 J	1.7 U	-	-	-	-	2.7 J	3.5 J	1.4 J
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	1,200 J	300 U	330 U	-	-	-	-	780 J	421 J	190 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.098 U	0.081 U	0.87 ∪	-	-	-	-	0.083 U	0.1 U	1 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	434	66.2	27.3	-	-	-	-	297	145	36.8
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	2,170	941	402	423	289	1,780	183	5,010	2,350	3,180
Fluoride	4,100	NS	7.4	GWP	3.95	7	168	8.7	1.7	-	_	, <u>-</u>	-	18.8	14.4	0.63 U
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	238	36.6	59.2	7	5.5	25	19.4	185	37.9	23.9
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	2,010	2,210	962	-	_	_	-	10,900	1,150	6,130
Total Phosphorus	NS	NS	NS	NS	NS	NS	8,390	850 U	85	_	_	_	-	16,300	3,700	4,980
рН	NS	NS	NS	NS	NS	NS	5.88	5.86	7.79	-	-	-	-	5.88	5.92	7.02
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	12.8	_	_	_	_	_	_	16.8	_
Gross beta	NS	NS	NS	NS	NS	NS	_	8.7	_	_	_	_	_	_	14.1	_
Radium 226	5 ^(surface) . 15	110	110	110	110	5 ^(surface) . 15	_	-	_	_	_	_	_	_	-	_
radiani 220	O , IO (subsurface)	NS	NC	NC	NC	O , IO (subsurface)										
Radium 229	,	NS NS	NS NS	NS NS	NS NC	,										
Radium 228	15	ONI	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-68	SB-69	SB-69	SB-69						
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE	INVE	INVE	INVE						
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	0-1	1-2	4-5	9-10	14-15	19-20	21.5-22.5	0-1	1-2	4-5

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve		epartment	Daalamaa a		SB-69 INVE	SB-69 INVE	SB-69 INVE	SB-69 INVE	SB-70 INVE	SB-70 INVE	SB-70 INVE	SB-70 INVE	SB-71 INVE	SB-71 DUP
	In the state of	Protection of		ental Quality (b)	Background	0		08/16/10	08/16/10	08/16/10	08/13/10	08/13/10		08/13/10	08/16/10	08/16/10
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10						08/13/10			
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	9-10	14-15	19-20	20-21	0-1	1-2	4-5	6.5-7.5	0-1	0-1
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	-	-	-	-	1.8 U	1.6 ∪	1.8 ∪	-	7.7 J	10.8
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.8	8.9	4.2	4.1	5.2 J	4.7 J	6.1 J	7.3	7 J	10.8
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	-	-	-	-	159 J	168	210	-	192 J	289
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	-	-	-	-	1.1 J	0.95 J	0.96 J	-	1.3 J	1.9
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	-	-	-	-	26.8	3.1	1.1 J	-	51.1	115
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	-	-	-	-	50,600	24,600	4,450	-	57,500	76,100
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	-	-	-	-	148	40.3	24	-	286	443
Lead	80	NS/14	50	GWP	NS/14	13.59	-	-	-	-	14.2	10.3	12.7	-	52	104
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	-	-	-	-	5,480	4,130	4,360 J	-	5,430	7,300
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	-	-	-	-	34.7	27.2	17.8	-	104	130
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	-	-	-	-	2,980 J	2,860 J	3,800 J	-	4,320 J	3,750
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	-	-	-	-	3.6 ∪	3.1 U	3.5 ∪	-	4.7 J	5.3
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	-	-	-	-	874 J	590 U	660 U	-	740 U	1,660
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	-	-	-	-	0.33 J	0.079 U	0.088 U	-	0.4 J	0.27
Vanadium	520	78/NS	NS	NS	78/NS	22.68	-	-	-	-	253	88.5	45.4	-	338	616
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	3,640	1,300	2,610	1,680	6.9	2	0.71 U	-	4,360	4,700
Fluoride	4,100	NS	7.4	GWP	3.95	7	, <u>-</u>	, <u>-</u>	, -	, <u>-</u>	50.1	17.5	0.6 U	-	46.7	93.4
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	40.7	8	27	39	32.1	32.4	29.6	10.7	333	287
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	-	_	-	_	1,890	742	380	-	5,360	10,500
Total Phosphorus	NS	NS	NS	NS	NS	NS	-	_	-	_	3,960	8,130	9,800	-	7,240	4,050
рH	NS	NS	NS	NS	NS	NS	-	-	-	-	5.32	4.92	4.57	-	5.91	5.77
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	-	_	-	_	_	6.8	-	-	_	_
Gross beta	NS	NS	NS	NS	NS	NS	-	-	-	_	-	5.5	-	-	_	_
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	-	-	-	_	-	-	-	-	_	_
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	_	-	_	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-69	SB-69	SB-69	SB-69	SB-70	SB-70	SB-70	SB-70	SB-71	SB-71
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE	DUP								
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10	08/13/10	08/13/10	08/13/10	08/13/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values ^(g)	9-10	14-15	19-20	20-21	0-1	1-2	4-5	6.5-7.5	0-1	0-1

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve		epartment			SB-71	SB-71	SB-71	SB-71
Sample Type:		Protection of		ental Quality (b)	Background	•	INVE	DUP	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values (g)</u>	1-2	1-2	4-5	5-6
Motolo (ma/ka)										
Metals (mg/kg)	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	1 U	_	0.12 J	
Antimony Arsenic	99,000 41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.9 J	-	2.4	2.4
Barium	1.6	1,200/82	896	GWP	1,200/82	4.7 170.2	147	-	186	2.4
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.94 J	_	0.62 J	_
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.94 J 1.3 J	-	0.62 3	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	64,700	-	62,700	-
Chromium		NS/180,000	2,135	GWP	NS/180,000	36,270 (I) 18.61	49.6	-	15	-
Lead	(j) 80	NS/14	2,133 50	GWP	NS/14	13.59	7.8	-	8.2	-
		NS/NS	NS	NS	NS/NS	7,146	7.6 3,400	-	2,910	-
Magnesium Nickel	(j)	20/NS		GWP	20/NS	7,146 15.71	3,400 44.9	-	32.2	-
Potassium	2,000	NS/NS	59 NS	NS	NS/NS	3,393 (I)	44.9 4,690 J	-	2,800	-
	(j) 510	0.4/0.26	2.0	GWP	0.4/0.26	3,393 (I) 1.04	4,690 J 2.9 J	-	2,000 0.24 U	-
Selenium		0.4/0.26 NS/NS	NS NS	NS	0.4/0.26 NS/NS		2.9 J 923 J	-	0.24 U 304 J	-
Sodium Thallium	(j) 1	0.0011/0.14		GWP	0.0011/0.14	289 (I) NS	923 J 0.1 U			-
	520	78/NS	1.6 NS	NS	78/NS			-	0.12 U	-
Vanadium	520	78/NS	NS.	INS	78/NS	22.68	96.8	-	29.5	-
Wet Chemistry (mg/kg)										
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	2,540	_	1,180	1,200
Fluoride	4,100	NS	7.4	GWP	3.95	7	12.8	-	0.64 U	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	1,670	-	640	1,010
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	2,030	-	1,260	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	3,730	-	6,960	_
pH	NS	NS	NS	NS	NS	NS	6.25	-	4.77	-
·										
Radiological (pCi/g)										
Gross alpha	NS	NS	NS	NS	NS	NS	19.8	66	-	-
Gross beta	NS	NS	NS	NS	NS	NS	24.2	51.9	-	-
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	-	-	-	-
	(subsurface)	NS	NS	NS	NS	(subsurface)				
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised. EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated a

Soil Sampling Analytical Results - 2010 Dry Products Storage East Ditch Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-71	SB-71	SB-71	SB-71
Sample Type:		Protection of	of Environmen	tal Quality (b)	Background		INVE	DUP	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/16/10	08/16/10	08/16/10	08/16/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	1-2	1-2	4-5	5-6

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoric

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cun

Page 16 of 16

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve		epartment	David services 1		SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-72 INVE	SB-73 INVE
Sample Type:	la la dalatat	Protection of		ental Quality (b)	Background	0										08/03/10
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative Values (g)	08/03/10 0-1	08/03/10 1-2	08/03/10 4-5	08/03/10 9-10	08/03/10 14-15	08/03/10 19-20	08/03/10 24-25	08/03/10 29-30	08/03/10 31-32	08/03/10 0-1
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDILS (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>values (3)</u>	<u>U-1</u>	1-2	4-5	9-10	14-15	19-20	24-25	29-30	31-32	<u>U-1</u>
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.53 U	0.45 U	0.42 U	-	-	-	-	-	-	0.49 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.8	4.0	4.1	4.9	4.2	3.5	5.8	4.3	8.4	4.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	126	132	132	-	-	-	-	-	-	133
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.71 J	0.70 J	0.76 J	-	-	-	-	-	-	0.74 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.99 J	0.34 J	0.28 J	-	-	-	-	-	-	0.42 J
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	7,540	6,600	4,990	-	-	-	-	-	-	5,090
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	23	19.2	21.5	-	-	-	-	-	-	19.8
Lead	80	NS/14	50	GWP	NS/14	13.59	10.8	10.8	10.9	-	-	-	-	-	-	11.5
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	3,980	3,740	4,140	-	-	-	-	-	-	3,230
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	17.8	16.6	18.5	-	-	-	-	-	-	15.3
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,540 J	2,290	1,960 J	-	-	-	-	-	-	3,030
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.1 U	0.9 ∪	1.2 J	-	-	-	-	-	-	0.99 ∪
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	715 J	484 J	160 U	-	-	-	-	-	-	190 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.11 U	0.09 U	0.083 U	-	-	-	-	-	-	0.99 ∪
Vanadium	520	78/NS	NS	NS	78/NS	22.68	30.2	24	26.4	-	-	-	-	-	-	24
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	68.3	65.5	49.5	9.1	8	5.9	6.1	2.9	3.5	99.9
Fluoride	4,100	NS	7.4	GWP	3.95	7	35.4	52.1	19.4	3.3	1.9	1.3	1.8	1.7	1.8	27.6
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	1.5	0.95 J	0.98 J	_	-	_	-	-	_	2.3
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	782	917	494	-	-	-	-	-	-	1,390
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,590	2,070	1,360	-	-	-	-	-	-	1,700
pH .	NS	NS	NS	NS	NS	NS	8.54	7.72	6.58	-	-	-	-	-	-	6.57
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	4.8	_	_	_	_	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	_	4.1	_	_	_	_	_	_	_	_
Radium 226	5 ^(surface) . 15					5 ^(surface) , 15	_	-	_	_	_	_	_	_	_	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS NS	NS NS	NS NS	15	_	_	_	_	_	_	_	_	_	_
	. •					. •										

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	partment			SB-72	SB-73								
Sample Type:		Protection of	of Environmer	ntal Quality ^(b)	Background		INVE									
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values (g)</u>	0-1	1-2	4-5	9-10	14-15	19-20	24-25	29-30	31-32	0-1

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects. The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Dooleanound		SB-73 INVE	SB-73 INVE	SB-73 INVE	SB-73 INVE	SB-73 INVE	SB-73 INVE	SB-73 INVE	SB-73 INVE	SB-74 INVE	SB-74 INVE
Sampling Date:	ا ماد، مدینا	Groundwater ^(e)	Risk Based	Critical	Background Concentrations	Commonative	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Industrial				(UTL 95-95) ^(f)	Comparative Values ^(g)	1-2	4-5	9-10	14-15	19-20	24-25	29-30	31.5-32.5	0-1	1-2
Бериі.	Soil (mg/kg)	Risk-Based/MCL-Based	IDTES (mg/kg)	Pathway	(UIL 95-95)\^	<u>values (*)</u>		4-3	9-10	14-13	19-20	24-25	29-30	31.5-32.5		<u> </u>
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.40 U	0.43 U							0.5 U	0.51 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.8	4.8	3.9	4.4	4	5.6	5	3	4.2	3.8
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	137	146	-	-	-	-	-	-	83.7	63.5
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.75 J	0.86 J	-	-	-	-	-	-	0.52 J	0.47 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.38 J	0.22 J	-	-	-	-	-	-	5	4.4
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	4,700	4,450	-	-	-	-	-	-	49,800	52,500
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	19	23.4	-	-	-	-	-	-	36.6	39.7
Lead	80	NS/14	50	GWP	NS/14	13.59	11.4	11.4	-	-	-	-	-	-	6.4	6.1
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	3,230	4,650	-	-	-	-	-	-	12,000	13,000
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	15.2	19.7	-	-	-	-	-	-	28.8	23.4
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	3,180	1,790 J	-	-	-	-	-	-	1,470 J	1,450 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	0.81 ∪	0.87 ∪	-	-	-	-	-	-	2.2 J	2.2 J
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	150 U	160 U	-	-	-	-	-	-	400 J	190 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.81 ∪	0.087 U	-	-	-	-	-	-	0.099 U	0.1 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	22.5	30	-	-	-	-	-	-	50.9	48.2
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	99	24.6	6.9	10.6	2.3	6.3	3.2	1.6	13.9	129
Fluoride	4,100	NS	7.4	GWP	3.95	7	34.5	2	-	-	-	-	-	-	7.1	111
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	3.3	1.9	-	-	-	-	-	-	1.9	1.3
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	1,330	412	-	-	-	-	-	-	316	363
Total Phosphorus	NS	NS	NS	NS	NS	NS	3,760	1,200	-	-	-	-	-	-	4,340	1,080
рН	NS	NS	NS	NS	NS	NS	6.44	6.76	-	-	-	-	-	-	6.63	6.64
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	3.4	_	_	_	_	_	_	_	_	13.2
Gross beta	NS	NS	NS	NS	NS	NS	3.35	_	_	_	_	_	_	_	_	8.5
Radium 226	5 ^(surface) . 15					5 ^(surface) , 15	-	_	_	_	_	_	_	_	_	2.97
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS NS	NS NS	NS NS	15	-	-	-	-	-	_	-	-	-	0.74 U

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated in the control of the contr

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-73	SB-74	SB-74							
Sample Type:		Protection of	of Environmen	ital Quality ^(b)	Background		INVE	INVE	INVE							
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values (g)</u>	1-2	4-5	9-10	14-15	19-20	24-25	29-30	31.5-32.5	0-1	1-2

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluori

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimur

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cui The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cui

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Daalaman d		SB-74 DUP	SB-74 INVE	SB-74 INVE	SB-74 INVE	SB-74 INVE	SB-74 INVE	SB-74 INVE	SB-75 INVE	SB-75 INVE	SB-75 INVE
	la dinatalal				Background	0	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative										
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	I IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	1-2	4-5	9-10	14-15	19-20	24-25	29-30	0-1	1-2	4-5
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.51 U	1.2 U	-	-	-	-	-	0.39 U	0.39 U	0.6 U
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	3.9	5.5 J	2.8	5.3	5.8	6.8	NR	3.9	4.1	4.5
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	59.5	154	-	-	-	-	-	119	139	117
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.45	0.81 J	-	-	-	-	-	0.5 J	0.58 J	0.66 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	5.1	1.6 J	-	-	-	-	-	1	0.39 J	0.3 U
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	65,100	23,900	-	-	-	-	-	35,200	4,730	4,500
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	44.4	41.4	-	-	-	-	-	24.1	18.1	23.6
Lead	80	NS/14	50	GWP	NS/14	13.59	5.7	12.2	-	-	-	-	-	9.3	11.9	12.1
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	18,300	4,000	-	-	-	-	-	10,100	3,660	4,600
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	26.1	16.6	-	-	-	-	-	15.9	15.5	18.8
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	1,330	4,020 J	-	-	-	-	-	2,460	3,180	2,830 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	2.7	2.4 U	-	-	-	-	-	0.78 ∪	0.77 ∪	1.2 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	190 U	440 U	-	-	-	-	-	150 U	150 U	220 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.1 U	1.2 U	-	-	-	-	-	0.78 ∪	0.077 U	0.12 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	53.7	56.1	-	-	-	-	-	32.3	22.4	28.2
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	86	96.7	7.4	7	5.9	2.8	2.2	62.1	100	49.3
Fluoride	4,100	NS	7.4	GWP	3.95	7	43.9	47.2	1.2	1.1 J	1.1 J		0.82 J		18	13
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	1.3	4.4	_	-	-	_	-	1.5	1.3	1.3
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	357	1,230	_	-	_	_	-	1,120	1,640	781
Total Phosphorus	NS	NS	NS	NS	NS	NS	2,930	1,100	_	-	-	_	-	686	515	1,150
рН	NS	NS	NS	NS	NS	NS	6.1	6.59	-	-	-	-	-	6.09	6.82	6.09
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	_	_	_	_	_	5.1	_
Gross beta	NS	NS	NS	NS	NS	NS	-	-	_	-	-	_	-	_	4.8	_
Radium 226	5 ^(surface) , 15	-	-	-	-	5 ^(surface) , 15	-	-	_	-	-	_	_	_	-	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	_

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathw GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised are considered as updated and revised.

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-74	SB-75	SB-75	SB-75						
Sample Type:		Protection of	of Environmen	ital Quality ^(b)	Background		DUP	INVE								
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values (g)</u>	1-2	4-5	9-10	14-15	19-20	24-25	29-30	0-1	1-2	4-5

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluori

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimur

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-75 DUP	SB-75 INVE	SB-75 INVE	SB-75 INVE	SB-76 INVE	SB-76 INVE	SB-76 INVE	SB-76 INVE	SB-76 INVE	SB-76 INVE
Sampling Date: Depth:	Industrial Soil (mg/kg)	Groundwater ^(e) Risk-Based/MCL-Based	Risk Based	Critical Pathway	Concentrations (UTL 95-95) ^(f)	Comparative <u>Values ^(g)</u>	08/03/10 4-5	08/03/10 9-10	08/03/10 14-15	08/03/10 19-20	08/03/10 0-1	08/03/10 1-2	08/03/10 4-5	08/03/10 9-10	08/03/10 14-15	08/03/10 19-20
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.57 U	-	_	-	0.42 U	0.52 U	0.48 U	-	_	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	4.9	3.2	4.3	5.7	4.6	5.0	5.0	3.3	3.7	4.3
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	120	-	-	-	134	147	126	-	-	-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.7	-	-	-	0.63 J	0.58 J	0.69 J	-	-	-
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.29 U	-	-	-	0.28 J	0.75 J	0.24 U	-	-	-
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	4,860	-	-	-	3,910	8,320	3,940	-	-	-
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	25.3	-	-	-	19.9	19.9	23.9	-	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	12.1	-	-	-	11.9	12.4	12.1	-	-	-
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	5,020	-	-	-	3,610	4,260	4,310	-	-	-
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	19.2	-	-	-	15	16	18.5	-	-	-
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,960	-	-	-	3,180	3,400	2,760	-	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.1 U	-	-	-	0.84 ∪	1 U	0.96 ∪	-	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	220 U	-	-	-	160 U	190 U	248 J	-	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.11 U	-	-	-	0.084 U	0.1 U	0.096 U	-	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	30	-	-	-	24.9	25.6	27.8	-	-	-
Wet Chemistry (mg/kg)																
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	57.1	3.4	7.1	6.6	131	83.7	64.5	7.1	5.2	2.7
Fluoride	4,100	NS	7.4	GWP	3.95	7	17.1	8.5	23.4	5.8	9.6	9.2	7.1	-	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	1.3	-	-	-	0.6 U	1.3	1.4	-	-	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	785	-	-	-	1,590	908	753	-	-	-
Total Phosphorus	NS	NS	NS	NS	NS	NS	654	-	-	-	988	908	312	-	-	-
рН	NS	NS	NS	NS	NS	NS	6.29	-	-	-	6.72	5.08	6.35	-	-	-
Radiological (pCi/g)																
Gross alpha	NS	NS	NS	NS	NS	NS	_	_	_	_	_	5.4	_	_	_	_
Gross beta	NS	NS	NS	NS	NS	NS	-	-	_	_	_	4.0	-	-	_	-
Radium 226	5 ^(surface) , 15	-	-	-	-	5 ^(surface) , 15	_	-	_	-	_	-	-	-	_	-
-	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve	Idaho Dep	artment			SB-75	SB-75	SB-75	SB-75	SB-76	SB-76	SB-76	SB-76	SB-76	SB-76
Sample Type:		Protection of	of Environmen	ıtal Quality (b)	Background		DUP	INVE								
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10	14-15	19-20

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluori

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimur

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	ional Screening Leve Protection of		epartment ental Quality ^(b)	Background		SB-76 INVE	SB-77 INVE						
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations	Comparative		08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	<u>Values ^(g)</u>	21.5-22.5	0-1	1-2	4-5	9-10	14-15	19-20	24-25
Metals (mg/kg)														
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.54 U	0.51 U	0.43 U	0.5 U	_	_	_	_
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	5.2	5.2	3.3	4.9	4.3	4.2	4.3 J	5.1
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	130	51.6	61.4	136				-
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.7 J	0.28 J	0.28 J	0.62 J	_	_	_	_
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.47 J	4.1	3.5	0.84 J	_	_	_	_
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	5,200	114,000	51,200	20,300	_	_	_	_
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	21	35.1	34.6	25.9	-	_	-	-
Lead	80	NS/14	50	GWP	NS/14	13.59	10.7	4.7	6.5	11.6	_	_	-	_
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	5,080	13,200	10,500	4,890	_	_	-	_
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	20.5	23.8	18	17.5	_	_	-	_
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,340 J	1,310 J	1,580 J	3,120	-	-	-	-
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	1.1 U	1.8 J	0.89 J	1 U	-	-	-	-
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	385 J	190 U	221 J	269 J	-	-	-	-
Thallium	í	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	1.1 U	0.1 U	0.085 U	0.1 U	-	-	-	-
Vanadium	520	78/NS	NS	NS	78/NS	22.68	30.1	45.5	41.3	33.9	-	-	-	-
Wet Chemistry (ma/ka)														
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	18.2	22.4	24.9	73.8	30.4	9.4	10.5	1.8
Fluoride	4,100	NS	7.4	GWP	3.95	7	1.4	3.0	4.7	5.7	-	-	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	5.4	2.0	0.54 U	1.5	_	_	_	_
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	187	369	766	1,170	_	_	_	_
Total Phosphorus	NS	NS	NS	NS	NS	NS	274	1,060	1,460	2,140	_	_	-	_
рН	NS	NS	NS	NS	NS	NS	5.96	7.13	7.44	7.77	-	-	-	-
Radiological (pCi/g)														
Gross alpha	NS	NS	NS	NS	NS	NS			11					
Gross beta	NS	NS NS	NS	NS	NS	NS NS	_	_	6.6	_	_	_	_	_
Radium 226	5 ^(surface) , 15	140	INO	INO	110	5 ^(surface) , 15	-	_	2.57	-	_	_	_	_
Nadialli 220	5 (subsurface)	NC	NC	NC	NC	5 (subsurface)			2.07					
Dadium 000		NS NG	NS	NS	NS				0.7.11					
Radium 228	15	NS	NS	NS	NS	15	-	-	0.7 U	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity; "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised. EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated in the control of the contr

Soil Sampling Analytical Results - 2010 Production Well NW-9 QC Lab Leach Field Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-76	SB-77						
Sample Type:		Protection of	of Environmen	tal Quality ^(b)	Background		INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10	08/03/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	21.5-22.5	0-1	1-2	4-5	9-10	14-15	19-20	24-25

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluori

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

- c/ US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/ Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the huma
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimur

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cur

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment	Daalamaa a		SB-78 INVE	SB-78 DUP	SB-78 INVE	SB-78 INVE	SB-78 INVE	SB-78 INVE	SB-78 INVE	SB-79 INVE	SB-79 INVE	SB-79 INVE
Sample Type: Sampling Date:	المطار والمعارف	Protection of Groundwater ^(e)	Risk Based	ental Quality ^(b) Critical	Background Concentrations	Commonative		08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10
Depth:	Industrial Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	(UTL 95-95) ^(f)	Comparative Values (g)	0-1	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5
Берин.	Son (mg/kg)	RISK-Daseu/IVICL-Daseu	IDTES (IIIg/kg)	Failiway	(OIL 93-93)**	values	<u> </u>		1-2		3-10	14-13	13-20			
Metals (mg/kg)																
Antimony	99,000	0.27/0.27	4.8	GWP	0.27/0.27	0.50	0.41 U	0.12	0.3 J	0.4 J	-	-	-	0.99 ∪	1.1 U	0.36 J
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	0.0013/0.29	4.7	2.8	2.7	1.4	0.33 J	-	-	-	4.4 J	3.7 J	0.39 J
Barium	1.6	1,200/82	896	GWP	1,200/82	170.2	130	133	82.1	188	-	-	-	120	170	76.3
Beryllium	19,000	13/3.2	1.63	GWP	13/3.2	0.89	0.57 J	0.66	0.25	0.14 J	-	-	-	0.85 J	0.86 J	0.13 J
Cadmium	200	NS/NS	1.4	GWP	NS/NS	0.869	0.53 J	0.52	4	0.072 J	-	-	-	1 J	0.76 J	0.27
Calcium	80	NS/NS	NS	NS	NS/NS	38,270 (I)	38,300	42,700	165,000	394,000	-	-	-	31,600	37,800	202,000
Chromium	(j)	NS/180,000	2,135	GWP	NS/180,000	18.61	13.9	15.5	20.1	1.3 J	-	-	-	24.8	25.7	2.2 J
Lead	80	NS/14	50	GWP	NS/14	13.59	7.8	9.3	8	0.58 J	-	-	-	10	12.5	0.87 J
Magnesium	(j)	NS/NS	NS	NS	NS/NS	7,146	6,390	7,260	-	4,530	-	-	-	10,300	7,230	2,220
Nickel	2,000	20/NS	59	GWP	20/NS	15.71	12.6	14.6	8.5	0.93 J	-	-	-	20.7	18.3	2.6
Potassium	(j)	NS/NS	NS	NS	NS/NS	3,393 (I)	2,010 J	2,000	1,330	144 J	-	-	-	2,370 J	2,970 J	337 J
Selenium	510	0.4/0.26	2.0	GWP	0.4/0.26	1.04	0.82 ∪	0.21 U	1.5 ∪	4.5 ∪	-	-	-	2 U	2.2 U	2.1 U
Sodium	(j)	NS/NS	NS	NS	NS/NS	289 (I)	150 U	160 U	59.7 J	113 J	-	-	-	370 U	410 U	39 U
Thallium	1	0.0011/0.14	1.6	GWP	0.0011/0.14	NS	0.082 U	0.11 U	0.29 U	0.45 ∪	-	-	-	0.099 U	0.11 U	0.1 U
Vanadium	520	78/NS	NS	NS	78/NS	22.68	18.6	20.4	15.3	1 J	-	-	-	26.7	29.1	2.5 J
Mat Chamiator (mar/km)																
Wet Chemistry (mg/kg) Ammonia (as N)	NS	NS	4.1	Subsurface Soil	NS	NS	21.8	24.7	84.5	6.5	2.5	6.8	1.6	8.9	46.4	50.5
Fluoride	4,100	NS	7.4	GWP	3.95	7	2.4	0.54	7	5.4		-	-	3.8	1.4	0.6 U
Nitrate as N	1,600,000	NS	18.4	GWP	NS	NS	1.9	2.7	4.2	7.3	_	_	_	0.52 U	2.0	3.3
Total Kjeldahl Nitrogen	NS	NS	NS	NS	NS	NS	825	446	2,710	301	_	_	_	355	991	1,440
Total Phosphorus	NS	NS	NS	NS	NS	NS	1,030	2,520	374	104	_	_	_	109	429	484
рН	NS	NS	NS	NS	NS	NS	8.02	7.83	7.3	8.03	-	-	-	7.78	7.66	7.62
Dedictorical (nCi/n)																
Radiological (pCi/g)	NC	214	NC	NC	NC	NC			4.7						4.4	
Gross alpha	NS	NS NS	NS	NS	NS NC	NS	-	-	4.7	-	-	-	-	-	4.4	-
Gross beta	NS - (surface)	IND	NS	NS	NS	NS - (surface)	-	-	6.3	-	-	-	-	-	3.7	-
Radium 226	5 ^(surface) , 15					5 ^(surface) , 15	-	-	2.55	-	-	-	-	-	-	-
	(subsurface)	NS	NS	NS	NS	(subsurface)										
Radium 228	15	NS	NS	NS	NS	15	-	-	0.91	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised. SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment			SB-78	SB-79	SB-79	SB-79						
Sample Type:		Protection of	of Environmen	ital Quality (b)	Background		INVE	DUP	INVE							
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	Concentrations	Comparative	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	ed IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/

- c/ Surface soil.
- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/
- f/ background as per EPA coresponance dated April 19, 2012, except where noted)
- g/Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV
- h/ provided by Accutest Laboratories.
- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Table D-11

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	ional Screening Leve		epartment	SB-79	SB-79	SB-79	SB-80	SB-80	SB-80	SB-80	SB-80	SB-81	SB-81	SB-81	SB-81	SB-81
Sample Type:		Protection of	of Environme	ental Quality (b)	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE	INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	d IDTLs (mg/kg)	Pathway	9-10	14-15	16-17	0-1	1-2	4-5	9-10	13-14	0-1	1-2	4-5	9-10	14-15
Metals (mg/kg)																	
Antimony	99,000	0.27/0.27	4.8	GWP	-	-	-	0.87 ∪	0.44 U	0.47 J	-	-	0.15 J	0.14 J	0.12 J	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	-	-	-	4 J	3	1.7	4.7	2.3	1.7	1.2	1	2.1 J	2.2
Barium	1.6	1,200/82	896	GWP	-	-	-	129	118	128	-	-	91.7	81.2	84.8	-	-
Beryllium	19,000	13/3.2	1.63	GWP	-	-	-	0.86 J	0.58 J	0.31	-	-	0.54	0.46	0.5 J	-	-
Cadmium	200	NS/NS	1.4	GWP	-	-	-	1 J	0.43 J	3.3	0.071 J	0.051 U	0.59	0.3	0.28	-	-
Calcium	80	NS/NS	NS	NS	-	-	-	41,900	9,370	245,000	-	-	38,200	106,000	49,400	-	-
Chromium	(j)	NS/180,000	2,135	GWP	-	-	-	20.7	15.8	11.1	-	-	10.9	15.20	12.8	-	-
Lead	80	NS/14	50	GWP	-	-	-	10	8.7	3.8	-	-	7.2	6.6	6.6	-	-
Magnesium	(j)	NS/NS	NS	NS	-	-	-	11,200	4,000	3,920	-	-	2,890	3,130	6,900	-	-
Nickel	2,000	20/NS	59	GWP	-	-	-	16.8	13.4	6.1 J	-	-	9.9	8.7	9.1	-	-
Potassium	(j)	NS/NS	NS	NS	-	-	-	2,410 J	2,080 J	1,450	-	-	2,660	1,460	1,080 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	-	-	-	1.7 U	0.88 ∪	3.5 ∪	-	-	7.9	1.5 ∪	0.6 ∪	-	-
Sodium	(j)	NS/NS	NS	NS	-	-	-	330 U	170 U	94.7 J	-	-	59 U	50.2 J	110 U		-
Thallium	1	0.0011/0.14	1.6	GWP	-	-	-	0.087 U	0.088 U	0.35 U	-	-	0.078 U	0.076 U	0.076 U	-	-
Vanadium	520	78/NS	NS	NS	-	-	-	23.2	18.7	11.4	-	-	12.6	8.4	11.1	-	-
Wet Chemistry (mg/kg)																	
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	4	2.7	3.1	23.3	40.8	53.7	0.88 J	0.66 J	76	49.8	10	13.9	3
Fluoride	4,100	NS	7.4	GWP	-	-	-	3.1	1.1	10.8	0.93 J	0.86 J	19.2	4.3	1.6	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	-	-	-	4.3	2.1	4.7	-	-	2.9	2.1	2	-	-
Total Kjeldahl Nitrogen	NS	NS	NS	NS	-	-	-	628	969	1,950	-	-	1,850	1,090	329	-	-
Total Phosphorus	NS	NS	NS	NS	-	-	-	545	804	641	-	-	585	1,100	301	-	-
рН	NS	NS	NS	NS	-	-	-	7.86	7.82	7.65	-	-	7.16	7.92	8.11	-	-
Radiological (pCi/g)																	
Gross alpha	NS	NS	NS	NS	_	_	_	_	6.1	_	_	_	_	4.0	_	_	_
Gross beta	NS	NS	NS	NS	_	_	_	_	4.6	_	_	_	_	4.0	_	_	_
Radium 226	5 ^(surface) , 15				-	-	-	-	-	-	-	-	-	-	-	-	-
	(subsurface)	NS	NS	NS													
Radium 228	15	NS	NS	NS	-	-	-	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL base "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicat GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = tresponsible of the standard not developed; SSLs = soil screening levels UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As upd SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and W EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As upc EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Wate

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment	SB-79	SB-79	SB-79	SB-80	SB-80	SB-80	SB-80	SB-80	SB-81	SB-81	SB-81	SB-81	SB-81
Sample Type:		Protection of	of Environmen	tal Quality (b)	INVE												
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	ed IDTLs (mg/kg)	Pathway	9-10	14-15	16-17	0-1	1-2	4-5	9-10	13-14	0-1	1-2	4-5	9-10	14-15

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurem

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to presoil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/c/ Surface soil.

- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/f/background as per EPA coresponance dated April 19, 2012, except where noted) g/Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compa
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified ε
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium cal m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published sc

Table D-11

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve		epartment	SB-81 INVE	SB-81 INVE	SB-82 INVE	SB-82 INVE	SB-82 DUP	SB-82 INVE	SB-82 INVE	SB-82 INVE	SB-82 INVE	SB-83 INVE	SB-83 INVE	SB-83 INVE	SB-83 INVE
Sample Type: Sampling Date:	lu de atrial	Protection of		ental Quality (b)	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10
Depth:	Industrial	Groundwater (e)	Risk Based	Critical	19-20	22-23	06/11/10	1-2	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5	5-6
рерии.	Soil (mg/kg)	Risk-Based/MCL-Base	ed IDTES (mg/kg)	Pathway	19-20	22-23			1-2		9-10	14-13	19-20				
Metals (mg/kg)																	
Antimony	99,000	0.27/0.27	4.8	GWP	-	-	1.6 ∪	1.8 U	2 U	0.64 J	-	-	-	0.95 ∪	2 U	0.37 J	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	3.6	6.7	3.3 J	3.9 J	4.4	0.76	0.72	0.94	3.7	10.7	9 J	1.4 J	2.7
Barium	1.6	1,200/82	896	GWP	-	-	64.9	60.8 J	56.5	115	-	-	-	170	71 J	82.7	-
Beryllium	19,000	13/3.2	1.63	GWP	-	-	1.6 J	0.92 U	1 U	1.5	-	-	-	0.72 J	1 U	0.13 J	-
Cadmium	200	NS/NS	1.4	GWP	-	-	13.3	16.3	12.2	1.1	-	-	-	5.1	102	0.22	-
Calcium	80	NS/NS	NS	NS	-	-	183,000	141,000	175,000	329,000	-	-	-	213,000	162,000	284,000	-
Chromium	(j)	NS/180,000	2,135	GWP	-	-	70.2	117	70.7	11	-	-	-	31.2	211	1.4 J	-
Lead	80	NS/14	50	GWP	-	-	5.8 J	6.1	8.1	1.1 J	-	-	-	7.4 J	6.4	0.35 J	-
Magnesium	(j)	NS/NS	NS	NS	-	-	1,920 J	1,360 J	979	2,940	-	-	-	5,080	2,120 J	2,550	-
Nickel	2,000	20/NS	59	GWP	-	-	12 J	20.6	13.1	6.1 J	-	-	-	17.1 J	56.2	0.68 J	-
Potassium	(j)	NS/NS	NS	NS	-	-	1,120 J	820 J	693	474 J	-	-	-	2,510 J	1,990 J	401 J	-
Selenium	510	0.4/0.26	2.0	GWP	-	-	8.4 J	6.4 J	4.9	3.9 ∪	-	-	-	1.9 U	8.4 J	3.3 ∪	-
Sodium	(j)	NS/NS	NS	NS	-	-	1,200 U	690 U	760 U	104 J	-	-	-	647 J	1,140 J	405 J	-
Thallium	1	0.0011/0.14	1.6	GWP	-	-	1.6 ∪	0.092 U	2 U	0.39 U	-	-	-	0.95 ∪	0.82	0.33 ∪	-
Vanadium	520	78/NS	NS	NS	-	-	74.4	132	75	11.9	-	-	-	56.5	302	1.4 J	-
Wet Chemistry (mg/kg)																	
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	2.4	1.3 J	17.3	13.7	17.7	19.1	6.7	94	125	135	161	82.3	58.9
Fluoride	4,100	NS	7.4	GWP	-	-	45.9	42.7	42.7	3.2	-	-	-	25.1	9.2	0.93 J	-
Nitrate as N	1,600,000	NS	18.4	GWP	-	-	0.53 U	11.2	0.53 U	20.3	22.1	1.6	0.59 U	0.55 U	0.58 U	0.58 U	_
Total Kjeldahl Nitrogen	NS	NS	NS	NS	-	-	343	611	338	367	-	-	-	851	911	454	-
Total Phosphorus	NS	NS	NS	NS	-	-	2,390	1,330	3,280	617	-	-	-	852	6,350	468	-
рН	NS	NS	NS	NS	-	-	6.95	6.17	6.7	7.4	-	-	-	7.35	5.45	8.06	-
Radiological (pCi/g)																	
Gross alpha	NS	NS	NS	NS	-	-	-	67	_	_	-	-	-	_	15.3	_	_
Gross beta	NS	NS	NS	NS	-	-	-	26.8	-	-	-	-	-	-	9.6	-	-
Radium 226	5 ^(surface) , 15				-	-	-	15.6	-	-	-	-	-	-	-	-	-
	(subsurface)	NS	NS	NS													
Radium 228	15	NS	NS	NS	-	-	-	0.3 U	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL base "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicat GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = tare "NS" = not applicable or standard not developed; SSLs = soil screening levels UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As upd SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and W EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As upc EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Wate

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment	SB-81	SB-81	SB-82	SB-83	SB-83	SB-83	SB-83						
Sample Type:		Protection of	of Environmen	tal Quality (b)	INVE	INVE	INVE	INVE	DUP	INVE							
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	ed IDTLs (mg/kg)	Pathway	19-20	22-23	0-1	1-2	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5	5-6

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurem

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to presoil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/c/Surface soil.

- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/f/background as per EPA coresponance dated April 19, 2012, except where noted) g/Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compa
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified ε
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium cal m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published sc

Table D-11

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Path	Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	SB-84 INVE	SB-84 INVE	SB-84 INVE	SB-84 INVE	SB-85 INVE	SB-85 INVE	SB-85 INVE	SB-85 INVE	SB-86 INVE	SB-86 DUP	SB-86 INVE	SB-86 DUP	SB-86 INVE
Antimony 99,000 0.27/0.27	Sampling Date:		Groundwater ^(e)	Risk Based	Critical													08/12/10 4-5
Arsenic 41 0.0013/0.29 0.39 Surficial Soil 3.9 J 3.3 J 5.9 4.9 3.6 J 7.1 4.4 1.3 5.1 5.1 5.1 6.6 J 6.1 Sarium 1.6 1.2018/22 896 GWP 160 140 133 - 132 264 150 - 152 J 144 192 181 3 Beryllium 19,000 13/3.2 1.63 GWP 1.J 1.J 0.67 J - 0.77 J 1.8 J 1.6 0.13 J 1.4 U 0.73 0.9 U 0.57 U 0.0 Cadmium 200 NSNS 1.4 GWP 1.2 J 0.96 J 0.66 J - 1.3 J 7.9 1.3 - 4.J 1.6 7.5 7.8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Metals (mg/kg)																	
Barulm	Antimony	99,000	0.27/0.27	4.8	GWP	0.81 ∪	0.98 ∪	0.95 ∪	-	0.79 ∪	0.94 ∪	0.47 J	-	2.8 ∪	0.98 ∪	1.8 U	1.1 U	2 U
Belgillum 19,000 13/3.2 16.3 GWP 1.J 1.J 0.67.J - 0.77.J 1.8.J 1.6 0.13.J 1.4.U 0.73 0.9.U 0.57.U 0.7.U 0.6.d 0.	Arsenic	41	0.0013/0.29	0.39	Surficial Soil	3.9 J	3.3 J	5.9	4.9	3.6 J	7.1	4.4	1.3	5.1 J	5.1	6.6 J	6.1	5 J
Cadmium 200 NS/NS 1.4 GWP 1.2 J 0.96 J 0.66 J - 1.3 J 7.9 1.3 - 4 J 1.6 7.5 7.8 1: Calcium 80 NS/NS NS NS NS 7.110 6,620 68,500 - 6,140 107,000 241,000 - 6,270 J 8,320 5,810 6,120 19,6 Chromium (j) NS/180,000 2,135 GWP 22 24.2 20.2 - 22.9 45.5 28 - 44.3 26.9 69.1 53.8 81 Lead 80 NS/14 50 GWP 11 11.4 7.8 - 19,9 10.4 3.3 - 12 11.5 11.9 12.3 11 Magnesium (j) NS/NS NS NS NS NS NS NS 5,020 5,280 5,360 - 3,870 4,750 2,110 - 3,200 J 3,910 2,560 J 2,790 4,0 Nickel 2,000 20/NS 59 GWP 17.6 17.5 14.1 - 15.3 21 13.6 - 12 14.7 11.1 13 Potassium (j) NS/NS NS NS NS NS 4,650 3,960 J 2,130 J - 3,480 J 4,720 916 - 4,350 J 3,740 4,960 J 4,760 9.1 Selenium 510 0.4/0.26 2.0 GWP 1.6 U 2 U 1.9 U - 1.6 U 1.9 U 3.4 U - 5.6 U 2 U 3.6 U 2.3 U 3. Sodium (j) NS/NS NS N	Barium	1.6	1,200/82	896	GWP	160	140	133	-	132	264	150	-	152 J	144	192	181	308
Calcium 80 NS/NS NS NS NS 7,110 6,620 68,500 - 6,140 107,000 241,000 - 6,270 J 8,320 5,810 6,120 19,6 Chromium (j) NS/180,000 2,135 GWP 22 24.2 20.2 - 22.9 45.5 28 - 44.3 26.9 69.1 53.8 81	Beryllium	19,000	13/3.2	1.63	GWP	1 J	1 J	0.67 J	-	0.77 J	1.8 J	1.6	0.13 J	1.4 U	0.73	0.9 U	0.57 U	0.98 U
Chromium (j)	Cadmium	200	NS/NS	1.4	GWP	1.2 J	0.96 J	0.66 J	-	1.3 J	7.9	1.3	-	4 J	1.6	7.5	7.8	11.4
Chromium (j)	Calcium	80	NS/NS	NS	NS	7,110	6,620	68,500	-	6,140	107,000	241,000	-	6,270 J	8,320	5,810	6,120	19,600
Magnesium (j) NS/NS NS NS S,	Chromium	(j)	NS/180,000	2,135	GWP	22	24.2	20.2	-		45.5	28	-	44.3	26.9		53.8	87.2
Nickel 2,000 20/NS 59 GWP 17.6 17.5 14.1 - 15.3 21 13.6 - 12 14.7 11.1 13 Potassium (i) NS/NS NS N	Lead	80	NS/14	50	GWP	11	11.4	7.8	-	19.9	10.4	3.3	-	12	11.5	11.9	12.3	10.2
Nicker 2,000 20/NS 59 GWP 17.6 17.5 14.1 - 15.3 21 13.6 - 12 14.7 11.1 13 Potassium 510 0.4/0.26 2.0 GWP 1.6 2 2 1.9 0 - 1.6 1.9 0 3.480 4.720 916 - 4.350 3.740 4.960 4.960 4.760 9.1 Selenium 510 0.4/0.26 2.0 GWP 1.6 2 2 1.9 0 - 1.6 1.9 0 3.4 0 - 5.6 2 2 3.6 2.3 0 3.5 Sodium (i) NS/NS NS NS NS NS 300 370 418 3 - 300 350 32 0 - 1.000 370 670 747 3.8 Thallium 1 0.0011/0.14 1.6 GWP 0.081 0.098 0.095 0 - 0.079 0.094 0.17 0 - 0.25 0.098 0.56 0.34 0.0 Vanadium 520 78/NS NS NS NS 25.2 25.5 31.8 - 24.6 61.6 11.6 - 48.1 32.3 70 66.5 1.4 Wet Chemistry (mg/kg) Armonia (as N) NS NS NS 7.4 GWP 5 5.6 7 - 50.5 30.1 1.3 - 436 136 1.310 1.330 2.3 Nitrate as N 1,600,000 NS 18.4 GWP 26.5 25.2 21.3 11.9 2.2 5.7 5.1 - 0.54 2.4 0.8 0.15 Total Kjeldahl Nitrogen NS NS NS NS NS NS NS N	Magnesium	(j)	NS/NS	NS	NS	5,020	5,290	5,360	-	3,870	4,750	2,110	-	3,200 J	3,910	2,560 J	2,790	4,070 J
Selenium		2,000	20/NS	59	GWP	17.6	17.5	14.1	-	15.3	21	13.6	-	12	14.7	11.1	13	13
Selenium S10 0.4/0.26 2.0 GWP 1.6 U 2 U 1.9 U - 1.6 U 1.9 U 3.4 U - 5.6 U 2 U 3.6 U 2.3 U 5.5 Sodium (i) NS/NS NS NS NS NS NS NS N	Potassium	(i)	NS/NS	NS	NS	4,650	3,960 J	2,130 J	-	3,480 J	4,720	916	-	4,350 J	3,740	4,960 J	4,760	9,100 J
Sodium (j) NS/NS	Selenium		0.4/0.26	2.0	GWP	1.6 ∪	2 U	1.9 ∪	-	1.6 ∪	1.9 ∪	3.4 U	_	5.6 ∪	2 U	3.6 ∪	2.3 ∪	
Thallium 1 0.0011/0.14 1.6 GWP 0.081 U 0.098 U 0.095 U - 0.079 U 0.094 U 0.17 U - 0.25 J 0.098 U 0.56 0.34 0.00 Vanadium 520 78/NS NS NS NS S NS S S S S S S S S S S S		(i)			NS				-									3,890 J
Wet Chemistry (mg/kg) NS NS NS 25.2 25.5 31.8 - 24.6 61.6 11.6 - 48.1 J 32.3 70 66.5 1 Wet Chemistry (mg/kg) S S 4.1 Subsurface Soil 70.9 52 8.2 3.8 61.3 79.6 37 1.4 98.2 71.5 216 270 6 Fluoride 4,100 NS 7.4 GWP 5 5.6 7 - 50.5 30.1 1.3 - 436 136 1,310 1,330 2,3 Nitrate as N 1,600,000 NS 18.4 GWP 26.5 25.2 21.3 11.9 2.2 5.7 5.1 - 0.54 U 2.4 0.58 U 5.5 0. Total Kjeldahl Nitrogen NS NS NS NS 1,860 1,920 281 - 1,500 2,280 1,210 - 1,010 1,500 848 1010	Thallium	1	0.0011/0.14	1.6	GWP	0.081 U	0.098 U	0.095 U	-	0.079 U	0.094 U	0.17 ∪	_	0.25 J	0.098 U	0.56	0.34	0.098 U
Ammonia (as N) NS NS NS 4.1 Subsurface Soil 70.9 52 8.2 3.8 61.3 79.6 37 1.4 98.2 71.5 216 270 6 Fluoride 4,100 NS 7.4 GWP 5 5.6 7 - 50.5 30.1 1.3 - 436 136 1,310 1,330 2,3 Nitrate as N 1,600,000 NS 18.4 GWP 26.5 25.2 21.3 11.9 2.2 5.7 5.1 - 0.54 U 2.4 0.58 U 5.5 0. Total Kjeldahl Nitrogen NS		520	78/NS	NS	NS				-									134
Ammonia (as N) NS NS NS 4.1 Subsurface Soil 70.9 52 8.2 3.8 61.3 79.6 37 1.4 98.2 71.5 216 270 6 Fluoride 4,100 NS 7.4 GWP 5 5.6 7 - 50.5 30.1 1.3 - 436 136 1,310 1,330 2,3 Nitrate as N 1,600,000 NS 18.4 GWP 26.5 25.2 21.3 11.9 2.2 5.7 5.1 - 0.54 U 2.4 0.58 U 5.5 0. Total Kjeldahl Nitrogen NS	Wet Chemistry (ma/ka)																	
Fluoride 4,100 NS 7.4 GWP 5 5.6 7 - 50.5 30.1 1.3 - 436 136 1,310 1,330 2,3 Nitrate as N 1,600,000 NS 18.4 GWP 26.5 25.2 21.3 11.9 2.2 5.7 5.1 - 0.54 U 2.4 0.58 U 5.5 0. Total Kjeldahl Nitrogen NS NS NS NS NS 1,860 1,920 281 - 1,500 2,280 1,210 - 1,010 1,500 848 1010 1,0 Total Phosphorus NS NS NS NS NS NS 1,350 663 543 - 290 290 320 - 3,150 582 8,350 2,940 2,8 pH NS		NS	NS	4.1	Subsurface Soil	70.9	52	8.2	3.8	61.3	79.6	37	1.4	98.2	71.5	216	270	616
Nitrate as N 1,600,000 NS 18.4 GWP 26.5 25.2 21.3 11.9 2.2 5.7 5.1 - 0.54 U 2.4 0.58 U 5.5 0. Total Kjeldahl Nitrogen NS NS NS NS 1,860 1,920 281 - 1,500 2,280 1,210 - 1,010 1,500 848 1010 1,0 Total Phosphorus NS NS NS NS 1,350 663 543 - 290 290 320 - 3,150 582 8,350 2,940 2,8 pH NS NS NS NS 7.04 7.12 7.86 - 7.26 7.47 7.83 - 5.32 6.38 4.77 4.78 4.	` ,				GWP													2,370
Total Kjeldahl Nitrogen NS NS NS NS 1,860 1,920 281 - 1,500 2,280 1,210 - 1,010 1,500 848 1010 1,010 Total Phosphorus NS NS NS NS NS 1,350 663 543 - 290 290 320 - 3,150 582 8,350 2,940 2,8 pH NS NS NS NS 7.04 7.12 7.86 - 7.26 7.47 7.83 - 5.32 6.38 4.77 4.78 4.	Nitrate as N			18.4	GWP	26.5		21.3	11.9		5.7		-	0.54 U				0.58 U
Total Phosphorus NS NS NS NS 1,350 663 543 - 290 290 320 - 3,150 582 8,350 2,940 2,8 pH NS NS NS NS 7.04 7.12 7.86 - 7.26 7.47 7.83 - 5.32 6.38 4.77 4.78 4.	Total Kieldahl Nitrogen												-	1.010				1,040
pH NS NS NS NS 7.04 7.12 7.86 - 7.26 7.47 7.83 - 5.32 6.38 4.77 4.78 4.									_	•			-	•				2,880
			NS	NS	NS				-				-					4.53
Kadiological (pCl/g)	Radiological (pCi/g)																	
Gross alpha NS NS NS - 5.7 8.7 15.4 13.8		NS	NS	NS	NS	_	5.7	_	_	_	8.7	_	_	_	_	15.4	13.8	_
Gross beta NS NS NS NS - 5.6 7.5 14.2 15.8	•					-		-	_	-		-	-	-	_			-
Radium 226 5 (surface), 15						-		_	_	_	-	_	_	_	_			-
(subsurface) NS NS NS		•	NC	NS	NC													
Radium 228 15 NS NS NS 0.92 1.28	Radium 228					_	_	_	_	_	_	_	_	_	_	0.92	1.28	_

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL base "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicat GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = ta "NS" = not applicable or standard not developed; SSLs = soil screening levels UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As upd SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and W EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As upc EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Wate

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment	SB-84	SB-84	SB-84	SB-84	SB-85	SB-85	SB-85	SB-85	SB-86	SB-86	SB-86	SB-86	SB-86
Sample Type:		Protection of	of Environmer	tal Quality (b)	INVE	DUP	INVE	DUP	INVE								
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/11/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	ed IDTLs (mg/kg)	Pathway	0-1	1-2	4-5	9-10	0-1	1-2	4-5	8-9	0-1	0-1	1-2	1-2	4-5

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurem

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to presoil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/c/Surface soil.

- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/f/background as per EPA coresponance dated April 19, 2012, except where noted) g/Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compa
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified ε
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium cal m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published sc

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type:	Region 9 Regi	onal Screening Leve Protection of		epartment ental Quality ^(b)	SB-86 DUP	SB-86 INVE	SB-86 INVE	SB-86 INVE	SB-87 INVE	SB-87 INVE	SB-87 INVE	SB-87 INVE	SB-87 INVE
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based		Pathway	4-5	9-10	14-15	18-19	0-1	1-2	4-5	9-10	10-11
•		2000402 20004											
Metals (mg/kg)													
Antimony	99,000	0.27/0.27	4.8	GWP	1.8 ∪	-	-	-	0.81 ∪	0.095 U	0.54 U	-	-
Arsenic	41	0.0013/0.29	0.39	Surficial Soil	5.9	4.6	3.9	2.9	2.4 J	2.2	5	3.9	4.9
Barium	1.6	1,200/82	896	GWP	289	199	110	74.6	121	107	111	-	-
Beryllium	19,000	13/3.2	1.63	GWP	0.89 U	-	-	-	0.71 J	0.58 J	0.75 J	-	-
Cadmium	200	NS/NS	1.4	GWP	13.9	-	-	-	1.7	0.35	0.63 J	-	-
Calcium	80	NS/NS	NS	NS	14,900	-	-	-	77,700	90,200	43,200	-	-
Chromium	(j)	NS/180,000	2,135	GWP	85.3	-	-	-	22.5	14.4	18.4	-	-
Lead	80	NS/14	50	GWP	10.4	-	-	-	8.7	7.6	9.2	-	-
Magnesium	(j)	NS/NS	NS	NS	3,170	-	-	-	5,340	7,220	9,270	-	-
Nickel	2,000	20/NS	59	GWP	15.4	-	-	-	11.5	9.9	12.5	-	-
Potassium	(j)	NS/NS	NS	NS	7,820	-	-	-	2,950 J	1,650 J	1,810 J	-	-
Selenium	510	0.4/0.26	2.0	GWP	3.6 ∪	-	-	-	1.6 ∪	0.95 ∪	1.1 U	-	-
Sodium	(j)	NS/NS	NS	NS	2,310	-	-	-	300 U	140 U	200 U	-	-
Thallium	1	0.0011/0.14	1.6	GWP	0.089 U	-	_	-	0.081 U	0.095 U	0.11 U	-	-
Vanadium	520	78/NS	NS	NS	109	-	-	-	28.4	19.7	28.3	-	-
Wet Chemistry (mg/kg)													
Ammonia (as N)	NS	NS	4.1	Subsurface Soil	416	9.7	7.3	2.8	55.7	37	8.3	3.4	3.4
Fluoride	4,100	NS	7.4	GWP	2,280	8.5	1.3	1.6	48.6	58.9	2.7	-	-
Nitrate as N	1,600,000	NS	18.4	GWP	0.58 U		-	-	15.6	2.2	2.2	_	_
Total Kjeldahl Nitrogen	NS	NS	NS	NS	1,150	_	_	_	1,790	1,210	279	_	_
Total Phosphorus	NS	NS	NS	NS	2,660	_	_	_	4,070	4,070	1,170	_	_
рН	NS	NS	NS	NS	4.51	-	-	-	7.57	7.78	7.93	-	-
5 " 1 ' 1 ' 6" \													
Radiological (pCi/g)	NO	NO	NO	NO									
Gross alpha	NS	NS	NS	NS	-	-	-	-	-	3.8	-	-	-
Gross beta	NS (gurfage)	NS	NS	NS	-	-	-	-	-	3.9	-	-	-
Radium 226	5 ^(surface) , 15				-	-	-	-	-	-	-	-	-
	(subsurface)	NS	NS	NS									
Radium 228	15	NS	NS	NS	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

INVE = Investigative sample.

DUP = Blind duplicate.

MS = Matrix spike.

NS = No regulatory criteria established.

GWP = Groundwater protection.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL base "m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicat GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = tare "NS" = not applicable or standard not developed; SSLs = soil screening levels UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As upd SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and W EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As upc EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Wate

Soil Sampling Analytical Results - 2010 Gypstack Area Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID:	Region 9 Regi	onal Screening Leve	Idaho Dep	artment	SB-86	SB-86	SB-86	SB-86	SB-87	SB-87	SB-87	SB-87	SB-87
Sample Type:		Protection of	of Environmen	tal Quality ^(b)	DUP	INVE							
Sampling Date:	Industrial	Groundwater (e)	Risk Based	Critical	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10	08/12/10
Depth:	Soil (mg/kg)	Risk-Based/MCL-Based	d IDTLs (mg/kg)	Pathway	4-5	9-10	14-15	18-19	0-1	1-2	4-5	9-10	10-11

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurem

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to presoil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/c/ Surface soil.

- d/ Idaho Risk Reduction Manual, July 2004.
- e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/f/background as per EPA coresponance dated April 19, 2012, except where noted) g/Comparative values are defined as:
 - the lower of the human health screening levels, unless this value is below background
 - if the human health screening level is above background, then it is the CV
 - if the human health screening level is below background, then background is the CV

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compa
- k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified ε
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium cal m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published sc

Soil Sampling Analytical Results - 2010 Trip and Equipment Blanks Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type: Sampling Date: Depth (feet):	EB08041 INVE 8/4/2010 -	EB08051 INVE 8/5/2010	EB08061 INVE 8/6/2010	EB08091 INVE 8/9/2010	EB08101 INVE 8/10/2010	EB08111 INVE 8/11/2010	EB08121 INVE 8/12/2010	EB08131 INVE 8/13/2010	EB08161 INVE 8/16/2010	EB08171 INVE 8/17/2010	EB08181 INVE 8/18/2010	TB08041 TB 8/4/2010	TB08051 TB 8/5/2010	TB08061 TB 8/6/2010
Volatile Organic Compunds	s (mg/l)													
Benzene	0.00021 U	0.00021 U	0.00021 U	-	-	-	-	-	-	-	-	0.00021 U	0.001 U	0.00021 U
Ethylbenzene	0.0002 U	0.0002 U	0.0002 U	-	-	-	-	-	-	-	-	0.0002 U	0.0011 U	0.0002 U
Toluene	0.0002 U	0.0002 U	0.0002 U	-	-	-	-	-	-	-	-	0.0002 U	0.0012 U	0.0002 U
TPH-GRO (C6-C10)	0.05 U	0.05 U	0.05 U	-	-	-	-	-	-	-	-	-	-	-
Xylene, (total)	0.00054 U	0.00054 U	0.00054 U	-	-	-	-	-	-	-	-	0.00054 U	0.0022 U	0.00054 U
Metals (mg/l)														
Antimony	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.0023	0.0021	0.002 U	-	-	-
Arsenic	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	-	-	-
Barium	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	0.005 U	-	-	-
Beryllium	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	-	-	-
Cadmium	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	-	-	-
Calcium	0.103	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.633	0.1 U	0.104	0.627	0.183	-	-	-
Chromium	0.001 U	0.001 U	0.0047	0.001 U	0.001 U	0.001 U	0.0064	0.001 U	0.001 U	0.001 U	0.0055	-	-	-
Lead	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	-	-	-
Magnesium	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	-	-	-
Nickel	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	-	-	-
Potassium	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	0.5 U	-	-	-
Selenium	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U	0.002 U			0.0022	0.002 U	-	-	-
Sodium	0.75 U	0.75 U	0.75 U	0.75 U	1	0.75 U	0.75 U				1.43	-	-	-
Thallium	0.0018	0.0019	0.0015 U	0.0015 U	0.0015 U	0.0015 U	0.0015 U	0.0018	0.0015 U	0.0015 U	0.0015 U	-	-	-
Vanadium	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	0.001 U	-	-	-
Wet Chemistry (mg/l)														
Ammonia (as N)	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	0.05 U	-	-	-
Fluoride	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U				0.1 U	-	-	-
Nitrate as N	0.05 U	0.05 U	0.12	0.05 U	0.11	0.15	0.05 U	0.05 U		0.05 U	0.1	-	-	-
Total Kjeldahl Nitrogen	0.11 U	0.11 U	0.11 U	0.11 U	0.11 U	0.11 U	0.22	0.49	0.38	0.14	0.11 U	-	-	-
Total Phosphorus	0.027	0.009	0.009 U	0.009 U	0.009 U	0.009 U	0.009 U	0.009 U	0.009 U	0.009 U	0.009 U	-	-	-
рН	4.75	7.6	7.79	7.48	7.3	3.87	6.11	7.23	6.7	6.56	6.56	-	-	-
Radiological (pCi/l)														
Gross alpha	-	-	-	-	-	-	-	-	0.22 U		-	-	-	-
Gross beta	-	-	-	-	-	-	-	-	0.9 U	1.7 U	-	-	-	-
Radium 226	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Radium 228	-	-	-	-	-	-	-	-	-	-	-	-	-	-

U = Analyte not detected above method detection limit.
J = Estimated result.
- = Not analyzed.
INVE = Investigative sample.
DUP = Blind duplicate.
MS = Matrix spike.

Bold = Above criteria

Soil Sample Summary - 2012 Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs Idaho

Conda Phosphate Operations Facility
Soda Springs, Idaho

Areas of Interest and Sample Information

Chemical Parameters

General Rad Chem ^(c) Sample Type

Soil Boring Sample ID Depths Date Time Metals ^(a) pH Chemistry ^(b) Gross a & B Ra-226/228 (INVE/DUP)

Phosphoric and Super Phosphoric Acid Area

Solid Boring Sample ID Debts Date Date	7.1.000 01 111.01.001 011.0	Guii	ipio iiiioii	nation				General	Rad C	hem ^(c)	Sample Type	
North Car Wash Sump Are SB-88 (I-1)	Soil Boring	Sample ID	Depths	Date	Time	Metals ^(a)	рΗ					Additional Comments
North Car Wash Sump Area SB-88 (3-1)							<u> </u>	<u></u>	<u> </u>	- 10	<u>,</u>	<u></u>
SB-88 SB-88 C-1 0-1 0-1 09111/12 810 X X X X X X X X X	тисорионо ши оброги											
SB-88 SB-88 C-1 0-1 0-1 09111/12 810 X X X X X X X X X	North Car Wash Sump Are	a										
SB-88 (H-2) 1-2 09/11/12 815			0-1	09/11/12	810	Χ	Χ	Χ	X	X	INVE	
SB-88 (4-5) 4-5 09/11/12 840 X X X X X X X X X		, ,										
SB-88 (9-10) 9-10 09/11/12 1530 X X X X X X X X X		, ,										
SB-89 SB-89 (0-1)		, ,										Refusal at 11 feet bas (not bedrock).
SB-90 SB-90 (0-1)		()		00/ / . =								(constant of the constant of
SB-90 SB-90 (0-1)	SB-89	SB-89 (0-1)	0-1	09/11/12	1530	Χ	Χ	Χ	Χ	Χ	INVE	Hand auger refusal at 1 feet bgs on gravel, 4 attempts
SB-90 (1-2) 1-2 09/11/12 1250 X X X X X X X X X		, ,										
SB-90 (4-5)	SB-90	SB-90 (0-1)	0-1	09/11/12	1245	Χ	Χ	Χ	X	X	INVE	
SB-90 (9-10) 9-10 09/11/12 1310 X X X X X X X X X		SB-90 (1-2)	1-2	09/11/12	1250	Χ	X	X	X	X	INVE	
SB-90 (9-10) 9-10 09/11/12 1310 X X X X X X X X X		SB-90 (4-5)	4-5	09/11/12	1255			Χ	Χ		INVE	
SB-90 (14-15) 14-15 09/11/12 1310 X X X X X X X X X		, ,										
SB-91 (SB-91 (0-1) 0-1 09/11/12 1320 X X X X X X INVE SB-91 (1-2) 1-2 09/11/12 1325 X X X X X X INVE SB-91 (1-5) 4-5 09/11/12 1335 X X X X X X INVE SB-91 (1-15) 14-15 09/11/12 1335 X X X X X X INVE SB-91 (14-15) 14-15 09/11/12 1345 X X X X X INVE SB-92 (1-1) 0-1 09/11/12 1345 X X X X X INVE SB-92 (1-1) 0-1 09/11/12 1040 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1040 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1045 X X X X X INVE SB-92 (1-1) 0-1 09/11/12 1045 X X X X X INVE SB-92 (1-1) 0-1 09/11/12 1055 X X X X X X INVE SB-92 (1-1) 0-1 09/11/12 1055 X X X X X X INVE SB-92 (1-15) 14-15 09/11/12 1105 X X X X X X INVE SB-92 (1-15) 14-15 09/11/12 1105 X X X X X X INVE SB-92 (1-15) 09/11/12 1110 X X X X X X INVE SB-92 (1-16) 09/11/12 1110 X X X X X X INVE SB-92 (24-25) 09/11/12 1110 X X X X X X X INVE SB-92 (24-25) 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X X X X X X X INVE SB-93 (1-1) 0-1 09/11/12 1120 X X X X X X X X X X X X X X X X X X X		` ,										Refusal at 15 feet bas (not bedrock).
SB-91 (1-2) 1-2 09/11/12 13325 X X X X X X X INVE SB-91 (9-10) 9-10 09/11/12 1330 X X X X X X INVE SB-91 (9-10) 9-10 09/11/12 1335 X X X X X X INVE SB-91 (14-15) 14-15 09/11/12 1345 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1040 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1046 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1055 X X X X X X INVE SB-92 (9-10) 9-10 09/11/12 1055 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (14-25) 24-25 09/11/12 1110 X X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1110 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1120 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1110 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1120 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1150 X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X X X X X X X X		()		00/ / . =								, , , , , , , , , , , , , , , , , , , ,
SB-91 (1-2) 1-2 09/11/12 13325 X X X X X X X INVE SB-91 (9-10) 9-10 09/11/12 1330 X X X X X X INVE SB-91 (9-10) 9-10 09/11/12 1335 X X X X X X INVE SB-91 (14-15) 14-15 09/11/12 1345 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1040 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1046 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1055 X X X X X X INVE SB-92 (9-10) 9-10 09/11/12 1055 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (14-25) 24-25 09/11/12 1110 X X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1110 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1120 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1110 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1120 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 1100 X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1150 X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X X X X INVE SB-93 (1-15) 14-15 09/11/12 1000 X X X X X X X X X X X X X X X X X	SB-91	SB-91 (0-1)	0-1	09/11/12	1320	Χ	Χ	Χ	X	X	INVE	
SB-91 (4-5)		, ,										
SB-91 (9-10) 9-10 09/11/12 1335 X X X X X X X INVE SB-91 (14-15) 14-15 09/11/12 1345 X X X X X X X X INVE Refusal at 15 feet bgs (not bedrock). SB-92 SB-92 (0-1) 0-1 09/11/12 1040 X X X X X X INVE SB-92 (1-2) 1-2 09/11/12 1050 X X X X X X INVE SB-92 (1-2) 4-5 09/11/12 1050 X X X X X X INVE SB-92 (9-10) 9-10 09/11/12 1055 X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1110 X X X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1120 X X X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 110 X X X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 110 X X X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 110 X X X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 915 X X X X X X X X INVE SB-93 (1-10) 09/11/12 925 X X X X X X X X INVE SB-93 (1-10) 09/11/12 925 X X X X X X X X INVE SB-93 (1-10) 09/11/12 925 X X X X X X X X INVE SB-93 (1-10) 19-20 09/11/12 1020 X X X X X X X X INVE SB-93 (1-10) 19-20 09/11/12 1020 X X X X X X X X X INVE SB-93 (1-10) 19-20 09/11/12 1020 X X X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X X X INVE SB-94 (1-2) 19-20 09/11/12 1020 X X X X X X X X X X X INVE SB-94 (1-2) 19-20 09/11/12 1155 X X X X X X X X X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X X X X X X X X X X		` ,										
SB-91 (14-15) 14-15 09/11/12 1345 X X X X X X X X X X X X X X X X X X X		, ,										
SB-92 SB-92 (0-1) 0-1 09/11/12 1040 X X X X X X X X X		, ,										Refusal at 15 feet has (not hedrock)
SB-92 (1-2) 1-2 09/11/12 1045 X X X X X X X INVE SB-92 (4-5) 4-5 09/11/12 1050 X X X X X X INVE SB-92 (9-10) 9-10 09/11/12 1055 X X X X X X INVE SB-92 (19-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1115 X X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1120 X X X X X X INVE SB-93 (24-25) 24-25 09/11/12 1120 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 910 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 915 X X X X X X INVE SB-93 (4-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (14-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 925 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X X INVE SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X X X X X X X X X X		30-91 (14-13)	14-13	09/11/12	1343	^	^	^	^	^	IIN V L	rtelusal at 13 leet bys (not bedrock).
SB-92 (1-2) 1-2 09/11/12 1045 X X X X X X X INVE SB-92 (4-5) 4-5 09/11/12 1050 X X X X X X INVE SB-92 (9-10) 9-10 09/11/12 1055 X X X X X X INVE SB-92 (19-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1115 X X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1120 X X X X X X INVE SB-93 (24-25) 24-25 09/11/12 1120 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 910 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 915 X X X X X X INVE SB-93 (4-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (14-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 925 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X X INVE SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X X X X X X X X X X	SB-92	SB-92 (0-1)	0-1	09/11/12	1040	Χ	Χ	X	X	X	INVE	
SB-92 (4-5)		, ,										
SB-92 (9-10) 9-10 09/11/12 1055 X X X X X X X INVE SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X INVE SB-92 (19-20) 19-20 09/11/12 1115 X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1120 X X X X X INVE SB-93 (24-25) 24-25 09/11/12 1120 X X X X X INVE SB-93 (1-2) 1-2 09/11/12 910 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 915 X X X X X INVE SB-93 (9-10) 9-10 09/11/12 925 X X X X X X INVE SB-93 (9-10) 9-10 09/11/12 935 X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1150 X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X INVE		, ,										
SB-92 (14-15) 14-15 09/11/12 1110 X X X X X X X X X X X X X X X X X		, ,					X					
SB-92 (19-20) 19-20 09/11/12 1115 X X X X X X X INVE SB-92 (24-25) 24-25 09/11/12 1120 X X X X X X X INVE Refusal at 25 feet bgs in basalt. SB-93 SB-93 (0-1) 0-1 09/11/12 910 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 915 X X X X X X INVE SB-93 (4-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (9-10) 9-10 09/11/12 935 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X INVE Refusal at 22 feet bgs in basalt; moist at 22 feet bgs SB-94 SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X INVE		, ,										
SB-92 (24-25) 24-25 09/11/12 1120 X X X X X X INVE Refusal at 25 feet bgs in basalt. SB-93 SB-93 (0-1) 0-1 09/11/12 910 X X X X X X INVE SB-93 (1-2) 1-2 09/11/12 915 X X X X X INVE SB-93 (4-5) 4-5 09/11/12 925 X X X X X INVE SB-93 (9-10) 9-10 09/11/12 935 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X INVE SB-94 SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X INVE		, ,										
SB-93 SB-93 (0-1) 0-1 09/11/12 910 X X X X X X X X X		, ,										Defugal at 25 feet has in becalt
SB-93 (1-2) 1-2 09/11/12 915 X X X X X X X INVE SB-93 (4-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (9-10) 9-10 09/11/12 935 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X INVE SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE		SB-92 (24-25)	24-25	09/11/12	1120	^	^	^	^	^	IINVE	Refusal at 25 feet bys in basalt.
SB-93 (1-2) 1-2 09/11/12 915 X X X X X X X INVE SB-93 (4-5) 4-5 09/11/12 925 X X X X X X INVE SB-93 (9-10) 9-10 09/11/12 935 X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X X INVE SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE	SB-93	SB-93 (0-1)	0-1	09/11/12	910	X	X	X	X	X	INVF	
SB-93 (4-5)	32 33	, ,										
SB-93 (9-10) 9-10 09/11/12 935 X X X X X X X INVE SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X INVE Refusal at 22 feet bgs in basalt; moist at 22 feet bgs SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE		, ,										
SB-93 (14-15) 14-15 09/11/12 1010 X X X X X X X INVE SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X X INVE Refusal at 22 feet bgs in basalt; moist at 22 feet bgs SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X X X INVE		, ,										
SB-93 (19-20) 19-20 09/11/12 1020 X X X X X X INVE Refusal at 22 feet bgs in basalt; moist at 22 feet bgs SB-94 (0-1) 0-1 09/11/12 1150 X X X X X X INVE SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X X INVE												
SB-94 SB-94 (0-1)												Deficient at 20 feet has in benefit, maint at 20 feet has
SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X INVE		SB-93 (19-20)	19-20	09/11/12	1020	X	Χ	X	Χ	X	INVE	Refusal at 22 feet bgs in basait; moist at 22 feet bgs
SB-94 (1-2) 1-2 09/11/12 1155 X X X X X X INVE	SR-94	SR-94 (0-1)	0-1	00/11/12	1150	×	X	X	×	×	INI\/F	
	OD 94	, ,										
		, ,										
		, ,										
SB-94 (9-10) 9-10 09/11/12 1210 X X X X X INVE		, ,										Duralizate completed OR OA (O. 40)
SB-94D (9-10) 9-10 09/11/12 1215 X X X X X DUP Duplicate sample of SB-94 (9-10)		, ,										Duplicate sample of SB-94 (9-10)
SB-94 (14-15) 14-15 09/11/12 1220 X X X X X X INVE												B (1 400 (41 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
SB-94 (19-20) 19-20 09/11/12 1230 X X X X X X X INVE Refusal at 22 feet bgs in basalt; moist at 22 feet bgs.		SB-94 (19-20)	19-20	09/11/12	1230	Х	Х	Х	Х	Х	INVE	Refusal at 22 feet bgs in basalt; moist at 22 feet bgs.

Soil Sample Summary - 2012 Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Areas of Interest and	Areas of Interest and Sample Information						cal Parameters Rad Cl				
Soil Boring Phosphoric and Super Ph	Sample ID	<u>Depths</u>	<u>Date</u>	<u>Time</u>	Metals (a)	рH	General Chemistry ^(b)	Gross a & B		Sample Type (INVE/DUP)	Additional Comments
i nospilone and ouper i n	ospiloric Acid A	ii Ca									
South Car Wash Sump Ar	ea										
SB-95	SB-95 (0-1)	0-1	9/10/2012		X	Χ	X	X	X	INVE	
	SB-95 (1-2)	1-2	9/10/2012	1455	Χ	Χ	Χ	Х	Χ	INVE	Hand auger refusal at 3 feet bgs on gravel
SB-96	SB-96 (0-1)	0-1	9/10/2012	1350	Χ	Χ	Χ	X	Χ	INVE	
	SB-96D (0-1)	0-1	9/10/2012		X	Χ	X	X	X	DUP	Duplicate sample of SB-96 (0-1)
	SB-96 (1-2)	1-2	9/10/2012		X	Χ	X	X	X	INVE	
	SB-96D (1-2)	1-2	9/10/2012		X	Χ	X	X	X	DUP	Duplicate sample of SB-96 (1-2)
	SB-96 (4-5)	4-5	9/10/2012		Χ	Χ	X	Χ	Χ	INVE	Refusal at 18 feet bgs.
SB-97	CD 07 (0 1)	0.1	00/44/40	1615	V	V	V	V	V	INVE	
3 D -97	SB-97 (0-1)	0-1	09/11/12	1615 1620	X X	X	X	X	X		
	SB-97 (1-2)	1-2	09/11/12		X	X X	X X	X X	X X	INVE	Parabala abandanad at E fact has
	SB-97 (4-5)	4-5	09/11/12	1625	Χ	۸	*	٨	Χ	INVE	Borehole abandoned at 5 feet bgs.
SB-98	SB-98 (0-1)	0-1	09/11/12	1640	X	X	Χ	Χ	Χ	INVE	
	SB-98 (1-2)	1-2	09/11/12	1645	Χ	X	X	Χ	Χ	INVE	
	SB-98 (4-5)	4-5	09/11/12	1650	Χ	Χ	Χ	Χ	Χ	INVE	Borehole abandoned at 5 feet bgs.
SB-99	SB-99 (0-1)	0-1	09/11/12	1515	Х	Х	Χ	Χ	Х	INVE	
	SB-99 (1-2)	1-2	09/11/12	1520	X	X	Χ	Χ	Χ	INVE	
	SB-99 (4-5)	4-5	09/11/12	1530	X	Χ	Χ	Χ	Χ	INVE	
	SB-99 (8-9)	8-9	09/11/12	1535	Χ	Χ	X	Χ	Χ	INVE	Refusal at 9 feet bgs (not bedrock).
SB-100	SB-100 (0-1)	0-1	09/11/12	1550	Х	Х	Χ	Χ	Х	INVE	
	SB-100 (1-2)	1-2	09/11/12	1555	Χ	X	X	Χ	Χ	INVE	
	SB-100 (4-5)	4-5	09/11/12	1600	Χ	Χ	X	Χ	Χ	INVE	
	SB-100 (8-9)	8-9	09/11/12	1605	Χ	Χ	X	Χ	Χ	INVE	Refusal at 9 feet bgs (not bedrock).
SB-101	SB-101 (0-1)	0-1	09/11/12	1440	X	Х	Χ	Χ	Х	INVE	
	SB-101 (1-2)	1-2	09/11/12	1445	X	X	Χ	Χ	Χ	INVE	
	SB-101 (4-5)	4-5	09/11/12	1450	X	X	Χ	Χ	Χ	INVE	
	SB-101 (9-10)	9-10	09/11/12	1455	X	Χ	X	Χ	X	INVE	
	SB-101D (9-10)		09/11/12	1500	Χ	Χ	Χ	X	Χ	DUP	Duplicate sample of SB-101 (9-10) Refusal at 10 feet bgs (not bedrock).
Additional QC Samples											
Equipment Blanks	EB0911A12		9/11/2012		X	Χ	X	X	Χ	INVE	Collected from Geoprobe cutting shoe, deconned 9/10/2012
	EB0911B12	-	9/11/2012	1715	Χ	Χ	X	Χ	X	INVE	Collected from Geoprobe cutting shoe, deconned 9/11/2012

a/ Metals include list presented in Table 3.

b/ General chemistry parameters include pH, total phosphorus, nitrate, Total Kjeldahl Nitrogen, ammonia, and fluoride.

c/ Radiological analysis

Bold = Above criteria

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

		A Region 9 eening Levels ^(c)	ldaho Dep of Environmer														
Sample ID:		Protection of			Background		SB-88	SB-88	SB-88	SB-88	SB-89	SB-90	SB-90	SB-90	SB-90	SB-90	SB-91
Sampling Date:	Industrial	Groundwater ^(e)	Risk Based	Critical	Concentrations		09/11/12	09/11/12	09/11/12	09/11/12	09/10/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12
Depth:	Soil (mg/kg) I	Risk-Based/MCL-Bas	e IDTLs (mg/kg)	Pathway	(UTL 95-95) ^(f)	Values (g)	0-1	1-2	4-5	9-10	0-1	0-1	1-2	4-5	9-10	14-15	0-1
Metals (mg/kg)																	
Aluminum	99,000	23.000/NS	- (g)	NS	15,041	23,000	16.900	31,300	16.400	32,800	12,700	13.400	19.700	17,300	13,200	20.000	24,000
Antimony	41	0.27/0.27	4.8	GWP	0.50	0.50	4.7 J	5.9	2.6 J	1.9 J	2.6 J	1.8 U	-,	0.83 J	0.39 U	0.56 U	2.7 J
Arsenic	1.6	0.0013/0.29	0.39	Surficial Soil	4.7	4.7	8.7	7.6	5.6	4.4 J	6.8	5.7 J	8.4 J	3.3	4.9	7.8	4.0 J
Barium	19,000	1,200/82	896	GWP	170.2	170.2	124	140	115	225	139	111 J	99.1 J	91	74.4	125	143
Beryllium	200	13/3.2	1.63	GWP	0.89	1.63	0.8 J	1.3 J	1.5 J	0.62 U	0.74 J	1.8 J	2.2 U	2.9	4.0	0.99 J	0.54 U
Cadmium	80	NS/NS	1.4	GWP	0.869	1.4	49.4	8.0	39.8	96.5	106	43.4	121	41.2	10.2	0.28 U	24.1
Calcium	(j)	NS/NS	NS	NS	38,270 (I)	38,270	38,600	6,510	7,190	13,500	88,200	122,000	152,000	51,900	70,000	4,470	19,100
Chromium	15,0000	NS/180,000	2,135	GWP	18.61	2,135	311	468	529	392	184	371	930	165	67.3	26.8	374
Iron	72,000	270/NS	5.76	GWP	14,811	14,811	19,200	22,100	14,200	7,000	18,500	12,800	12,300	9,640	6,990	25,400	25,600
Lead	80	NS/14	50	GWP	13.59	14	135	101	15.4	16.4	19.1	7.3	6.6	8.0	8.3	14.9	36.9
Magnesium	(j)	NS/NS	NS	NS	7,146	'7,146	2,070 J	819 J	490 J	695 J	4,060	2,830 J	2,230 J	721 J	973 J	7,100	1,650 J
Manganese	2,300	NS/NS	223	GWP	742 (I)	742	172	36	43	37.7	326	379	155	258	181	404	337
Nickel	2,000	20/NS	59	GWP	15.71	20	9.4 J	1.7 J	2.9 J	2.1 J	22.3 J	10.7 J	36.6 J	15.1	12	30	5.3 J
Potassium	(j)	NS/NS	NS	NS	3,393 (I)	3,393	3,220 J	5,350	3,780 J	2,810 J	4,440 J	3,820 J	3,580 J	4,150	1,730 J	2,230 J	3,870 J
Selenium	510	0.4/0.26	2.0	GWP	1.04	1.04	2.1 U	1.7 J	2.1 U	2.5 U	2.5 U	4.0 J	10.3 J	1.4 J	0.78 U	1.1 J	2.2 U
Sodium	(j)	NS/NS	NS	NS	289 (I)	289	1,860 J	1,080 J	742 J	773 J	2,350 J	1,290 J	2,400 U	404 J	299 J	310 U	712 J
Thallium	1	0.0011/0.14	1.6	GWP	NS	0.5	2.9 J	13.1	1.3 U	1.6 U	3.7 J	2.3 U	5.7 U	0.78 U	0.51 U	0.73 U	1.4 U
Vanadium	520	78/NS	NS	NS	22.68	78	372	382	363	559	204	481	890	482	207	37	280
Wet Chemistry (mg/kg)	1																
Ammonia (as N)	NS	NS	4.15	Subsurface Soi	I NS	NS	402	504	196	112	342	412	332	98.1	46.3	49.1	697
Fluoride	41,000	NS	7.36	GWP	3.95	7	155	165	40.8	157	181	20.5	66.8	3.1 U	4.7 J	43.8	26.5
Nitrate as N	1.600.000	NS	18.4	GWP	NS	NS	1.2	5.6 J	3.0 U	3.3 U	1.1 J	2.7 U		3.2 J	3.9 J	18.2 J	4.4 J
Total Kieldahl Nitrogen	NS	NS	NS	NS	NS	NS	1.580	1,380	380	615	1.250	441	612	313	195	403	1.660
Total Phosphorus	NS	NS	NS	NS	NS	NS	49,200	52,200	47,200	20,800	67.100	3,020	90,300	139.000	66,000	10,000 U	62,400
рН	NS	NS	NS	NS	NS	NS	4.16	3.55	3.20	3.10	4.12	6.57	6.75	4.14	5.05	6.39	3.43
Dadialasiaal (n0:/a)																	
Radiological (pCi/g) Gross alpha	NS	NO	NS	NS	NS	NS	50.9	49.6	222	81	81	106	00	51	5.3	2.39	114
Gross alpha Gross beta	NS NS	NS NS	NS NS	NS NS	NS NS	NS NS	30.9	49.6 36.2	119	45.2	45.6	78	89 59	52.7	5.3 5.7	2.39	66
Radium-226	5 ^(surface) , 15	INO	INO	NS	NO		7.7	4.98	4.22	3.62	29.4	34.9	16.5	1.46	1.46	1.71	27.8
Naululii-220	5 (subsurface)					5 ^(surface) , 15 (subsurface)	1.1	4.98	4.22	3.02	29.4	34.9	6.01	1.40	1.40	1.71	Z1.8
		NS	NS	NS	NS												
Radium-228	15	NS	NS	NS	NS	15	0.9	1.19	1.67	0.95	0.69	0.44	1.1	1.11	0.7	1.25	0.62

U = Analyte not detected above method detection limit.

g/Comparative values are defined as:

J = Estimated result.

^{- =} Not analyzed.

NS = No regulatory criteria established.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per gram; "n" indicates RSL based on non-carcinogenic toxicity;

[&]quot;m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on carcinogenic toxicity; SS indicates surficial soil as the critical pathway;

GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gram; RL = reporting limit

IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C = degrees Celsius; TCL = target compound list;

[&]quot;NS" = not applicable or standard not developed; SSLs = soil screening levels

UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percentile of the distribution

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physical/Chemical Methods. As updated and revised.

SM source: American Public Health Association. 1998. Standard Methods for Examination of Water and Wastewater. As updated and revised.

EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Waste. EPA 600/4-70-020. As updated and revised.

EPA. 1980. EPA Prescribed Procedures for Measurement of Radioactivity in Drinking Water. EPA 600 4-80-032. As updated and revised

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-300). Environmental Measurements Laboratory. 28th Edition.

ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste with Water, will be used to prepare samples for analysis of fluoride.

c/ Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health/conmedia/soil/cleanup.htm

US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/region9/superfund/prg/

d/ Idaho Risk Reduction Manual, July 2004.

e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov/region9/superfund/prg/

f/ background as per EPA coresponance dated April 19, 2012, except where noted)

[•] the lower of the human health screening levels, unless this value is below background

[•] if the human health screening level is above background, then it is the CV

[•] if the human health screening level is below background, then background is the CV

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc.
Conda Phosphate Operations Facility Soda Springs, Idaho

h/ provided by Accutest Laboratories.

- i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for manganese.
- j/ To determine potential impacts from the releases, sample concentrations for these parameters will be compared to background concentrations.
 k/ Thallium was not detected in any of the background samples. The laboratory reporting limit was identified as the screening value as the human health values are lower than the reporting and method detection limits.
- I/ Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesium, potassium, and sodium calculated by WSP.
- m/ Radiochemistry labs do not use method detection limts and reporting limits that conventional labs use. Radiochemistry labs only use minimum detection concentrations as shown.

The EPA screening values provided for non-carcinogenic parameters (n) are 1/10th of the published screening levels to account for cumulative adverse effects.

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

	EF	PA Region 9	Idaho De	9																				
	Regional Sci	reening Levels ^(c)	of Environme	_																				
Sample ID:		Protection of		SB-91	SB-91	SB-91	SB-91	SB-92	SB-92	SB-92	SB-92	SB-92	SB-92	SB-92	SB-93	SB-93	SB-93	SB-93	SB-93	SB-93	SB-94	SB-94	SB-94	SB-94
Sampling Date:	Industrial	Groundwater (e)	Risk Based	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	e IDTLs (mg/kg)	1-2	4-5	9-10	14-15	0-1	1-2	4-5	9-10	14-15	19-20	24-25	0-1	1-2	4-5	9-10	14-15	19-20	0-1	1-2	4-5	9-10
Matala (ma/ka)																								
Metals (mg/kg) Aluminum	99,000	23,000/NS	- (g)	22,300	22,700	9,680	19,000	11,100	21,700	14,600	11,800	6,530	19,000	17,700	11,000	4,630	17,300	8,770	5,500	23,700	13,400	26,800	19,100	12,600
Antimony	41	0.27/0.27	- (g) 4.8	3.2 U	22,700 2.6 J	9,000 1.0 U	0.38 U	3.8 J	21,700 2.8 J	0.55 U	0.48 U	0.34 U	19,000 I 0.37 L	J 0.57 L	J 1.7 J	4,630 2.7 L		0,770 0.42 U	0.34 U		4.0 J	20,600 2.3 J	0.79 J	J 0.44
Arsenic	1.6	0.0013/0.29	0.39	9.7 J	5.5 J	2.6 J	6.9	5.3	4.2	12	9.1	2.9	5.3	6.3	3.6 J	2.7 L		3.1	4.3	5.8	6.6	4.0	7.1	8.3
Barium	19,000	1,200/82	896	77.9 J	156 J	135	99.5	125	87.6	117	84.8	42.7	144	96.1	85.7 J	83.7 J		72.5	40.1	132	109	4.0 77	125	93.6
Bervllium	200	13/3.2	1.63	1.6 U		1.8 J	0.89 J	0.54 J	0.67 J	1.0 J	0.84 J	42.7 0.47 J	1.2	0.9 J	0.55 U	1.3 L		72.3 0.49 J	40.1 0.4 J	1.2	0.55 J	0.45 J	3.1	0.73
Cadmium	80	NS/NS	1.4	81.8	34.8	14.3	0.03 J	13	20.5	13.1	3.9	2.7	0.41 J	0.28 L	J 20.5	1.5 J	35.1	2.2	5.4	0.23 U	23	5.0	45.2	1.5
Calcium	(i)	NS/NS	NS	71,700	19,900	77,700	6,030	38,500	69,200	70,600	99,400	41,000	4,770	3,960	69,400	872 J	42,900	53,500	86,200	4,170	28,100	59,700	66,900	60,200
Chromium	15,0000	NS/180,000	2,135	658	557	130	27.4	335	319	24.9	22.9	19.1	20.4	16.9	236	100	163	21.8	26.7	21.5	367	352	47.1	27
Iron	72,000	270/NS	5.76	13,100	17,000	3,150	20,500	25,300	16,600	2,410	5,690	6,330	15,600	26,700	7,300	6,610	10,700	7,500	4,750	17,800	32,600	19,300	10,300	11,900
Lead	80	NS/14	50	13.7	12.5	8.3	14.4	29.4	16.9	8.1	6.9	8.7	13.5	8.9	24.9	3.1 J	8.2	7,500	7.1	15.2	54.4	11.4	9.2	11.0
Magnesium	(i)	NS/NS	NS	2,070 J	779 J	410 J	6,060	388 J	567 J	617 J	2730	9,450	5,860	6,750	496 J	130 L	J 1,230 J	9.470	19,300	6,330	385 J	511 J	1160 J	J 8,820
Manganese	2,300	NS/NS	223	196	200	100	397	106	40.9	200	394	170	499	342	28.6	4.0 J	216	158	271	235	51.3	29.6	197	550
Nickel	2,000	20/NS	59	1.7 J	9.9 J	4.3 J	32.8	2.9 J	1.7 J	100	65	19.1	42.6	37.3	1.7 J	1.3 J	11.6 J	29.5	22.1	31.7	2.8 J	1.1 J	15.9	38.2
Potassium	(i)	NS/NS	NS	3.510 J	3,690 J	2,760 J	2,240	4,080	3,290 J	1.800 J	1,570 J	1,000 J	2,030	2,350 J	1,490 J	1,940 J	2,620 J	1,310 J	953 J	2,230 J	5,830	2,990	2,450 J	J 1,820
Selenium	510	0.4/0.26	2.0	6.6 J	4.5 U	2.1 U	0.8 J	3.0 J	2.1 J	1.1 U	0.95 U	0.69 U	,	,	,	5.3 L	,	,	0.69 U	,	2.6 J	2.2 J	1.1 U	
Sodium	(i)	NS/NS	NS	1,800 U	1,200 U	570 U	225 J	262 J	360 U			272 J	200 L			1500 L		230 U	343 J	260 U	230 U	290 U		
Thallium	1	0.0011/0.14	1.6	4.2 U	2.9 U	13 U	0.49 U	7.3	3.3	0.71 U	0.62 U	0.45 U	0.48 L	J 0.74 L		3.5 L	1.0 U	0.55 U	0.45 U		11.6	4.8	0.71 U	
Vanadium	520	78/NS	NS	681	540	308	45.4	260	317	113	24.4	22.6	25.8	28.9	158	70.7	349	24.9	29.5	29.9	414	287	547	33.5
Wet Chemistry (mg/kg	-																							
Ammonia (as N)	NS	NS	4.15	301	78.8	74.3	45.5	467	340	70	61	47.1	104	69.1	168	150	75.9	46	34	94	810	503	68	144
Fluoride	41,000	NS	7.36	123	6.2 J	3.7 J	26.6 J	332	67	3 U		2.8 U		24.1	969	48.2	14.6	3.7 J	3.4 J	3 U	186	151	5.9	5.2
Nitrate as N	1,600,000	NS	18.4	4.2 J	3.3 J	3.5 J	16.5 J	3.4 U				2.8 U			J 3.4 U	2.7 L	J 5.1 J	2.9 U	2.8 U		3.2 U	3.6 U		3.5
Total Kjeldahl Nitrogen	NS	NS	NS	712	886	271	442	1,130	566	1,440	491	360	442	237	654	389	226	509	345	365	2,680	1,910	607	601
Total Phosphorus	NS	NS	NS	60,000	66,000	102,000	10,700	38,500	60,900	60,900	50,400	19,200	7,490	39,500	59,100	55,100	46,800	7,930	25,800	8,640	48,000	35,100	70,300	31,000
рН	NS	NS	NS	4.15	3.91	4.29	6.32	2.85	4.98	4.98	5.37	7.24	6.46	6.66	2.79	7.43	4.80	7.03	7.19	6.62	4.31	3.29	4.80	6.52
Radiological (pCi/g)																								
Gross alpha	NS	NS	NS	63	78	48	3.76	79	86	35	56.5	9.9	3.52	3.32	121	67	91	5.6	9.1	2.69	80	45	41.9	6.3
Gross beta	NS	NS	NS	60	49.7	58	3.07	67	60	38.8	36.6	8.3	3.51	2.85	76	48.6	37.4	3.6	7.2	3.09	61	30.4	46.6	4.3
Radium-226	5 ^(surface) , 15		110	1.64	3.93	1.8	1.24	24.1	8.1	1.44	15.9	2.79	1.25	1.13	24.6	29.3	2.21	1.92	3.16	1.25	29.2	21.9	1.16	2.23
radidili 220	o , lo (subsurface)			1.04	0.00	1.0	1.27	۷ ۱	0.1	1	10.9	2.13	1.20	1.13	24.0	20.0	2.21	1.52	5.10	1.20	25.2	21.9	1.10	2.20
	(NS	NS																					
Radium-228	15	NS	NS	1.14	0.99	0.91	0.95	0.29	1.61	0.79	0.88	0.81	1.36	0.64	1.13	0.43	2.05	0.81	0.71	1.17	0.71	0.66	0.93	1.25

U = Analyte not detected above method detection limit.

J = Estimated result.

^{- =} Not analyzed.

NS = No regulatory criteria established.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per g
"m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on ca
GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gra
IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C =
"NS" = not applicable or standard not developed; SSLs = soil screening lev
UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percenti

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physics SM source: American Public Health Association. 1998. Standard Methods for EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Wast EPA. 1980. EPA Prescribed Procedures for Measurement of F

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-30 ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste c/ Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening Levels (revised November 2010), http://www.epa.gov/superfund/health US EPA Regional Screening (revised November 2010), https://www.epa.gov/superfund/health US EPA Regional Screening (revised November 2010), http

d/ Idaho Risk Reduction Manual, July 2004.

e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov f/ background as per EPA coresponance dated April 19, 2012, except where not g/Comparative values are defined as:

[•] the lower of the human health screening levels, unless this value is belo

[•] if the human health screening level is above background, then it is the (

[•] if the human health screening level is below background, then background

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc.
Conda Phosphate Operations Facility Soda Springs, Idaho

- h/ provided by Accutest Laboratories.
 i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for n
- j/ To determine potential impacts from the releases, sample concentrations for th k/ Thallium was not detected in any of the background samples. The laboratory
- // Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesiur m/ Radiochemistry labs do not use method detection limts and reporting limits the

The EPA screening values provided for non-carcinogenic parameters (n) are

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

		A Region 9 eening Levels ^(c)	Idaho De <u>of Environme</u>																				
Sample ID:		Protection of		SB-94	SB-94	SB-95	SB-95	SB-96	SB-96	SB-96	SB-97	SB-97	SB-97	SB-98	SB-98	SB-98	SB-99	SB-99	SB-99	SB-99	SB-100	SB-100	SB-100
Sampling Date:	Industrial	Groundwater (e)	Risk Based	09/11/12	09/11/12	09/10/12	09/10/12	09/10/12	09/10/12	09/10/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12
Depth:	Soil (mg/kg) I	Risk-Based/MCL-Base	e IDTLs (mg/kg)	14-15	19-20	0-1	1-2	0-1	1-2	4-5	0-1	1-2	4-5	0-1	1-2	4-5	0-1	1-2	4-5	8-9	0-1	1-2	4-5
B. (. 1 . / / / /																							
Metals (mg/kg)	00.000	22.000/NC	(~)	0.07	40.400	7.050	0.750	0.500	0.400	0.000	20.700	05.000	44.000	44.000	40.000	00.000	40.400	47.000	04.000	44.000	45.000	44.000	00.400
Aluminum	99,000 41	23,000/NS 0.27/0.27	- (g) 4.8	6,97	,	7,950 1.7 J	8,750 1.9 J	9,530	3,460	3,820 4.1	39,700 5.6 J	35,600 2.2 J	14,000	14,300 1.8 J	16,800 0.6 U	20,600 3.9 J	16,400	17,000	24,300 2.2 J	14,000 1.4 U	15,000 0.9 J	14,800 3.5 J	23,100
Antimony	1.6	0.0013/0.29	0.39	U 0.4		7.4	8.2	4.0 2.7	1.5 J 2.7	2.3	7.3	2.2 J 4.4	1.0 J 6.8	3.8 J	3.4	9.5	3.9 J 7.6	4.1 J 5.5	3.4	6.8	3.9 J	3.5 3	2.5 U 7.3 J
Arsenic	19.000	1.200/82	896			7. 4 77.2 J	0.2 115	163	35.3	42.2	302	321	274	3.6 J 194	220	9.5 359	248	399		90.8 J	210	3.7 140	7.3 J 249 J
Barium	200	1,200/82	1.63	J 0.4		0.83 J	0.96 J	0.38 J	ან.ა 0.16 U	42.2 0.19 U	0.84 J	0.65 J	0.43 J	0.42 U	0.57 J	0.89 J		0.64 J	233 0.52 J	90.8 J 1.1 J	0.89 J	0.46 J	249 J 1.6 J
Beryllium	80	NS/NS	1.4	J 0.4		34.2						0.65 J 0.49 J		98.4	15		0.5 J 42.9	7.0	3.8	58.7	47.8	93.7	
Cadmium	(i)	NS/NS	NS	18,70		28,200	39.1 31.400	15.7 2,570	1.0 413 J	0.34 J 750 J	3.8 696 J	3,980	102 20,200	9,390	37,000	2.2 949 J	42.9 14,500	133,000		78,900	47.8 64,100	16,000	52 176,000
Calcium	15,0000	NS/180,000	2,135	34.	,	176	218	156	70.7	58.2	570	3,960	20,200	196	292	949 J 479	278	1,670	499 J 253	306	758	224	526
Chromium	72,000	270/NS	2,135 5.76	9,22		8,850	10,900	14,300	4.470	4,640	26,900	21,800	31,200	26,000	36,300	33,800	38,900	76,100	12,200	15,000	31,000	14,200	28,000
Iron	80	NS/14	50	9,22		28	38.4	11.4	4,470	4,640 5.7	13.8	8.7	9.1	5.5	7.2	18.3	8.8	10.9	10.1	5.0	12.0	6.5	25.7
Lead	(i)	NS/NS	NS	6,34		26 1,240 J	1,310 J	647 J	4.6 108 J	107 J	368 J	6.7 454 J	9.1 949 J	898 J	8,610	236 J	897 J	15,000	294 J	5,460	5,250	492 J	12,700
Magnesium	2,300	NS/NS	223	21	,	1,240 3	1,310 3	39.9	2.9	2.9	32	434 3	470	359	1,830	66	497	15,900	13.9	590	3,230	83	3,400
Manganese Nickel	2,000	20/NS	59	27.		13.5 J	17.4	9.4	2.9 0.87 J	2.9 0.48 J	2.6 J	43 1.2 J	8.5 J	4.7 J	1,030	2.6 J	7.4 J	62.1	4.6 J	74.6	3,330 13.9 J	8.2 J	3,400 44.3 J
Potassium	2,000 (i)	NS/NS	NS	J 87		1,300 J	1,380 J	2,940	1.370 J	1,250 J	8,830	7,000	4,930	3.850 J	2,750 J	11,400	5.840	772 J	5,850	2.100 J	3,260 J	4,970	6,770 J
Selenium	510	0.4/0.26	2.0	U 1.	,	1,300 J	,	,	0.65 U	,	1.2 J	1.9 J	4,930 1.6 U	-,	2,730 J	2.3 J	2.3 J	4.7 J	,	2,100 J 2.7 U	3,200 J 1.8 U	4,970 2.2 J	4.9 U
Sodium	(i)	NS/NS	NS	J 27		1,210 J	1,330 J	482 J	180 U	210 U	310 U	509 J	2,500 J	1,870 J	3.470	2.3 J	1,760 J	4.7 J	290 U	750 U	1,060 J	2.2 J 240 U	4,500 J
Thallium	(I) 1	0.0011/0.14	1.6	U 0.6		1,210 J	1,330 J 1.2 J	4.9		1.7 J	3.6	1.8 J	2,300 J	,	3,470 1.4 J	2.8	1,700 3	477 J	1.2 J	1.8 U	1,000 J	7.5	4,300 J 16 U
Vanadium	520	78/NS	NS	22.		154	1.2 3	186	2.1 52.1	26.8	542	476	448	257	282	415	550	265	415	483	367	7.3 598	554
variaulum	320	70/110	140	22.	23	134	170	100	32.1	20.0	342	470	440	251	202	413	330	203	413	403	307	390	334
Wet Chemistry (mg/kg))																						
Ammonia (as N)	NS	NS	4.15	63.	3 29	25	23.6	690	120	139	338	503	478	480	132	626	917	13	478	87	498	818	209
Fluoride	41,000	NS	7.36	J 4.		29.3	31.3	1960	236	35.7	20	21.9	21.1	28	5.3 J	81.6	12.3	27.3	6.1 J	3.8 J	13.6	2.6 U	
Nitrate as N	1,600,000	NS	18.4	J 3.		4.4	5.3	0.57 J	1.3 J	26.5	6.1 J	3.9 J	3.2 U	J 3.1 U	3.1 J	5.6 J	3 U	13.6	5.5 J	4.1 J	12.2	20.5	42.6
Total Kjeldahl Nitrogen	NS	NS	NS	23	347	225	244	1210	889	268	1,400	907	1,330	1,400	271	2,670	4,600	132	955	343	1,270	1,760	684
Total Phosphorus	NS	NS	NS	6,02	14,200	13,000	10,500	21,800	4,220	6,070	57,700	37,800	272,000	51,100	24,200	64,700	93,200	130,000	49,200	132,000	74,200	54,500	79,600
pН	NS	NS	NS	7.1	6.57	6.44	6.52	4.04	4.03	4.43	2.78	2.76	3.38	3.33	5.06	2.42	3.63	7.70	3.56	5.59	6.20	4.45	6.19
Radiological (pCi/g)																							
Gross alpha	NS	NS	NS	6.		72	71	3.97	2.45	17.5	29.4	63	69	56	31.7	32	64	107	31.8	4.8	57	54	54
Gross beta	NS	NS	NS		1 2.27	47.8	43.8	2.2	1.65	14.7	23.3	42	44.9	43.8	26	27.1	54.7	56	18.3	5	43.4	37.4	42.8
Radium-226	5 ^(surface) , 15			2.3	9 1.37	11.9	12.5	1.23	0.89	9.2	1.92		3.01	2.25	5.2	2.34	13.8	8.4	1.85	0.41	3.83	7.6	12.2
	(subsurface)																						
		NS	NS									8.2											
Radium-228	15	NS	NS	1.1	2 1.15	0.71	0.49	0.76	0.71	0.52	0.84	0.3	0.03	0.25	0.43	1.08	0.34	1.05	0.93	0.54	0.55	0.47	0.53

U = Analyte not detected above method detection limit.

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-30 ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste c/ Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/healtt US EPA Regional Screening Levels (revised November 2010), http://www.epid/ Idaho Risk Reduction Manual, July 2004.

e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov f/ background as per EPA coresponance dated April 19, 2012, except where not g/Comparative values are defined as:

- the lower of the human health screening levels, unless this value is belo
- if the human health screening level is above background, then it is the (
- if the human health screening level is below background, then background

J = Estimated result.

^{- =} Not analyzed.

NS = No regulatory criteria established.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per g
"m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on ca
GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gra
IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C =
"NS" = not applicable or standard not developed; SSLs = soil screening lev
UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percenti

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physics SM source: American Public Health Association. 1998. Standard Methods for EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Wast EPA. 1980. EPA Prescribed Procedures for Measurement of F

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc.
Conda Phosphate Operations Facility Soda Springs, Idaho

- h/ provided by Accutest Laboratories.
 i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for n
- j/ To determine potential impacts from the releases, sample concentrations for th k/ Thallium was not detected in any of the background samples. The laboratory
- // Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesiur m/ Radiochemistry labs do not use method detection limts and reporting limits the

The EPA screening values provided for non-carcinogenic parameters (n) are

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

		PA Region 9 reening Levels ^(c)	Idaho De of Environme					
Sample ID:		Protection of		SB-100	SB-101	SB-101	SB-101	SB-101
Sampling Date:	Industrial	Groundwater (e)	Risk Based	09/11/12	09/11/12	09/11/12	09/11/12	09/11/12
Depth:	Soil (mg/kg)	Risk-Based/MCL-Base	IDTLs (mg/kg)	8-9	0-1	1-2	4-5	9-10
Metals (mg/kg)								
Aluminum	99,000	23,000/NS	- (g)	30.600	15.700	14.100	9,900	17,600
Antimony	41	0.27/0.27	4.8	1.3 U	0.96 U	1.9 J	0.47 U	1.7 U
Arsenic	1.6	0.0013/0.29	0.39	5.8 J	2.4 J	5.0 J	9.5	2.3 J
Barium	19.000	1.200/82	896	121 J	187	155	67.7	117 J
Beryllium	200	13/3.2	1.63	4.1	0.73 J	1.2 J	1.4	0.91 J
Cadmium	80	NS/NS	1.4	22.1	29.3	41.9	7.5	35.9
Calcium	(j)	NS/NS	NS	17,300	21,200	92,100	124,000	43,900
Chromium	15,0000	NS/180,000	2,135	283	207	218	29.6	204
Iron	72,000	270/NS	5.76	12,400	34,100	14,800	14,900	9,350
Lead	80	NS/14	50	11.9	5.8	19.9	5.6	7.9
Magnesium	(i)	NS/NS	NS	814 J	5,290	4,800	2,940	407 J
Manganese	2,300	NS/NS	223	111	580	323	598	82
Nickel	2,000	20/NS	59	69.3	4.7 J	19.2 J	114	4.6 J
Potassium	(j)	NS/NS	NS	6,480 J	2,560 J	6,260 J	1,280 J	2,330 J
Selenium	510	0.4/0.26	2.0	2.6 U	1.9 U	5.4 J	0.93 U	3.4 U
Sodium	(j)	NS/NS	NS	720 U	2,920 J	3,690 J	341 J	950 U
Thallium	1	0.0011/0.14	1.6	17 U	24 U	1.9 U	0.61 U	2.2 U
Vanadium	520	78/NS	NS	524	262	282	100	454
Wet Chemistry (mg/kg)								
Ammonia (as N)	NS	NS	4.15	496	130	180	103	151
Fluoride	41,000	NS	7.36	10.8	14.4	76.3	2.8 U	3.7 J
Nitrate as N	1,600,000	NS	18.4	9.0	2.9 J	2.8 J	5.2 J	4.4 J
Total Kjeldahl Nitrogen	NS	NS	NS	3,890	912	241	525	133
Total Phosphorus	NS	NS	NS	242,000	61,900	46,300	111,000	154,000
рН	NS	NS	NS	4.48	3.74	5.06	5.43	4.48
Radiological (pCi/g)								
Gross alpha	NS	NS	NS	68	53	53.4	55	40.4
Gross beta	NS	NS	NS	62	38.7	44.2	38.4	24.6
Radium-226	5 ^(surface) , 15			3.54	2.94	4.29	2.29	4.35
	(subsurface)							
		NS	NS					
Radium-228	15	NS	NS	1.53	0.44	1.29	1.32	1.18

U = Analyte not detected above method detection limit.

HASL source: U.S. Department of Energy. EML Procedures Manual (HASL-30 ASTM source: American Society for Testing and Materials.

Methods for sample preparation include SW-846 3050B for metals.

ASTM D3987-85, Standard Test Method for Shake Extraction of Solid Waste c/ Soil Cleanup Criteria in 40 CFR Part 192, http://www.epa.gov/superfund/healtt US EPA Regional Screening Levels (revised November 2010), http://www.epid/ Idaho Risk Reduction Manual, July 2004.

e/ US EPA Regional Screening Levels (revised May 2012), http://www.epa.gov f/ background as per EPA coresponance dated April 19, 2012, except where not g/Comparative values are defined as:

- the lower of the human health screening levels, unless this value is belo
- if the human health screening level is above background, then it is the (
- if the human health screening level is below background, then backgrou

J = Estimated result.

^{- =} Not analyzed.

NS = No regulatory criteria established.

a/ mg/kg = milligrams per kilogram; s.u. = standard units; pCi/g = picocuries per g
"m" indicates RSL may exceed the ceiling limit; "c" indicates RSL based on ca
GWP indicates groundwater is the critical pathway; pci/g = picoCuries per gra
IDTL = Idaho default target level; G = glass; Poly = polyethylene; °C =
"NS" = not applicable or standard not developed; SSLs = soil screening lev
UTL 95-95 = upper tolerance limits 95% confidence limit of the 95 th percenti

b/ SW-846 source: EPA. 1986. Test Methods for Evaluating Solid Waste, Physics SM source: American Public Health Association. 1998. Standard Methods for EPA sources: EPA. 1983. Methods for Chemical Analysis of Water and Wast EPA. 1980. EPA Prescribed Procedures for Measurement of F

Soil Sampling Analytical Results - 2012 North and South Car Wash Sump Areas Nu-West Industries, Inc.
Conda Phosphate Operations Facility Soda Springs, Idaho

- h/ provided by Accutest Laboratories.
 i/ The RSL for diet is reported for cadmium; the RSL for non-diet is reported for n
- j/ To determine potential impacts from the releases, sample concentrations for th k/ Thallium was not detected in any of the background samples. The laboratory
- // Values not provided by EPA; UTL 95-95 concentrations for calcium, magnesiur m/ Radiochemistry labs do not use method detection limts and reporting limits the

The EPA screening values provided for non-carcinogenic parameters (n) are

Soil Sampling Analytical Results - 2012 Equipment Blank Nu-West Industries, Inc. Conda Phosphate Operations Facility Soda Springs, Idaho

Sample ID: Sample Type: Sampling Date:	EB0911A12 INVE 9/11/2012	EB0911B12 INVE 9/11/2012
Depth (feet):	<u>-</u>	
Metals (mg/l)		
Aluminum	0.025 U	0.025 U
Antimony	0.0039 J	0.020 U
Arsenic	0.0020 U	0.0020 U
Barium	0.0050 U	0.0050 U
Beryllium	0.0010 U	0.0010 U
Cadmium	0.0010 U	0.0010 U
Calcium	0.10 U	0.10 U
Chromium	0.0010 U	0.0010 U
Iron	0.035 U	0.035 U
Lead	0.0010 U	0.0010 U
Magnesium	0.10 U	0.10 U
Manganese	0.0010 U	0.0010 U
Nickel	0.0020 U	0.0020 U
Potassium	0.50 U	0.50 U
Selenium	0.0020 U	0.0020 U
Sodium	1.9 U	1.9 U
Thallium	0.0019 U	0.0019 U
Vanadium	0.0010 U	0.0010 U
Wet Chemistry (mg/l)		
Ammonia (as N)	0.050 U	0.050 U
Fluoride	0.050 U	0.050 U
Nitrate as N	0.090 J	0.089 J
Total Kjeldahl Nitrogen	0.11 U	0.11 U
Total Phosphorus	0.028 U	0.093 J
рН	6.73	6.60
Radiological (pCi/l)		
Gross alpha	-	_
Gross beta	-	-
Radium 226	-	-
Radium 228	-	-

U = Analyte not detected above method detection limit.

INVE = Investigative sample.

WSP Environment & Energy

1 of 1

J = Estimated result.

^{- =} Not analyzed.